


Small Molecule Screen Reveals Regulation of Survival Motor Neuron Protein Abundance by Ras Proteins

Reka R. Letso,[†] Andras J. Bauer,[†] Mitchell R. Lunn,^{†,§} Wan Seok Yang,[†] and Brent R. Stockwell^{*,†,‡}

[†]Howard Hughes Medical Institute, Department of Biological Sciences and [‡]Department of Chemistry, Columbia University, Northwest Corner Building, MC 4846, 550 West 120th Street, New York, New York 10027, United States

Supporting Information

ABSTRACT: Small molecule modulators of protein activity have proven invaluable in the study of protein function and regulation. While inhibitors of protein activity are relatively common, small molecules that can increase protein abundance are rare. Small molecule protein upregulators with targeted activities would be of value in the study of the mechanisms underlying loss-of-function diseases. We developed a high-throughput screening approach to identify small molecule upregulators of the Survival of Motor Neuron protein (SMN), whose decreased levels cause the neurodegenerative disease spinal muscular atrophy (SMA). We screened 69,189 compounds for SMN upregulators and performed mechanistic studies on the most active compound, a bromobenzophenone analogue designated cuspin-1. Mechanistic studies of cuspin-1 revealed that increasing Ras signaling upregulates SMN protein abundance via an increase in translation rate. These findings suggest that controlled modulation of the Ras signaling pathway may benefit patients with SMA.



Bioactive small molecules have enabled major advances in biological research and drug development. Although many efforts have focused on the identification of small molecule inhibitors of protein function, the ability to upregulate protein abundance or otherwise increase protein activity can provide complementary insights that cannot be obtained through loss of function studies alone.¹ Although the number of compounds known to function as protein upregulators is small, they have important research and therapeutic applications. Examples include histone deacetylase inhibitors, used in the treatment of psychiatric conditions, neurologic disorders, and cutaneous T cell lymphoma,^{2–4} and proteasome inhibitors, used in the treatment of multiple myeloma⁵ and for the study of the ubiquitin-proteasome system and its targets.⁶

Unfortunately, existing upregulating compounds lack specificity, altering the levels of hundreds to thousands of gene products.⁷ Such broad spectrum effects, in addition to potentially being therapeutically undesirable, make these compounds unsuitable for investigations of mechanisms regulating the abundance of an individual or limited number of proteins. Small molecules that can upregulate specific protein targets are needed, and would be especially advantageous for the study of loss of function diseases. However, we lack systematic methods for identifying such compounds.

In order to address this need, we developed a high-throughput screening approach to identify small molecule upregulators of a specific protein. This assay detects increases in the abundance of a target protein after treatment with small molecules. We selected the Survival Motor Neuron (SMN) protein as a target for modulation. Decreased SMN protein

levels result in the loss of function neurodegenerative disease spinal muscular atrophy (SMA). In SMA patients, SMN protein levels are decreased to ~10% of normal levels, due to homozygous deletion of the *survival of motor neuron 1* (*smn1*) gene but retention of the *smn2* gene. The *smn2* gene is nearly identical to *smn1* but contains a translationally silent C to T transition that causes the *smn2* pre-mRNA to be aberrantly spliced, generating only ~10% of the full-length, functional protein produced by the *smn1* gene.⁸ This small amount of SMN protein prevents the embryonic lethality seen in an SMN null background⁹ but cannot prevent the motor neuron degeneration observed in SMA. Small molecule upregulators of SMN could provide novel insights into the regulation of SMN protein levels, leading the way to new therapeutic targets. A relevant example is the screen for SMN-upregulating compounds performed by Makhortova et al, which identified the RTK-PI3K-AKT-GSK-3 signaling cascade as an important modulator of SMN protein stability.¹⁰

We developed an assay for detecting increased SMN protein abundance and screened 69,189 small molecules. We discovered three SMN-upregulating compounds. Mechanistic investigations of the most effective compound, Chemical Upregulator of SMN Protein-1 (cuspin-1), revealed that increasing Ras signaling upregulates SMN protein levels by enhancing the rate of SMN translation.

Received: July 19, 2012

Accepted: March 15, 2013




Figure 1. High-throughput screen (HTS) and hit compound cuspin-1. (A) Schematic of HTS assay for SMN protein upregulators. (B) Structure of cuspin-1, the optimal SMN-upregulating small molecule discovered in our screen. (C) Cytoblot dose-response curve for cuspin-1 in 9677 SMA patient fibroblast cells treated for 48 h. The data represent the average of triplicates \pm SEM. RLU: relative luminescence units. *p*-values were calculated using a two-tailed *t* test (* = *p* < 0.01, ** = *p* < 0.001). (D) Western blot analysis of SMN and actin in 9677 cells treated with indicated concentrations of cuspin-1 for 48 h. Fold increase of SMN to actin ratio normalized to nontreated control (NT) is graphed below. Data represent average of duplicates \pm standard deviation.

RESULTS AND DISCUSSION

Development of a High-Throughput Screen That Detects Endogenous SMN Protein Abundance. In order to identify compounds upregulating SMN protein levels, we developed a high-throughput screen for monitoring the levels of endogenous SMN protein in SMA patient fibroblast cells. Previous screens searching for SMN-upregulating compounds focused on altering either the transcription or splicing of *smn2*.^{11,12} Our screen focuses on the desired final product, SMN protein abundance, in order to be inclusive of all aspects of SMN protein regulation, from transcription through degradation. This method permits identification of small molecule upregulators of SMN protein without bias toward their mechanism of action.

To quantify SMN protein levels in a 384-well format, the cyto blot assay,¹³ a modified enzyme-linked immunosorbent assay (ELISA), was optimized for microtiter-based screening using SMA patient fibroblasts. A schematic of the primary screening workflow is shown in Figure 1A. The optimized assay protocol involved the seeding of cells and subsequent treatment with small molecules for 48 h. This incubation period was selected to allow sufficient time for the compounds to increase SMN protein levels, while limiting the depletion of nutrients and accumulation of toxins in the microtiter format during prolonged incubation.¹⁴ After incubation, cells were fixed,

permeabilized, and incubated with primary and secondary antibodies in-plate. Horseradish-peroxidase-derived chemiluminescence was detected using an automated plate reader. The optimized SMN cyto blot is a suitably robust format for detecting SMN levels in cells, as indicated by a *Z'* factor¹⁵ of 0.35–0.5 in the optimized assay, using the 3814 SMA carrier cell line as a positive control.

Small Molecule Libraries and *in Silico* Filtering for Blood–Brain Barrier Penetrating Small Molecule Libraries. We screened compounds from two libraries that we assembled in our laboratory. The first was our previously reported small molecule library,¹⁶ described in Methods. The second library was composed of compounds specifically chosen for their predicted ability to cross the blood–brain barrier (BBB), as the development of CNS-active compounds is impeded by the inability of a large portion of compounds to penetrate the BBB. We addressed both BBB-penetration and toxicity issues during the selection of small molecules for our library by *in silico* filtering for compounds likely to be BBB-penetrant and filtering out known toxic and reactive species, thus increasing the ultimate therapeutic potential of this “BBB library”.

To predict the BBB penetration potential of each compound, we utilized the logarithm of brain to blood partitioning ratio (log BB). We used a training set of 109 compounds with experimentally determined log BB values¹⁷ in creating a log BB




Figure 2. Structure–activity relationship of cuspin-1. (A) Structures and activities of the highest similarity commercially sourced cuspin-1 analogues. (B) Friedel–Crafts acylation scheme for synthesis of cuspin-1. The same synthetic scheme was utilized for synthesis of cuspin-1 analogues. (C) Structures and activities of analogues synthesized.

predictor based on simple physicochemical descriptors, such as log P (logarithm of the octanol–water partitioning coefficient of a compound). The compound set was split into two sets, one with 73 compounds for creating a predictor and one with 36 compounds for validation. Partial least-squares regression was used to correlate experimental log BB values of the training set with physicochemical descriptors. Our choice of descriptors was influenced by the planned use of the log BB predictor, as we needed it to be able to evaluate millions of candidate compounds with relatively little computational expense. For this reason, we avoided the use of semiempirical energy terms and instead chose to use descriptors that were calculated using the atomic contribution model. This approach, pioneered by Wildman and Crippen,¹⁸ uses a statistical data set of known atoms and functional groups to predict the groups' contribution to molecular descriptors. The descriptors in the model were

made up of the number of rotatable bonds, van der Waals surface area, molar refractivity, number of nitrogen atoms, and log P. To enable better sampling of these descriptors, we calculated log P and van der Waals descriptors multiple times with different weights (Supplementary Table S1). The obtained model gave $R^2 = 0.82$ for the training set and $R^2 = 0.70$ for the validation set. The predictions gave no false negatives (Supplementary Figure 1).

Screening and Hit Validation. A total of 69,189 small molecules were screened in the SMN cyto blot assay. Primary screening was performed in triplicate in 9677 SMA patient fibroblast cells in 384-well format. To remain inclusive of compounds with moderate efficacy, a “hit” was defined as having a median signal greater than 2 standard deviations from the median of the untreated wells. The 1,106 compounds passing hit criteria in the primary screen were retested in dose–

response series in three SMA patient fibroblast lines (232, 3813, and 9677). The 105 candidates showing dose–response activity in at least two cell lines were tested by Western blot. Three structurally distinct compounds were found to significantly increase endogenous SMN protein levels (Figure 1B,C and Supplementary Figure 2). The EC_{50} values for the three scaffolds were in the range of 10–20 μ M, typical for hits from high-throughput, cell-based screens. A summary of the screen can be found in Supplementary Table S2.

We focused on the most robust of these three hits, a bromobenzophenone analogue we designated cuspin-1 (Chemical Upregulator of SMN Protein-1, Figure 1B), whose identity was confirmed by proton NMR (Supplementary Figure 3). Cuspin-1 showed strong activity in the plate-based dose–response assay (Figure 1C) and increased SMN levels by 50% compared to DMSO-treated controls at 5 μ g/mL (18 μ M) in SMA patient fibroblast cells (Figure 1D). As the 9677 SMA patient cell line expresses approximately one-fourth of the SMN protein expressed by the 3814 carrier cell line,¹⁹ a 50% increase in SMN protein levels is modest but of interest. Toxicity was observed at concentrations at or above 80 μ g/mL (720 μ M), providing a robust effective concentration (EC) to lethal concentration (LC) ratio for cell culture studies. It has proven challenging to find small molecules that significantly upregulate endogenous SMN protein levels.²⁰ Thus, cuspin-1, despite its modest overall effect, was selected as a suitable probe for further characterization as it showed moderate activity and low cellular toxicity.

The ability of cuspin-1 to upregulate SMN protein levels in diverse genetic backgrounds was tested in a panel of cell lines, including wild-type and SMA human and rodent cells (Supplementary Table S3). Although cuspin-1 was active in the majority of human cell lines, it was inactive in the rodent lines tested, suggesting that its target or mechanism may not be conserved across species.

Structure–Activity Relationship of Cuspin-1. To evaluate the features necessary for the activity of cuspin-1, a structure–activity relationship (SAR) analysis was undertaken through analogue synthesis and testing. Although few structural analogues were commercially available (Figure 2A), one highly similar analogue (5-bromo-3-pyridinyl)(4-ethylphenyl)-methanone, designated cuspin-2, increased SMN protein abundance to comparable levels as cuspin-1. Although cuspin-2 was active, it did not improve upon the potency or efficacy of cuspin-1. All other cuspin-1 analogues that we synthesized, including analogues lacking either the bromine, nitrogen, or methyl group, were inactive (Figure 2B,C). Additionally, while cuspin-2 demonstrated that extension of the methyl to an ethyl group retained activity, extension to a propyl group rendered the compound inactive, indicating a possible steric clash with cuspin-1's target. As modifications to the cuspin-1 scaffold generally resulted in loss of activity, further work is required to identify modifications that would improve the potency and efficacy of the compound, prior to testing in SMA mouse models. However, the current cuspin-1 scaffold represents a valuable tool for the study of SMN regulation.

Cuspin-1 Reveals Link between Ras Signaling and SMN Protein Levels. During our efforts to identify the mechanism of action of cuspin-1 we utilized phospho-specific antibodies to determine whether any common signaling pathways were activated by cuspin-1 treatment. We observed that treatment with cuspin-1 increased phosphorylation of extracellular signal-related kinase (Erk), while proteins involved




Figure 3. Increased Ras signaling upregulates SMN protein abundance. (A) Western blot analysis of 9677 cells treated with 10 μ g/mL cuspin-1, harvested at indicated time points, and blotted for p-Erk, total Erk, p-Akt, total Akt, and tubulin. Graph shows quantification of phospho-protein to respective total protein ratios, normalized to tubulin as loading control. Data represents the average of duplicates \pm standard deviation. (B) Western blot analysis for NRas, p-Erk, SMN, and tubulin in 3813 SMA patient fibroblast cells transduced with either empty vector or constitutively active $NRas^{Q61K}$ viral plasmid. Graph shows fold increase of SMN compared to vector-only control, normalized to tubulin as loading control. (C) Comparison of the engineered tumor cell line BJeLR (eLR), expressing $HRas^{V12}$, versus the parental line (eH) and the isogenic line BJeHLT (eHLT), which contains all genes ectopically expressed in BJeLR with the exception of $HRas^{V12}$. Samples were blotted for SMN and several control proteins. Graph shows the fold increase of SMN in BJeHLT and BJeLR compared to BJeH, normalized to actin as loading control. Data represent average of duplicates \pm standard deviation.

in other signaling pathways, such as Akt, were not activated (Figure 3A). As the canonical pathway resulting in phosphorylation of Erk (p-Erk) is the Ras-Raf-MEK signaling cascade,²¹ we endeavored to determine whether increased Ras signaling could be the downstream event responsible for the cuspin-1 mediated upregulation of SMN protein levels. We increased Ras signaling in 3813 SMA patient fibroblast cells by expression of a constitutively active form of Ras ($NRas^{Q61K}$). $NRas^{Q61K}$ expression resulted in a 2.7-fold increase in SMN protein level compared to vector alone (Figure 3B), demonstrating that direct activation of the Ras signaling pathway was able to strongly increase SMN protein levels. The 5-fold increase in p-Erk in the $NRas^{Q61K}$ -expressing cells confirmed that Ras signaling was elevated in these cells. Similar increases in SMN and p-Erk levels were observed in 9677 and 232 SMA

patient fibroblasts, indicating this effect of *NRas*^{Q61K} is not unique to one SMA patient cell line (data not shown).

At this stage, *cuspin-1*'s utility as a probe was overshadowed by the more powerful effect of direct Ras activation on SMN protein levels. Therefore, we chose to continue characterizing the effect using constitutively active Ras rather than *cuspin-1*. However, further experiments investigating the effect of activated Ras on SMN protein levels could not be performed in the SMA patient cell lines, as nontransformed cells senesce shortly after introduction of oncogenic Ras.²¹ Therefore, we utilized the tumorigenic *HRas*^{V12}-expressing BJeLR cell line for subsequent experiments. The G12V mutation in the *HRas*^{V12} protein is functionally equivalent to the Q61K mutation used previously, as they both inactivate GTPase activity, rendering the proteins constitutively active. BJeLR cells are an engineered cell line created by sequential introduction of viral and mammalian genetic elements into parental BJeH human foreskin fibroblasts.²² The BJeHLT cell line contains all genetic elements found in the BJeLR cell line, with the exception of *HRas*^{V12}. Comparison of SMN protein levels in the BJ cell lines demonstrated a 3-fold increase in SMN protein in the BJeLR cell line compared to the parental cell line, BJeH (Figure 3C). The lack of increase in SMN in the BJeHLT cell line indicates that activated Ras is the sole genetic element responsible for upregulation of SMN protein abundance in this cell line (Figure 3C). This upregulation in SMN protein could be the result of either global upregulation of protein levels or via a more specific mechanism. To exclude the possibility that the observed effect on SMN protein levels was due to a nonspecific mechanism, the levels of several control proteins were examined by Western blot. These proteins are involved in diverse cellular processes, including glucose metabolism (GAPDH), the cytoskeleton (actin), translation (eIF4E), and cell death (Smac). The abundance of these proteins remained unaltered by oncogenic Ras expression (Figure 3C), indicating a level of specificity to Ras-mediated SMN upregulation. In hindsight, it is perhaps not so surprising that a pro-growth factor such as Ras would upregulate SMN, a protein vital for the splicing of mRNAs, as cellular growth would require this process. Although activating Ras signaling likely increases the levels of other pro-growth proteins in addition to SMN in these cell lines, at this stage we are more interested in the mechanism by which Ras increases SMN protein levels, rather than identifying the other proteins affected. As increasing Ras signaling would be therapeutically undesirable due to the likelihood of tumorigenesis, the specificity of its effect is not clinically relevant.

Mechanistic Studies. To investigate the mechanism underlying SMN protein upregulation by activated Ras, we utilized the BJ cell lines to determine the cellular process through which Ras exerts its effect on SMN protein levels. First, we tested whether alterations in transcription or splicing of *smn* mRNA were responsible for the increase in SMN protein abundance. The BJeLR (+*HRas*^{V12}) cell line did not show increased levels of either full-length or $\Delta 7$ *smn* mRNA, nor did it have an altered ratio of the spliced isoforms, as compared to the parental line BJeH (*-HRas*^{V12}) (Figure 4A). This lack of increase in *smn* mRNA levels suggests that the upregulation of SMN protein levels was not due to activation of p38 mitogen-activated kinase, a Ras effector protein, which was recently shown to upregulate SMN protein levels via stabilization of *smn* mRNA.²³ Unexpectedly, the BJeLR line showed consistently lower levels of *smn* mRNAs than the BJeH parental line. This




Figure 4. Increased Ras signaling enhances SMN translation rate. (A) Quantification of RT-qPCR of full-length (*smnFL*) and exon 7-deleted (*smnΔ7*) *smn* mRNA levels from total RNA harvested from BJeH (*-HRas*^{V12}) and BJeLR (+*HRas*^{V12}) cells. Transcript cycle times were normalized to *hprt* to control for total mRNA. Data represent average of triplicates \pm standard deviation. (B) Western blot analysis of SMN and actin in BJeH (*-HRas*^{V12}) and BJeLR (+*HRas*^{V12}) cells treated with the protein synthesis inhibitor cycloheximide (CHX, 10 μ M) for indicated times. Quantification of average of duplicates of SMN to actin ratio normalized to respective untreated control (0 h) is shown below each lane. (C) Phosphorimager bands and quantification of translation rate analysis of SMN, quantified as SMN to actin ratio normalized to BJeH (*-HRas*^{V12}) cells. Data represent mean \pm standard deviation. *p*-values were calculated using a two-tailed *t* test (* = *p* < 0.01). (D) Phosphorimager lanes showing total ³⁵S protein labeling in treated and untreated samples.

could be due to a feedback mechanism downregulating *smn* transcripts in response to strongly elevated SMN protein levels.

Next, we examined the effect of increased Ras signaling on SMN turnover. Cycloheximide (CHX), a protein synthesis inhibitor, was used to inhibit the translation of nascent SMN peptides in order to quantify the half-life of SMN. The half-life of SMN in the BJ cell lines was \sim 3 h, similar to previous reports.^{24,25} The degradation rate of SMN protein remained unchanged in the *HRas*^{V12}-expressing cells, indicating that increased Ras signaling does not affect the rate of SMN degradation (Figure 4B).

Having ruled out alterations in transcription, splicing and turnover, we tested whether *HRas*^{V12} affected the translation rate of SMN. A translation rate assay, using radioactive ³⁵S-methionine and cysteine to label newly synthesized protein with subsequent immunoprecipitation of proteins of interest, was performed to quantify the translation rate of SMN. *HRas*^{V12}-expressing cells showed a significant 2.5-fold increase (*p* < 0.01) in the rate of SMN translation compared to parental cells (Figure 4C). No differences were observed in the total amount of label incorporated in each cell line (Figure 4D), indicating the increased translation rate of SMN protein is not due to a global increase in translation. As we do not consider utilization of CA-Ras to be a therapeutic option, we are not concerned with the specificity of the effect of Ras activation. Rather, we are excited to discover such a robust effect on SMN protein levels via an unexpected pathway. Future work determining the cellular components downstream of Ras

directly responsible for affecting the translation rate of SMN may identify therapeutic targets of interest for SMA, at which stage the specificity of the effect will be of paramount importance.

In this work, we describe the development of a high-throughput screen for identification of small molecule upregulators of the SMN protein and identify three novel SMN-upregulating compounds. Mechanistic studies of cusp-1 led to the discovery that Ras signaling regulates SMN protein abundance at the level of translational regulation. In addition to the novel SMN regulatory pathway identified, this study provides a proof of principle that mechanistically unbiased screens can uncover unexpected connections between cellular pathways and provide novel small molecule tools for the study of biological processes.

Small molecule upregulators of protein abundance are an underutilized class of bioactive compounds, with the potential to elucidate protein regulation and function. The work presented here defines a straightforward method for identifying small molecule upregulators of a target protein's abundance. These compounds have two important uses. First, on the basis of the premise that upregulating the abundance of a target protein will increase its level of cellular activity, such compounds could act as protein function activators, allowing dose-dependent and reversible studies on the effects of increasing protein function. Discovery of small molecule activators has mainly focused on identifying compounds that increase the intrinsic activity of the steady-state levels of proteins. Small molecule activators have been identified for a handful of targets, such as caspases,²⁶ the proteasome,²⁷ and several cell surface receptors.^{28–30} However, the number and classes of protein targets for these direct activators are likely as limited as those for inhibitors of protein functions. In comparison to inhibitors, upregulators are not restricted to specific protein families, as they do not rely on the protein of interest being directly “druggable”, generally defined as the presence of a catalytic region that interacts with small molecules or an endogenous ligand binding site.³¹ Upregulators have the advantage of targeting members of the regulatory pathways of the protein of interest, greatly increasing the range of protein families that can be modulated in this manner.

The second advantage of small molecule upregulators is their utility as probes to investigate the pathways involved in regulating the levels of a target protein. As many currently intractable conditions fall into the category of loss of function diseases, such as spinal muscular atrophy, cystic fibrosis, and Duchenne muscular dystrophy,³² elucidation of the pathways governing the abundance of the disease gene products may identify new therapeutic candidates. Additionally, small molecule upregulators of sufficient potency and efficacy have the potential to be developed into lead therapeutics.

Using a high-throughput screening approach for identification of small molecule upregulators of protein abundance, we discovered three small molecule upregulators of the Survival of Motor Neuron (SMN) protein. SMN has been shown to be involved in many cellular processes, such as pre-mRNA splicing³³ and β -actin mRNA localization to neuronal growth cones.³⁴ SMN protein levels are decreased in the loss of function neurodegenerative disease spinal muscular atrophy,³⁵ making it a target of interest for both research and biomedical applications. The most active scaffold identified in our screen, a bromobenzophenone analogue designated cusp-1 (Chemical Upregulator of SMN Protein-1) was utilized as a probe to

reveal a novel regulatory pathway governing SMN protein abundance. Treatment of Type I SMA fibroblast cells with cusp-1 resulted in a 50–100% increase in endogenous SMN protein levels, as well as a concomitant increase in the phosphorylation of Erk. Since the major pathway responsible for phosphorylation of Erk is the Ras-Raf-MEK cascade,²¹ we expressed constitutively active isoforms of Ras and determined that increasing Ras signaling resulted in a robust ~270–300% increase in SMN protein abundance. Mechanistic studies revealed that this upregulation was due to Ras-mediated enhancement of SMN translation rate.

Increasing the translation rate of a target protein is an unusual mechanism of action for a small molecule; however, it may be a versatile one, as recent studies suggest that translation regulation may have a role in diverse cellular processes such as cancer, aging, and mitochondrial function.³⁶ Future studies defining the mechanism by which increased Ras signaling affects translation of SMN protein could identify pathways of interest for SMA and perhaps other loss of function diseases. Although Ras is best known for its oncogenic properties, its ability to upregulate non-oncogenic proteins, such as SMN, demonstrate potential therapeutically beneficial downstream consequences of Ras activation. There is also significant evidence for a pro-survival effect of Ras activation, especially in neurons, where Ras has been shown to be activated by pro-survival neurotrophic factors.³⁷ Expression of constitutively active isoforms of Ras (CA-Ras) has shown beneficial effects on the survival and outgrowth of cultured chick embryonic neurons.³⁸ Additionally, in a mouse model of motor neuron degeneration, neuron-selective expression of CA-Ras resulted in complete rescue from lesion-induced degeneration.³⁹ The identification of the mechanisms downstream of Ras activation resulting in cell survival could lead to novel therapeutic targets for diseases resulting from pathological cell death.

METHODS

Cell Culture. Type I spinal muscular atrophy (SMA)-affected human primary fibroblasts were obtained live with low passage number (Coriell Cell Repositories; nos. GM03813, GM09677, GM00232). The cells were cultured in Minimum Essential Medium (MEM) with Earle's salts and nonessential amino acids (Invitrogen; no. 10370-021) supplemented with 15% (v/v) fetal bovine serum (FBS) (Invitrogen; no. 26140-079), 2 mM L-glutamine (Invitrogen; no. 25030-081), and antibiotics (100,000 units penicillin L⁻¹ MEM, 100 mg streptomycin L⁻¹ MEM) (Invitrogen; no. 15140-122). Cells were allowed to grow in 50 mL of the aforementioned medium at 37 °C with 5% carbon dioxide in 175 cm² tissue culture treated flasks (Corning; no. 431080). The foreskin fibroblast-derived BJ cell lines were cultured in DMEM (Invitrogen; no. 11995-040) supplemented with 10% (v/v) FBS, 2 mM L-glutamine and antibiotics. Cells were grown under the same conditions as SMA cell lines.

Small Molecule Libraries Selected for Screening. The commercially available small molecule libraries selected for screening consisted of the National Institute of Neurological Disorders and Stroke (NINDS) Custom Collection of Known Bioactives (1,040 compounds), the Annotated Compound Library (ACL) (2,337 compounds), the TIC library (a select composite from TimTech, InterBioScreen and Chembridge totaling 23,685 compounds),¹⁵ and the Comgenex library (20,000 compounds).

A second unique set of small molecules, designated the blood-brain barrier (BBB) libraries, were selected from five commercially available compound libraries. These libraries contain a total of 1,322,183 unique structures that were subjected to *in silico* filtering using MOE 2004.03. All compounds were preprocessed by assigning polar hydrogen atoms and charges according to the MMFFx forcefield. Salts were converted to free-base or free-acid form. These preprocessed libraries were first

screened for reactive functional groups and unsuitable scaffolds, as defined by Hann et al.⁴⁰ using a SMILES-string-based search method. We then proceeded to assess the BBB-penetrating potential of the compounds that passed the first two filtering steps by constructing a set of filters to predict log BB (logarithm of brain to blood partitioning ratio). The molecules' log BB values were predicted using a model trained on 73 compounds with experimentally determined log BB values and validated on a set of 36 CNS positive and negative drugs. The obtained model gave an $R^2 = 0.8225$ for the training and $R^2 = 0.6989$ for the validating set. Of note, the predictions gave no false negatives in this data set.

From this screen, 126,877 compounds with predicted log BB values greater than zero (on a log scale from -3 to $+6$) were selected. A value of zero indicates equal partitioning between the blood and brain, while a positive value indicates that the compound would partition preferentially into CNS tissues compared to plasma. Of these compounds, we discarded those with molecular weights less than 250 or containing three or more aromatic rings. These filters were designed to eliminate very simple compounds or planar compounds that might intercalate into DNA.

The final 79,691 compounds passing these filters were submitted to clustering to assess their diversity and to remove close structural analogues, which are frequently created during syntheses used by commercial vendors. This clustering was achieved by annotating the compounds with MACCS (Molecular Design Limited) structural keys; the clustering itself was performed using the Jarvis–Patrick method. Molecular similarity was determined by the Tanimoto coefficient, the threshold for which was set at 0.7. The procedure yielded 27,577 structurally distinct clusters from four vendors. Three vendors (Asinex, Chembridge, and Life Chemicals) were chosen on the basis of pricing, and one compound from each cluster was purchased, yielding a library of 22,795 compounds likely to cross the blood–brain barrier and to be drug-like.

Small Molecule Screening. See Supporting Information for details on the screening protocols.

Analogue Synthesis. Solvents and starting materials were purchased from Sigma-Aldrich and used without further purification. Analogues were synthesized using chlorination and Friedel–Crafts acylation. Briefly, 50–100 mg of 5-bromonicotinic acid, or desired nicotinic acid, was dissolved in ~ 10 mL of anhydrous dichloromethane. Thionyl chloride (1.2 equiv) was added to the solution. Reaction progress was monitored using pH paper. Once complete, solvent was removed under high vacuum. Acyl chlorides were coupled to desired benzene derivative in anhydrous dichloromethane with aluminum trichloride (1.3 equiv) overnight. Unreacted starting materials were removed by aqueous extraction, and products were purified by chromatography in hexanes/ethyl acetate. Identity and purity was confirmed by mass spectrometry. Yields ranged between 20% and 75%, providing sufficient material for testing. Synthetically prepared cuspis-1 and cuspis-2 showed similar activity to that of the commercially sourced compounds, verifying the identity of the compounds. Cuspis-1: $^1\text{H NMR}$ (CDCl_3 , 400 MHz) 8.89 (t, $J = 2.2$ Hz, 2H), 8.25 (t, $J = 2.0$ Hz, 1H), 7.74 (d, $J = 8.2$ Hz, 2H), 7.35 (d, $J = 8.0$ Hz, 2H), 2.49 (s, 3H).

Western Blot Analysis. SMA patient fibroblast lines were plated at 30,000 cells per well in a 6-well plate ~ 24 h prior to compound treatment. Cells were treated with desired concentration of compound for 48 h. Cells were harvested at $<75\%$ confluence; confluence may be lower due to compound toxicity. Cells were washed twice in ice-cold PBS and lysed by scraping in 100 μL of ice-cold lysis buffer (50 mM HEPES, 40 mM NaCl, 2 mM EDTA, 0.5% Triton-X, 1.5 mM Na_3VO_4 , 50 mM NaF, 10 mM Na-pyrophosphate, 10 mM $\text{Na}\beta$ -glycerophosphate, Roche protease inhibitor tablet). Insoluble materials were spun out by centrifugation at 10,621g for 10 min at 4 $^\circ\text{C}$. Supernatants were boiled with SDS-PAGE sample buffer prior to being separated on a 4–12% Bis-Tris NuPage gel in 1X NuPage MES running buffer. Proteins were transferred to a PVDF membrane using Invitrogen iBlot blotting system, setting P3 for 6 min and 40 s. Membranes were blocked in Odyssey blocking buffer (LI-COR; no. 927-40010) for 20 min at RT and then incubated overnight at 4 $^\circ\text{C}$ with primary antibodies (see

below for dilutions). Membranes were washed three times in TBS with 0.1% Tween-20 and then incubated for 45–60 min with 680 nm-conjugated anti-mouse IgG secondary antibody at 1:3,000 (Invitrogen; no. A21058) and 800 nm-conjugated anti-rabbit IgG secondary antibody at 1:3,000 (LI-COR; no. 926-322110). Signals were detected and quantified using the LiCor Odyssey system and software (LI-COR Biosciences).

Antibodies. Mouse anti-SMN (BD Biosciences; no. 610647) (1:1,000), rabbit anti-actin (Santa Cruz Biotechnologies; no. sc-1616-R) (1:3,000), mouse anti-alpha tubulin (Santa Cruz Biotechnologies; no. sc-32293) (1:5,000), rabbit anti-phospho-Erk1/2 (Cell Signaling; no. 9101S) (1:1,000), rabbit anti-Erk1/2 (Cell Signaling; no. 9102) (1:3,000), rabbit anti-phospho-Akt (Cell Signaling; no. 9271) (1:2,000), rabbit anti-Akt (Cell Signaling; no. 9272) (1:2,000), mouse anti-NRas (EMD Biosciences; no. OP25) (1:2,000), mouse anti-eIF4E (BD Biosciences; no. 610269) (1:2,000), mouse anti-Smac/DIABLO (BD Biosciences; no. 612244) (1:1,000), and mouse anti-GAPDH (Santa Cruz Biotechnologies; no. sc-47724) (1:5,000).

Retroviral Preparation and Transduction. To prepare the retroviral supernatants, 2×10^6 Plat-GP cells (Cell Biolabs; no. RV-103) were seeded per 10 cm culture dish in DMEM supplemented with 10% (v/v) fetal bovine serum (FBS), 2 mM L-glutamine, antibiotics (100,000 units penicillin L^{-1} MEM, 100 mg streptomycin L^{-1} MEM), and 5 $\mu\text{g mL}^{-1}$ blasticidin (Invitrogen; no. 11811-031). Cells were allowed to attach overnight. The following morning pVSV-G helper plasmids with the MigR1 vector either empty or containing *NRas*^{Q61K} were cotransfected into the Plat-GP cells using FuGENE 6 Transfection Reagent (Roche; no. 11-814-443-001). Thirty hours post-transfection, culture media was replaced by Viral Collection Media (VCM, normal culture media with 30% (v/v) FBS). VCM was collected and replaced twice at 12 h intervals to collect viral particles released into the supernatant. Viral supernatant was filtered through 0.45 μm filters prior to use or storage at -80 $^\circ\text{C}$. For transduction, 3813 SMA patient fibroblast cells were plated at 2×10^5 cells per well into a 6-well plate and allowed to attach overnight. The following day, the infection cocktail was prepared by mixing 0.75 mL of viral supernatant, 1.25 mL of growth media, and 2 μL of 8 mg mL^{-1} Polybrene (Sigma; no. H9268). The culture media in each well was replaced with the infection cocktail, and then plates were spun at 1,089g (2,250 rpm with 19.2 cm rotor radius) for 90 min before incubation. Cells were cultured for 2 days prior to harvesting.

RT-qPCR of *smn* Transcripts. RNA was harvested from 5×10^6 cells using the Qiagen RNeasy Kit as per manufacturer's protocol. RNA from each sample (2 μg) was reverse transcribed using Applied Biosystems Taqman kit (N808-0234) as per manufacturer's protocol. Quantitative PCR was performed on a 7300 Real-Time PCR System (Applied Biosystems) using 1X Power SYBR Green PCR Master Mix (Applied Biosystems; no. 4367659), 77 ng cDNA and 0.15 μM primers per well in a MicroAmp Optical 96-well plate (Applied Biosystems; no. N801-0560). The cycle times of the *smn* transcripts were normalized to cycle times for hypoxanthine guanine phosphoribosyltransferase (*hprt*) as loading control. Amplification conditions: 50 $^\circ\text{C}$ for 2 min, 95 $^\circ\text{C}$ for 10 min, 40 cycles of 95 $^\circ\text{C}$ for 15 s and 60 $^\circ\text{C}$ for 1 min. $\Delta\Delta\text{CT}$ was calculated after normalization to *hprt* expression. Primer sequences: *smnFL*-forward: 5'-CAAAAAGAAGG-AAGGTGCTCA-3', *smnFL*-reverse: 5'-TGGTGTCATTTAGTGC-TGCTC-3', *smn* Δ 7-forward: 5'-ATTCTCTTGATGATGCTGATGCT-3', *smn* Δ 7-reverse: 5'-TATGCCACCAATTCATATAATAG-3', *hprt*-forward: 5'-TGACACTGCAGCAAAATGCA-3', *hprt*-reverse: 5'-GGTCCTTTTACCAGCAAGCT-3'

^{35}S -Methionine/Cysteine Translation Rate Assay. Cells were seeded at 5×10^6 cells per 10 cm dish and allowed to attach overnight. For methionine/cysteine starvation, cells were rinsed twice in PBS and incubated for 30 min in starvation media: Met/Cys-free DMEM (Invitrogen; no. 21013-024), 10% dialyzed FBS (v/v), and 2 mM L-glutamine. After the starvation period, cells were pulsed with 100 μCi TRAN35S-Label Metabolic Labeling Reagent (MP; no. 015100907) for 1 h. Cells were lysed in 1 mL of IP Buffer (50 mM Tris [pH 7.75], 250 mM NaCl, 200 mM EDTA, 150 mM MgCl_2 , 0.3% NP-40, 1% Empigen BB and Roche protease inhibitor tablet) and scraped, and

insoluble materials were spun out at 10,621g for 10 min at 4 °C. Lysates were precleared with 20 μ L Protein A/G-agarose bead slurry (Pierce; no. 20421) for 30 min, and supernatants were transferred to tubes containing 30 μ L of Protein A/G-agarose precoupled to 2.5 μ g of mouse anti-SMN antibody (BD Biosciences; no. 610647) and immunoprecipitated overnight at 4 °C. After SMN IP, lysates were transferred to tubes containing 30 μ L of Protein A/G-agarose precoupled to 2 μ g of rabbit anti-actin antibody (Santa Cruz Biotechnologies; no. sc-1616-R) and immunoprecipitated for 4 h at 4 °C. Beads were washed three times in IP buffer, and bound proteins were eluted by boiling in SDS-PAGE buffer. Samples were run on NuPAGE 4–12% Bis-Tris gels (Invitrogen; no. WG1402BOX) in 1X NuPAGE MES running buffer, fixed in 50% methanol +10% acetic acid, and dried before exposing overnight to phosphorimager screen. Bands were imaged using a Storm phosphorimager (Molecular Dynamics) and quantification was performed using ImageQuant software (Molecular Dynamics). Values were reported as SMN to actin ratio normalized to BJeH ($-HRas^{V12}$).

■ ASSOCIATED CONTENT

● Supporting Information

Supplemental figures illustrating the validation of *in silico* screening filters, the SMN-upregulating activity of hit compounds c72 and c81, and the ^1H NMR of cusp-1; three tables listing the summary of the screening data, a list of descriptors used for *in silico* calculations, as well as results of genotype-specificity testing of cusp-1; supplemental methods describing the methods used for the cyto blot screening procedure. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: bstockwell@columbia.edu.

Present Address

[§]Brigham and Women's Hospital, 75 Francis Street, Boston, Massachusetts 02115, United States

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful to G. Morris (Centre for Inherited Neuromuscular Disease, RJA Orthopaedic Hospital; Oswestry, U.K.) for his gift of mouse monoclonal SMN antibodies. We kindly thank C. Henderson (Center for Neurobiology and Behavior, Columbia University, NY) for the experiments involving mouse embryonic stem cell-derived motor neurons. We thank B. Burnett (Neurogenetics Branch, NINDS, NIH, Bethesda, MD) for confirmatory testing of cusp-1. B.R.S. is an Early Career Scientist of the Howard Hughes Medical Institute. This work was supported by The SMA Foundation and the National Institutes of Health [R01GM085081, R01CA161061, and R01 CA097061].

■ REFERENCES

- (1) Sopko, R., Huang, D., Preston, N., Chua, G., Papp, B., Kafadar, K., Snyder, M., Oliver, S. G., Cyert, M., Hughes, T. R., Boone, C., and Andrews, B. (2006) Mapping pathways and phenotypes by systematic gene overexpression. *Mol. Cell* 21, 319–330.
- (2) Atmaca, M. (2009) Valproate and neuroprotective effects for bipolar disorder. *Int. Rev. Psychiatry* 21, 410–413.
- (3) Trinka, E. (2007) The use of valproate and new antiepileptic drugs in status epilepticus. *Epilepsia* 48 (Suppl 8), 49–51.
- (4) Marks, P. A. (2007) Discovery and development of SAHA as an anticancer agent. *Oncogene* 26, 1351–1356.
- (5) Richardson, P. G., Hideshima, T., and Anderson, K. C. (2003) Bortezomib (PS-341): a novel, first-in-class proteasome inhibitor for the treatment of multiple myeloma and other cancers. *Cancer Control* 10, 361–369.
- (6) Sun, L., Trausch-Azar, J. S., Ciechanover, A., and Schwartz, A. L. (2007) E2A protein degradation by the ubiquitin-proteasome system is stage-dependent during muscle differentiation. *Oncogene* 26, 441–448.
- (7) Chateauvieux, S., Morceau, F., Dicato, M., and Diederich, M. (2010) Molecular and therapeutic potential and toxicity of valproic acid. *J. Biomed. Biotechnol.*, 2010.
- (8) Monani, U. R., Lorson, C. L., Parsons, D. W., Prior, T. W., Androphy, E. J., Burghes, A. H., and McPherson, J. D. (1999) A single nucleotide difference that alters splicing patterns distinguishes the SMA gene SMN1 from the copy gene SMN2. *Hum. Mol. Genet.* 8, 1177–1183.
- (9) Burghes, A. H., and Beattie, C. E. (2009) Spinal muscular atrophy: why do low levels of survival motor neuron protein make motor neurons sick? *Nat. Rev. Neurosci.* 10, 597–609.
- (10) Makhortova, N. R., Havhurst, M., Cerquiera, A., Sinor-Anderson, A. D., Zhao, W. N., Heiser, P. W., Arvanites, A. C., Davidow, L. S., Waldon, Z. O., Steen, J. A., Lam, K., Ngo, H. D., and Rubin, L. L. (2011) A screen for regulators of survival of motor neuron protein levels. *Nat. Chem. Biol.* 7, 544–552.
- (11) Jarecki, J., Chen, X., Bernardino, A., Coovert, D. D., Whitney, M., Burghes, A., Stack, J., and Pollok, B. A. (2005) Diverse small-molecule modulators of SMN expression found by high-throughput compound screening: early leads towards a therapeutic for spinal muscular atrophy. *Hum. Mol. Genet.* 14, 2003–2018.
- (12) Andreassi, C., Jarecki, J., Zhou, J., Coovert, D. D., Monani, U. R., Chen, X., Whitney, M., Pollok, B., Zhang, M., Androphy, E., and Burghes, A. H. (2001) Aclarubicin treatment restores SMN levels to cells derived from type I spinal muscular atrophy patients. *Hum. Mol. Genet.* 10, 2841–2849.
- (13) Stockwell, B. R., Haggarty, S. J., and Schreiber, S. L. (1999) High-throughput screening of small molecules in miniaturized mammalian cell-based assays involving post-translational modifications. *Chem. Biol.* 6, 71–83.
- (14) Root, D. E., Flaherty, S. P., Kelley, B. P., and Stockwell, B. R. (2003) Biological mechanism profiling using an annotated compound library. *Chem. Biol.* 10, 881–892.
- (15) Zhang, J. H., Chung, T. D., and Oldenburg, K. R. (1999) A simple statistical parameter for use in evaluation and validation of high throughput screening assays. *J. Biomol. Screening* 4, 67–73.
- (16) Kelley, B. P., Lunn, M. R., Root, D. E., Flaherty, S. P., Martino, A. M., and Stockwell, B. R. (2004) A flexible data analysis tool for chemical genetic screens. *Chem. Biol.* 11, 1495–1503.
- (17) Rose, K., Hall, L. H., and Kier, L. B. (2002) Modeling blood-brain barrier partitioning using the electrotopological state. *J. Chem. Inf. Comput. Sci.* 42, 651–666.
- (18) Wildman, S. A., and Crippen, G. M. (1999) Prediction of Physicochemical Parameters by Atomic Contributions. *J. Chem. Inf. Comput. Sci.* 39, 868–873.
- (19) Coovert, D. D., Le, T. T., McAndrew, P. E., Strasswimmer, J., Crawford, T. O., Mendell, J. R., Coulson, S. E., Androphy, E. J., Prior, T. W., and Burghes, A. H. (1997) The survival motor neuron protein in spinal muscular atrophy. *Hum. Mol. Genet.* 6, 1205–1214.
- (20) Lunn, M. R., and Wang, C. H. (2008) Spinal muscular atrophy. *Lancet* 371, 2120–2133.
- (21) Vojtek, A. B., and Der, C. J. (1998) Increasing complexity of the Ras signaling pathway. *J. Biol. Chem.* 273, 19925–19928.
- (22) Hahn, W. C., Counter, C. M., Lundberg, A. S., Beijersbergen, R. L., Brooks, M. W., and Weinberg, R. A. (1999) Creation of human tumour cells with defined genetic elements. *Nature* 400, 464–468.
- (23) Farooq, F., Balabanian, S., Liu, X., Holcik, M., and MacKenzie, A. (2009) p38 Mitogen-activated protein kinase stabilizes SMN mRNA through RNA binding protein HuR. *Hum. Mol. Genet.* 18, 4035–4045.

- (24) Burnett, B. G., Munoz, E., Tandon, A., Kwon, D. Y., Sumner, C. J., and Fischbeck, K. H. (2009) Regulation of SMN protein stability. *Mol. Cell. Biol.* 29, 1107–1115.
- (25) Cho, S., and Dreyfuss, G. (2010) A degron created by SMN2 exon 7 skipping is a principal contributor to spinal muscular atrophy severity. *Genes Dev.* 24, 438–442.
- (26) Wolan, D. W., Zorn, J. A., Gray, D. C., and Wells, J. A. (2009) Small-molecule activators of a proenzyme. *Science* 326, 853–858.
- (27) Huang, L., Ho, P., and Chen, C. H. (2007) Activation and inhibition of the proteasome by betulinic acid and its derivatives. *FEBS Lett.* 581, 4955–4959.
- (28) Grimm, C., Jors, S., Saldanha, S. A., Obukhov, A. G., Pan, B., Oshima, K., Cuajungco, M. P., Chase, P., Hodder, P., and Heller, S. (2010) Small molecule activators of TRPML3. *Chem. Biol.* 17, 135–148.
- (29) Mruk, K., and Kobertz, W. R. (2009) Discovery of a novel activator of KCNQ1-KCNE1 K channel complexes. *PLoS One* 4, e4236.
- (30) Small, K. M., McGraw, D. W., and Liggett, S. B. (2003) Pharmacology and physiology of human adrenergic receptor polymorphisms. *Annu. Rev. Pharmacol. Toxicol.* 43, 381–411.
- (31) Hopkins, A. L., and Groom, C. R. (2002) The druggable genome. *Nat. Rev. Drug Discovery* 1, 727–730.
- (32) Segalat, L. (2007) Loss-of-function genetic diseases and the concept of pharmaceutical targets. *Orphanet J. Rare Dis.* 2, 30.
- (33) Pellizzoni, L., Kataoka, N., Charroux, B., and Dreyfuss, G. (1998) A novel function for SMN, the spinal muscular atrophy disease gene product, in pre-mRNA splicing. *Cell* 95, 615–624.
- (34) Rossoll, W., Jablonka, S., Andreassi, C., Kroning, A. K., Karle, K., Monani, U. R., and Sendtner, M. (2003) Smn, the spinal muscular atrophy-determining gene product, modulates axon growth and localization of beta-actin mRNA in growth cones of motoneurons. *J. Cell Biol.* 163, 801–812.
- (35) Lefebvre, S., Burlet, P., Liu, Q., Bertrand, S., Clermont, O., Munnich, A., Dreyfuss, G., and Melki, J. (1997) Correlation between severity and SMN protein level in spinal muscular atrophy. *Nat. Genet.* 16, 265–269.
- (36) Sonenberg, N., and Hinnebusch, A. G. (2009) Regulation of translation initiation in eukaryotes: mechanisms and biological targets. *Cell* 136, 731–745.
- (37) Segal, R. A., and Greenberg, M. E. (1996) Intracellular signaling pathways activated by neurotrophic factors. *Annu. Rev. Neurosci.* 19, 463–489.
- (38) Borasio, G. D., John, J., Wittinghofer, A., Barde, Y. A., Sendtner, M., and Heumann, R. (1989) ras p21 protein promotes survival and fiber outgrowth of cultured embryonic neurons. *Neuron* 2, 1087–1096.
- (39) Heumann, R., Goemans, C., Bartsch, D., Lingenhohl, K., Waldmeier, P. C., Hengerer, B., Allegrini, P. R., Schellander, K., Wagner, E. F., Arendt, T., Kamdem, R. H., Obst-Pernberg, K., Narz, F., Wahle, P., and Berns, H. (2000) Transgenic activation of Ras in neurons promotes hypertrophy and protects from lesion-induced degeneration. *J. Cell Biol.* 151, 1537–1548.
- (40) Hann, M., Hudson, B., Lewell, X., Lively, R., Miller, L., and Ramsden, N. (1999) Strategic pooling of compounds for high-throughput screening. *J. Chem. Inf. Comput. Sci.* 39, 897–902.