

A mitochondrial DNA phylogeny of African clawed frogs: phylogeography and implications for polyploid evolution

Ben J. Evans,^{a,*} Darcy B. Kelley,^b Richard C. Tinsley,^c Don J. Melnick,^d
and David C. Cannatella^e

^a Department of Biology, McMaster University, Life Sciences Building, Room 328, 1280 Main Street West, Hamilton, Ont., Canada L8S 4K1

^b Department of Biological Sciences, Fairchild Building, Columbia University, New York, NY, USA

^c School of Biological Sciences, University of Bristol, Bristol B58 1UG, UK

^d Center for Environmental Research and Conservation and Department of Ecology, Evolution, and Environmental Biology,
1200 Amsterdam Ave, Columbia University, New York 10027, USA

^e Section of Integrative Biology, C0930, and Texas Memorial Museum, The University of Texas, Austin, TX 78712-0253, USA

Received 11 January 2004; revised 20 April 2004

Available online 8 June 2004

Abstract

The African clawed frogs (*Silurana* and *Xenopus*), model organisms for scientific inquiry, are unusual in that allopolyploidization has occurred on multiple occasions, giving rise to tetraploid, octoploid, and dodecaploid species. To better understand their evolution, here we estimate a mitochondrial DNA phylogeny from all described and some undescribed species. We examine the timing and location of diversification, and test hypotheses concerning the frequency of polyploid speciation and taxonomy. Using a relaxed molecular clock, we estimate that extant clawed frog lineages originated well after the breakup of Gondwana, about 63.7 million years ago, with a 95% confidence interval from 50.4 to 81.3 million years ago. *Silurana* and two major lineages of *Xenopus* have overlapping distributions in sub-Saharan Africa, and dispersal–vicariance analysis suggests that clawed frogs originated in central and/or eastern equatorial Africa. Most or all extant species originated before the Pleistocene; recent rainforest refugia probably acted as “lifeboats” that preserved existing species, rather than “species pumps” where many new successful lineages originated. We estimate that polyploidization occurred at least six times in clawed frogs.

© 2004 Elsevier Inc. All rights reserved.

Keywords: *Xenopus*; *Silurana*; Pipidae; Genome duplication; Hybridization; Polyploidy; Rainforest refugia

1. Introduction

The clawed frogs *Xenopus laevis* and *Silurana tropicalis* are the most intensively studied amphibians; they are readily maintained in captivity, have large oocytes that facilitate investigation and manipulation of embryogenesis, transgenesis is now possible, and the DNA sequence of the complete genome of *Silurana tropicalis* soon will be completed. However, a comprehensive phylogeny is still unavailable because of the challenge of sampling these animals throughout their distribution in sub-Saharan Africa, and also because of an unusual

mode of evolution by allopolyploidization, which complicates phylogeny estimation. A phylogeny is central to understanding the timing and placement of diversification of this interesting group, and can be used to tease apart details of allopolyploid speciation.

1.1. Natural history and systematics

Clawed frogs comprise *Xenopus* and *Silurana*, which are sister taxa in the subfamily Xenopodinae, family Pipidae (Cannatella and de Sá, 1993; de Sá and Hillis, 1990). The other pipid subfamily, Pipinae, includes the New World *Pipa* (7 species), and the African sister genera *Hymenochirus* (4 species), and *Pseudhymenochirus* (1 species; Cannatella and Trueb, 1988a,b). Pipoid frogs include the Pipidae and the New World sister

* Corresponding author. Fax: +1-905-522-6066.

E-mail address: evansb@mcmaster.ca (B.J. Evans).

family Rhinophryinae (Ford and Cannatella, 1993) among living forms.

Pipid frogs are aquatic anurans that live principally in still or slow moving water. A derived morphology facilitates aquatic life, and is characterized by a lateral line system that persists through metamorphosis, laterally positioned limbs and dorsoventral compression of the body, pelvic modifications that facilitate swimming but impede terrestrial jumping, and modifications of the head and cranium including the loss of a tongue (Trueb, 1996). The larynx is exceptionally sexually dimorphic (larger in males; Sassoon and Kelley, 1986) and produces underwater vocalizations using a unique mechanism of sound production (Yager, 1996). At least one species has an elaborate repertoire of communication characterized by multiple call types in males and in females, and dueting between the sexes and multiple male call types (Tobias et al., 1998, 2004). Clawed frogs probably aestivate underground during dry periods and may migrate over land and through flowing water (McCoid and Fritts, 1980; Tinsley and McCoid, 1996).

Silurana includes one diploid species ($2N=20$) and one tetraploid species ($2N=40$); additionally one diploid (referred to here as *S. cf. tropicalis*) and two tetraploids are suggested by divergent mtDNA, and are undescribed (see below). *Xenopus* includes 10 tetraploid species ($2N=36$), 5 octoploid species ($2N=72$), and 2 dodecaploid species ($2N=108$) (Schmid and Steinlein, 1991; Tymowska, 1991). One of the 10 tetraploids, “*Xenopus muelleri* west” (Kobel et al., 1996, 1998), is referred to as *X. “new tetraploid”* in this paper. We identified two tetraploids similar to *Xenopus fraseri*, and refer to them as *X. cf. fraseri* 1 and 2. *Xenopus* has been divided into five species groups based on similarities in morphology, advertisement call, and/or molecular characters: (1) the *laevis* group, which includes the tetraploids *X. laevis*, *X. gilli*, and *X. largeni*; (2) the *muelleri* group, which includes the tetraploids *X. muelleri*, *X. borealis*, and *X. clivii*; (3) the *vestitus-wittei* group, which includes the octoploids *X. wittei* and *X. vestitus*, (4) the *fraseri*-like group, which includes the tetraploids *X. fraseri*, *X. pygmaeus*, the octoploids *X. amieti*, *X. andrei*, *X. boumbaensis*, and the dodecaploid *X. ruwenzoriensis*, and (5) the *longipes* group which includes only the dodecaploid *X. longipes* (Kobel et al., 1996). Additional information is necessary to test whether some of the divergent mtDNA lineages do constitute separate species; we view species assignments in this study that are based only on mtDNA as working hypotheses (Evans et al., 2003).

A mitochondrial DNA phylogeny of 7 species was estimated and reanalyzed from mapped restriction endonuclease sites of 11 enzymes in the complete mtDNA genome (Cannatella and de Sá, 1993; Carr et al., 1987). Another mtDNA phylogeny for 11 species was estimated from about 600 bp of mitochondrial ribosomal

DNA (rDNA) sequence (Kobel et al., 1998). Relationships among some species have also been estimated from protein data (Bisbee et al., 1977; Graf, 1996; Mann et al., 1982), and among extant and fossil taxa from morphology (Trueb and Báez, 1997).

1.2. Allopolyploid speciation and mitochondrial DNA

Speciation of clawed frogs can occur without genome duplication, as is common in other organisms; an ancestor gives rise to two descendants with the same number of chromosomes. Alternatively, speciation of clawed frogs can occur by reticulation via allopolyploidization; here two species hybridize and give rise to a descendant species that inherits the complete nuclear genome of both ancestors. Allopolyploidization rather than autopolyploidization is probably the principal mode of genome duplication in clawed frogs (Kobel, 1996a). Some polyploid species, for example, are morphologically similar to one species but carry mtDNA more closely related to another (Carr et al., 1987; Kobel et al., 1998). Hybrids are also known to occur in nature. The widely distributed *X. laevis* hybridizes with three other sympatric species: *X. gilli* in Cape Province, South Africa (Evans et al., 1997, 1998; Picker et al., 1996), *X. muelleri* in northern South Africa (Fischer et al., 2000), and *X. borealis* in Kenya (Yager, 1996). Hybrid crosses have yielded polyploid progeny in the laboratory (Kobel, 1996a; Kobel and Du Pasquier, 1986) and functional diploidization of some polyploid species is suggested by the formation of bivalents during meiosis (Kobel et al., 1996).

In *Xenopus* a diploid ancestor with 18 chromosomes is suggested by extant tetraploid karyotypes ($2N=36$), but presumed extinct. Thus, polyploidization in *Xenopus* is inferred to have occurred at least once from 18 to 36 chromosomes, at least once from 36 to 72 chromosomes, and at least once from a combination of genomes with 36 and 72 chromosomes to 108 chromosomes. In *Silurana*, polyploidization is inferred to have occurred at least once from 20 to 40 chromosomes.

As an initial step to unraveling the evolutionary history of this allopolyploid group, we have constructed a phylogeny based on mitochondrial DNA. In allopolyploid speciation, a polyploid descendant inherits the complete nuclear genome of two ancestors, but mitochondrial DNA is inherited from only one ancestor. Because mitochondrial DNA is maternally inherited and haploid, a mitochondrial DNA phylogeny only reflects a portion of the ancestry of a species that originated from allopolyploidization (species C, Figs. 1a and b). A mitochondrial DNA phylogeny also offers only a minimum estimate of the number of times polyploidization has occurred, because a single mtDNA lineage might occur in polyploids that have multiple independent origins (Figs. 1c and d). A mitochondrial DNA phylogeny also




Fig. 1. Mitochondrial DNA phylogenies under various scenarios of allopolyploid diversification of species A–E. Maternal lineages are solid lines, paternal lineages are dashed. Elimination of the dotted lines in the species relationships (left side) produces a topology identical to the mtDNA phylogeny (right side). Species are diploid, tetraploid ($4N$), octoploid ($8N$) or dodecaploid ($12N$). (a) Allopolyploid speciation generates tetraploid species C from a female ancestor of species B and a male ancestor of species D. Only half of the ancestry of species C is reflected by the mtDNA phylogeny. (b) Evolution of a dodecaploid species from tetraploid ancestors. In this example, only a third of the ancestry of the dodecaploid is reflected by the mtDNA phylogeny. (c) Multiple contemporaneous episodes of allopolyploidization could generate a phylogeny with a poorly resolved node or a hard polytomy. (d) Extinction could result in mtDNA monophyly of extant polyploids with different paternal ancestors.

offers insight into phylogeographic patterns of differentiation, irrespective of ploidy level, over the large range of this group.

1.3. Goals

To better understand the evolution of African clawed frogs here we: (1) estimate phylogenetic relationships among mitochondrial DNA (mtDNA) of all known and some undescribed species, (2) elucidate phylogeographic patterns of this group over sub-Saharan Africa, (3) use a

molecular clock calibrated by geology and fossil pipids to estimate the age of mtDNA clades of clawed frogs, and (4) evaluate the minimum number of polyploidization events needed to account for the mtDNA phylogeny.

2. Methods

2.1. Genetic samples

We sequenced a broad taxonomic sample of 52 in-group individuals and 1 outgroup individual, including all known species of clawed frog and intraspecific samples from multiple localities (Fig. 2 and Table 1). Although some of the recognized subspecies of *X. laevis* may merit full species status (Kobel et al., 1996; Measey and Channing, 2003), we treated them all as subspecies of *X. laevis*.

Most samples in this study were obtained from the live collection at the Institute of Zoology at the University of Geneva (Table 1). These individuals were collected and identified by B. Colombelli, L. Du Pasquier, M. Fischberg, H. Kobel, C. Loumont, J. Perret, D. Rungger, E. Rungger-Brandle, C. Thiebaud, J. Tymowska, and R.C.T. Species identifications were verified using characters compiled in Tinsley and Kobel (1996). Additional samples were collected by B.J.E., R.C.T., D.B.K., D.C.C., M. Picker, and M. Tobias, and obtained from the tissue archives of CAS, AMNH, TNHC, and ROM (museum abbreviations follow Leviton et al., 1985). Sequences from other pipoids (*Pipa pipa*, *Pipa parva*, *Hymenochirus* sp., and *Rhinophrynus dorsalis*) were included, and a sequence from a pelobatid frog, *Scaphiopus hurterii* was used as an outgroup (Table 1). We also analyzed a portion of the complete mtDNA sequence of *S. tropicalis* published online by the Joint Genome Institute (genome.jgi-psf.org/xenopus0/X.tropicalisMtGenome.gb). Because this sequence was identical to our sequence of *S. tropicalis* from Nigeria, we did not include it in subsequent analyses.

2.2. Ploidy

For many species, ploidy level was ascertained previously by karyotyping or measuring DNA content of related individuals (e.g., Thiebaud and Fischberg, 1977; Tymowska, 1991). To assess ploidy level of all samples in this study, alleles of the nuclear gene RAG-1 were co-amplified with PCR, cloned, and independently sequenced (B.J.E., unpublished data). Because it was not possible to obtain a karyotype or to measure DNA content from two ethanol preserved tissues (*S. cf. tropicalis* and *S. new tetraploid 2*), we inferred the ploidy level of these individuals solely by cloning and counting the number of divergent RAG-1 alleles. We view these




Fig. 2. Sample locations of *Xenopus* and *Silurana*. Information on distributional ranges of each species can be found in Tinsley et al. (1996).

ploidy levels as provisional until karyotypes of other conspecific individuals can be performed.

2.3. Molecular data

Data for this study include 2335 bp of DNA (including gaps that were introduced through alignment but not including regions of ambiguous homology that were excluded from analysis) from portions of the 12S gene, tRNA^{val}, and 16S rDNA. The gene order of the sequenced region (5′–3′) is 12S rDNA, tRNA^{val}, and 16S rDNA. These sequences include the 12S rDNA gene beginning approximately 106 bp after the 5′ start of the gene, the entire tRNA^{val} gene, and most of the 16S rDNA gene ending about 246 bp before the 3′ end. This region is homologous to positions 2816–5056 in the *Rana nigromaculata* GenBank Accession No. AB043889. To sequence this region, we used ABI Prism 3100 and 3730XL automated capillary sequencers and primers described elsewhere (Evans et al., 2003). Alignment was done with ClustalX (Thompson et al., 1997) and then adjusted by eye using MacClade, version 4.06 (Maddison and Maddison, 2000). Autapomorphies such as single site

insertions in a species were verified by examining the chromatograms. Alignment was generally unambiguous with the exception of two regions that were excluded: one region of ~13 bp begins about 80 bp upstream of the 3′ end of the 12S gene; the other region of ~10 bp begins at base 18 in the 16S gene. These data have been submitted to GenBank (Table 2).

2.4. Phylogenetic analyses

Phylogenetic analyses used maximum parsimony (MP), maximum likelihood (ML), and Bayesian methods. For MP analysis we performed a heuristic search for the most parsimonious tree with 1000 replicates of random taxon-addition and TBR branch swapping, with gaps treated as missing characters, using PAUP* version 4.0 (Swofford, 2002). Non-parametric bootstrap values were obtained with 2000 replicates, each with a single replicate of random taxon-addition. Decay index values were calculated for each node with Autodecay, version 4.0.2 (Eriksson, 1999).

For ML and Bayesian analysis, we used Modeltest version 3.06 (Posada and Crandall, 1998) to evaluate

Table 1
Samples used in this study

Species	Museum catalog	Country of origin	Specific locality	Coordinates (lat long)		GenBank
Outgroup						
<i>Scaphiopus huerterii</i>	TNHC 61954	United States	Aransas County, Texas	28.05	−97.05	AY581619
Silurana						
<i>S. tropicalis</i> *	—	Nigeria	Uyere, near Benin	6.72	5.77	AY581668
<i>S. tropicalis</i> *	—	Nigeria	Unknown	6.72	5.77	AY581669
<i>S. tropicalis</i> *	MHNG 2644.55	Sierra Leone	Near Freetown	8.50	−13.24	AY581670
<i>S. tropicalis</i> *	—	Ivory Coast	Adiopo Doume	5.32	−4.12	AY581671
<i>S. cf. tropicalis</i>	ROM 19161	Liberia	Sinoe Province, Sapo National Park, Gbaborne Creek	5.01	−9.04	AY581666
<i>S. epitropicalis</i> *	MHNG 2644.56	DRC	Kinshasa	−4.30	15.30	AY581665
<i>S. new tetraploid 1</i>	—	Republic of the Congo (Brazzaville)	Malemba	−5.83	12.57	AY581660
<i>S. new tetraploid 1</i>	—	Cameroon	Longyi	2.95	9.92	AY581661
<i>S. new tetraploid 1</i>	MHNG 2644.58	Cameroon	Nkoemvone	2.88	11.15	AY581664
<i>S. new tetraploid 1</i>	—	Gabon	Makokou	0.57	12.87	AY581662
<i>S. new tetraploid 1</i>	—	Gabon	Cap Esterias, near Libreville	0.37	9.43	AY581663
<i>S. new tetraploid 2</i>	CAS 207759	Equatorial Guinea	Bioko Island, Arena Blanca Rd.	3.53	8.58	AY581667
Xenopus						
<i>X. amieti</i>	MHNG 2644.54	Cameroon	Galim (type locality)	5.63	10.33	AY581634
<i>X. cf. fraseri 1</i>	CAS 207765	Equatorial Guinea	Bioko Id. Arena Blanca Rd.	3.53	8.58	AY581632
<i>X. cf. fraseri 2*</i>	—	Cameroon	Yaounde	3.87	11.52	AY581631
<i>X. new tetraploid</i>	AMNH 158377	Chad	Prifecture du Borkou-Ennedi-Tibesti, Sub-Prifecture Ennedi, Guelta d'Archei	16.91	21.77	AY581651
<i>X. new tetraploid*</i>	MHNG 2644.60	Nigeria	near Jos	8.7–9.0	11.2–12.0	AY581652
<i>X. andrei</i>	MHNG 2644.51	Cameroon	Longyi (type locality)	2.95	9.91	AY581627
<i>X. borealis</i> *	—	Kenya	Marsabit	2.33	38.01	AY581653
<i>X. borealis</i> *	—	Kenya	Kiambu	−1.17	38.01	AY581654
<i>X. borealis</i> *	MHNG 2644.64	Kenya	Samburu Range	0.66	37.50	AY581656
<i>X. borealis</i> *	—	Kenya	Maralal	1.08	36.68	AY581655
<i>X. boumbaensis</i>	MHNG, uncataloged	Cameroon	Yaounde	3.87	11.52	AY581635
<i>X. boumbaensis</i>	MHNG 2644.57	Cameroon	Moloundou (type locality)	2.02	15.13	AY581633
<i>X. clivii</i> *	MHNG 2644.50	Ethiopia	Near Addis Ababa	9.02	38.70	AY581637
<i>X. gilli</i>	—	South Africa	Cape Province, Cape Point Peninsula, Cape Point Nature Reserve	−34.25	18.41	AY581649
<i>X. gilli</i>	—	South Africa	Cape Province, Betty's Bay	−34.37	18.92	AY581650
<i>X. laevis</i> *	MHNG 2644.61	Malawi	Near Blantyre	−15.78	35.00	AY581648
<i>X. laevis</i> *	MHNG 2644.67	Republic of the Congo (Brazzaville)	Between Loubono and Pointe Noire near town of Koukila	−4.30	12.40	AY581638
<i>X. laevis</i> *	—	South Africa	Cape Province, Cape Point Peninsula, Lewis Gay Dam	−34.20	18.40	AY581639
<i>X. laevis</i> *	—	Botswana	Okavango	−18.97	22.57	AY581642
<i>X. laevis</i> *	MHNG 2644.53	Zambia	Near Lusaka	−15.50	28.20	AY581644
<i>X. laevis</i> *	—	Cameroon	Ngaoundere	7.32	13.58	AY581640
<i>X. laevis</i> *	MHNG 2644.52	Nigeria	Jos	9.90	8.90	AY581641
<i>X. laevis</i> *	CAS 168711	Tanzania	Tanga Region, Muheza Dist., East Usambara Mts, Amani, Amani Pond	−5.07	38.72	AY581647
<i>X. laevis</i> *	—	Rwanda	Shama	−1.95	30.06	AY581645
<i>X. laevis</i> *	—	Uganda	Kitanga	−1.12	30.05	AY581646
<i>X. largeni</i> *	MHNG 2644.59	Ethiopia	Sidamo Province, 400 km S of Addis Ababa, near Kibre Mengist	5.60	38.90	AY581643
<i>X. longipes</i>	AMNH A168447	Cameroon	Lake Oku (type locality)	6.21	10.47	AY581625
<i>X. muelleri</i> *	—	Swaziland	Nkambeni area	−26.05	31.66	AY581657
<i>X. muelleri</i> *	—	Malawi	Near Blantyre	−15.78	35.00	AY581658
<i>X. muelleri</i> *	MHNG 2644.63	Tanzania	Ifakara	−8.13	36.68	AY581659

Table 1 (continued)

Species	Museum catalog	Country of origin	Specific locality	Coordinates (lat long)		GenBank
<i>X. pygmaeus</i> *	—	DRC	Boende	-0.25	20.60	AY581626
<i>X. ruwenzoriensis</i>	—	Uganda	Rain forest near Bundibugyo in the Semliki Valley at the foot of the Ruwenzori Mountains (type locality)	0.70	30.00	AY581624
<i>X. vestitus</i>	—	Uganda	Lake Mutanda, near Kisoro	-1.22	29.71	AY581636
<i>X. wittei</i> *	CAS 201664	Uganda	Rukungiri Dist., Bwindi Impenetrable National Park, Kabale-Kayonza Rd.	-0.98	29.69	AY581629
<i>X. wittei</i> *	—	Rwanda	Cyamudongo Forest, Nyakabuye	-2.57	28.98	AY581628
<i>X. wittei</i> *	MHNG 2644.62	Uganda	Chelima Forest, near Echuya and Mulehe (type locality)	-1.06	29.90	AY581630
Other pipoids						
<i>Rhinophrymus dorsalis</i>	LACM 129913	Mexico	24 mi. WNW Coyuca de Benitez, 0.6 mi. NW El Papayo	17.03	-100.07	AY581620
<i>Hymenochirus</i> sp.	—	Africa, country unknown	Unknown	—	—	AY581623
<i>Pipa parva</i>	—	Venezuela	Merida	8.60	-71.15	AY581622
<i>Pipa pipa</i>	KU 205801	Peru	Madre de Dios, Cusco Amazonico, 15 km E of Puerto Maldonado	-12.60	-70.08	AY581621

There is no voucher for some samples and GPS coordinates (in decimal degree format) of some localities are approximate. Taxa demarcated by an asterisk were analyzed by Kobel et al. (1998).

Table 2

Estimated divergence times of nodes in millions of years and 95% confidence interval (min–max)

Node	Age	95% confidence
Pipidae	129.8	114.5–146.1
Pipinae	112.0	(Fixed)
Xenopodinae	63.7	50.4–81.3
<i>Pipa</i>	55.7	46.5–65.7
<i>Xenopus</i>	41.6	31.8–54.6
<i>Silurana</i>	17.6	12.1–24.8
Clade A	32.3	22.7–45.3
Clade B	19.4	13.7–25.2
Clade C	12.0	7.8–17.7
MRCMA <i>X. longipes</i> , <i>X. ruwenzoriensis</i>	6.0	2.8–8.3
MRCMA <i>X. longipes</i> , <i>X. boumbaensis</i>	4.8	1.9–6.7
MRCMA <i>X. longipes</i> , <i>X. amieti</i>	3.7	0.9–15.0
MRCMA <i>X. pygmaeus</i> , <i>X. cf. fraseri 1</i>	6.8	4.2–10.5
MRCMA <i>X. pygmaeus</i> , <i>X. wittei</i>	10.0	2.3–13.8
MRCMA <i>X. pygmaeus</i> , <i>X. andrei</i>	9.1	6.0–13.4
MRCMA <i>X. largeni</i> , <i>X. pygmaeus</i>	20.1	13.3–30.1
MRCMA <i>X. vestitus</i> , <i>X. pygmaeus</i>	25.5	17.7–36.1
MRCMA <i>X. laevis</i> , <i>X. gilli</i>	16.7	9.6–26.2
MRCMA <i>S. epitropicalis</i> , <i>S. new tetraploid 1</i>	7.7	4.3–12.0
MRCMA <i>S. tropicalis</i> , <i>S. cf. tropicalis</i>	13.8	8.9–20.3
<i>S. new tetraploid 1</i>	3.2	0.9–6.3
<i>X. laevis</i>	11.2	6.0–19.3
<i>X. gilli</i>	8.5	4.8–13.4
<i>X. wittei</i>	2.7	1.2–4.6

Age of the root (Pipoidea) is not shown because of inaccuracy of method in determining age of root. Some nodes are referred to as the most recent common maternal ancestor (MRCMA) of two species. The average rate of evolution was 0.00249 substitutions per site per million years, $s = 0.00029$, with a range of 0.00184–0.00299.

different models of evolution. Maximum likelihood analysis was performed with PAUP* (Swofford, 2002) under the model selected by Modeltest (GTR + I + Γ , see below), with two rounds of successive approximation, each with five replications of random taxon-addition. Initial parameters were set to those estimated from a neighbor-joining topology, and then re-estimated from the resulting topology and set to these new values for the second iteration. Only two rounds of successive approximation were performed because the topology recovered from both rounds was identical.

Bayesian analysis was performed with MrBayes, version 3.0b4 (Huelsenbeck and Ronquist, 2001), under the same model as the ML analysis. We ran four Metropolis coupled Markov chain Monte Carlo analyses starting with random trees for each of four simultaneous chains, with flat Dirichlet prior distributions set to 1.0 for each rate substitution type, and the differential heating parameter set to 0.2. Four runs were performed, each for 5,000,000 generations. Post-run analysis suggested that parameter estimates reached stationarity well before 200,000 generations; trees recovered from these initial rounds were discarded as burn-in. The joint posterior probability distributions and parameter estimates of each of the independent runs were similar, suggesting that the chains were run for a sufficient number of generations to sample the posterior probability landscape adequately. Posterior probabilities were calculated from post-burn-in trees from all runs.

2.5. Molecular clock

In order to estimate divergence times from molecular data, we used a likelihood-ratio test (Goldman, 1993) to determine whether the likelihood of a tree without a molecular clock enforced was significantly greater than one with a molecular clock enforced under the GTR + I + Γ model of evolution, in which degrees of freedom are equal to the number of terminals–2 (Huelsenbeck and Rannala, 1997). Because the data did not conform to expectations of clock-like evolution (see below), we used a semi-parametric method to estimate divergence times of various clades using r8s, version 1.60 (Sanderson, 1997, 2002). This approach, penalized likelihood, permits a different rate of evolution on every branch of the tree, but imposes a cost on the model if rates among branches change quickly.

A first step in estimating divergence times with r8s is to use a cross-validation procedure to empirically test various values for a smoothing parameter that determines the relative importance of the independent rates of each branch versus the cost of big changes in rates among branches (Sanderson, 1997, 2002). Larger smoothing parameters are appropriate for phylogenies that more closely approximate clock-like behavior. Using the TN algorithm, we tested a range of smoothing parameter values from 1 to about 3100. A value of 10 had the lowest error interval and the smoothing parameter was set to this value for analysis of divergence times. *Scaphiopus hurterii* was used to root the phylogeny with branch lengths recovered from the likelihood analysis. This taxon was pruned from the tree for analysis of divergence times.

Under the assumption that pipid ancestors did not cross marine barriers, the rifting of Africa and South America is a likely source of vicariance that initiated the diversification of the Pipinae subfamily, leading to the New World genus *Pipa* and the African genera *Hymenochirus* and *Pseudhymenochirus* (Báez, 1996). The complete separation of Africa and South America by ocean is dated in the late Aptian about 105–119 million years ago (Maisey, 2000; McLoughlin, 2001). To calibrate a molecular clock, the divergence time of Pipinae was fixed at 112 million years ago (mya). If the Pipinae diversified prior to this time, the estimated divergence times will underestimate the true divergence times.

The extinct species *Xenopus arabiensis*, represented by fossils from the late Oligocene of Yemen (23.8–28.5 mya), has a number of morphological features that support its inclusion in the genus *Xenopus*, including a prominent cone-shaped prehallux, a characteristic of the *muelleri* subgroup (Henrici and Báez, 2001; Kobel, 1996b). This species also has been allied with *X. muelleri* on the basis of the presence of a dentate vomer, a condition that is known among extant pipids only in *X.*

muelleri (Estes, 1977), although this character state could be plesiomorphic (Henrici and Báez, 2001). We used the age of this species to provide a minimum age limit for the genus *Xenopus* of 23.8 mya.

It has been proposed that pipoid frogs are sister to all other Anura (Maglia et al., 2001; Púgner et al., 2003). While other information does not support this configuration (Biju and Bossuyt, 2003; Ford and Cannatella, 1993), to be conservative we also set a maximum age limit for the root to the age of the earliest known anuran fossil (*Prosalirus*), estimated at 195 mya (Shubin and Jenkins, 1995).

Confidence limits for divergence times were calculated by bootstrapping the data, generating a phylogram for each bootstrap dataset, estimating a smoothing parameter for each bootstrap phylogram with r8s, and then analyzing each phylogram with the appropriate smoothing parameter and other settings as described above (Sanderson, 1997). For these analyses, we used Perl scripts to make input files and to parse output files. Confidence limits (95%) were estimated as the mean estimated age of a node in all bootstrap replicates \pm 1.96 standard deviations of these estimates.

2.6. Phylogeographic analysis

Area cladograms were constructed from ML and MP topologies by changing the name of each sample to the location(s) of that species or, in *X. laevis*, the location of intraspecific clades (western, central, eastern, and/or southern Africa). These regions were delimited by three putative biogeographic barriers that may or may not have affected clawed frogs: the Rift Valley, the Dahomey Gap (where contemporary savannah habitat extends southward to the coast from Ghana, through Togo and Benin, to west Nigeria), and the southern limit of tropical rainforest. Western, central, and eastern Africa, have and/or had tropical rainforest in the past, whereas southern Africa was dominated by drier habitat during clawed frog evolution (Stokes et al., 1997). To better understand phylogeographic evolution of this group, we included the additional geographic information that *X. new tetraploid* is also known from western Africa, even though genetic material from this species was not available from this population (Tinsley et al., 1996). The distribution of *Hymenochirus* was set to western and central Africa and the distributions of *Pipa*, *Rhinophrynus*, and *Scaphiopus* were designated as New World. We also generated a chronogram from the ML topology using r8s.

We used dispersal–vicariance analysis to estimate the ancestral locations of various clades in all of the MP topologies, and in the ML topology, with DIVA, version 1.1 (Ronquist, 1996, 1997). The maximum number of ancestral areas was set to five (the total number of areas in the analysis), and ambiguous optimal solutions

were reported. This approach has the advantages that it does not assume a hierarchical (or any) area relationship, it permits an ancestor to occupy multiple areas, and it permits dispersal between areas without speciation (Ronquist, 1996, 1997). DIVA reconstructs ancestral areas by minimizing the number of dispersal and vicariance events needed to explain an extant distribution. Vicariance is the preferred (no cost) mode of speciation; dispersal and extinction each impose an equal cost on the optimization model. Most differences between the topologies recovered from each phylogenetic analysis did not affect reconstruction of ancestral areas because the topological differences occurred within a geographical region. One difference in the topology within *Silurana* in some of the MP trees affected reconstruction of ancestral areas within this genus, but not on other parts of the tree, and is thus not discussed.

2.7. Testing of phylogenetic hypotheses

To estimate the minimum number of polyploidization events that occurred in clawed frogs, we assume that genome duplication is unidirectional—that reversions to a lower ploidy level do not occur. Five phylogenetic hypotheses concerning the frequency of polyploidization were tested with parametric bootstrapping (Goldman et al., 2000; Hillis et al., 1996; Huelsenbeck et al., 1996). Hypothesis 1 postulates a single origin of octoploids. Because dodecaploids could potentially inherit mtDNA from a tetraploid ancestor rather than an octoploid ancestor (Fig. 1b), this hypothesis was tested with a backbone constraint in which dodecaploids were allowed to assume any phylogenetic position with respect to octoploids and tetraploids. Hypothesis 1 also leaves open the possibility that one or both dodecaploids were derived from the union of two hexaploid ($2N = 54$) genomes that are now extinct. Intermediate ploidy levels (triploid or hexaploid) may be precludes to allopolyploid speciation of clawed frogs (Kobel, 1996a). Hypothesis 2 postulates a single origin of dodecaploids and monophyly of dodecaploid mtDNA. Hypothesis 3 postulates a single origin of *Silurana* tetraploids and monophyly of these mtDNA sequences.

Phylogenetic analyses recovered a clade that contains all of the octoploids except *X. vestitus* and both dodecaploids (see below). However, ML and Bayesian analyses also nest the tetraploids (*X. pygmaeus* + *X. cf. fraseri* 1) in this clade, and MP analysis nests the tetraploid *X. cf. fraseri* 2 in this clade. To examine whether further episodes of polyploidization are supported, two hypotheses were tested that postulate mtDNA monophyly of both dodecaploids and monophyly of all octoploids except *X. vestitus*. The first one, Hypothesis 4, postulates monophyly of dodecaploid mtDNA within a clade containing all octoploid mtDNA except *X. vestitus*. The second, Hypothesis 5, postulates one clade

containing dodecaploid mtDNA and a separate clade containing all octoploid mtDNA except *X. vestitus*. Both of these hypotheses are consistent with two origins of octoploids and one origin of dodecaploids.

Additionally, we tested for mtDNA monophyly of each species group with more than one species. Hypotheses 6–9 postulate monophyly of the *laevis*, *muelleri*, *vestitus-wittei*, and *fraseri*-like groups, respectively.

For parametric bootstrapping, a heuristic search for the most parsimonious tree consistent with each biogeographic (null) hypothesis was performed using 100 replications of random taxon-addition. The set of most parsimonious trees consistent with each null hypothesis differed primarily in relationships among short terminal branches. Because of this, simulation on any of these trees should produce similar results. We ranked these trees under the K2P + Γ model of evolution and selected the most likely tree for data simulation. Using this tree, which is the null hypothesis of the parametric bootstrap test (Goldman et al., 2000; Huelsenbeck and Rannala, 1997), a hierarchical likelihood-ratio test was used to select a model of evolution for data simulation using Modeltest (GTR + I + Γ was independently selected for each test). One thousand datasets were simulated under this more complex model for each null hypothesis with Seq-Gen version 1.2.6 (Rambaut and Grassly, 1997). For each simulated dataset, we calculated the treelength difference in parsimony score between searches unconstrained and constrained for the null hypothesis. The Logreader program (D. Zwickl, unpublished) was used to parse the output files. The probability (P) of the null hypothesis is equal to the proportion of treelength differences from the simulated data that is equal to or more extreme than the observed treelength difference. We applied the sequential Bonferroni procedure to adjust a critical value of 0.05 for hypothesis rejection in multiple tests (Rice, 1989).

The posterior probability of each hypothesis was estimated by filtering post-burn-in trees from Bayesian analysis with a constraint tree (Huelsenbeck et al., 2000). We also evaluated the prior probability of each hypothesis (the expected probability of each hypothesis before data collection) by generating 400,000 random trees with PAUP and computing the proportion of these trees that satisfy each hypothesis.

3. Results

3.1. Phylogeny

Results from MP, ML, and Bayesian analyses were similar, with exceptions discussed below. Maximum parsimony searches recovered 55 equally parsimonious trees of 3355 steps (CI = 0.477, RI = 0.766). For model-based phylogeny estimation, the GTR + I + Γ model was




Fig. 3. Maximum likelihood tree. Number of chromosomes of each species is indicated in parentheses. Non-parametric MP bootstrap values and posterior probabilities are separated by a forward slash above branches, and decay indexes are below them. An asterisk indicates bootstrap values or posterior probabilities greater than 95, a dash indicates values below 50, and for clarity some bootstrap and decay values of terminal branches are omitted. Scale bars indicate the number of substitutions per site to the left and to the right of the dotted line. Clades A–C referred to in text are indicated; these clades are also in all MP trees. Initial parameters estimated for the GTR + I + Γ model on a neighbor-joining tree were: base frequencies (A C G) = (0.3861 0.2381 0.1588), rate matrix (A–C A–G A–T C–G C–T G–T) = (4.1564 11.1392 10.1821 0.4933 40.2440 1), shape parameter = 0.4155, and proportion of invariant sites = 0.3135. Parameters estimated on a likelihood tree after a successive approximation search were: base frequencies (A C G) = (0.3848 0.2374 0.1605), rate matrix (A–C A–G A–T C–G C–T G–T) = (4.2356 11.1240 10.4404 0.4606 41.4022 1), shape parameter = 0.4103, and proportion of invariant sites = 0.3089.

selected by the hierarchical likelihood-ratio test. The hypothesis that these sequences conformed to a molecular clock was rejected ($P < 0.001$). Maximum likelihood analysis and Bayesian analyses recovered very similar topologies (Fig. 3).

Most clades with high bootstrap values or high posterior probabilities are present in trees recovered from all analyses. These analyses all support mtDNA monophyly of *Silurana* with respect to *Xenopus*, and monophyly of *X. borealis*, *X. muelleri*, and *X. new tetraploid* (Clade B, Fig. 3) with respect to other *Xenopus* (Clade A, Fig. 3). In all analyses, within Clade A the sequence of *X. clivii* is sister to other sequences, sequences of *X. gilli* and *X. laevis* form a clade, and mtDNA of *X. pygmaeus*, *X. cf. fraseri* 1 and 2, all octoploids except *X. vestitus*, and both dodecaploids form a clade (Clade C, Fig. 3). All analyses support monophyly of sequences of *X. ruwenzoriensis*, *X. longipes*, *X. amieti*, and *X. boumbaensis*, and monophyly of mtDNA of *X. pygmaeus* and *X. cf. fraseri* 1. Within *X. laevis*, mtDNA from South Africa is the sister to all other intraspecific sequences, and mtDNA from Malawi is sister to mtDNA of *X. laevis* to the North and West (Cameroon, Nigeria, Botswana, Zambia, Rwanda, Uganda, and Tanzania). *X. laevis* mtDNA from western Africa (Cameroon and Nigeria) and the center of southern Africa (Botswana and Zambia) form a clade that is sister to *X. laevis* mtDNA from East Africa (Uganda, Rwanda, and Tanzania). Of the five species groups in *Xenopus*, the only one that includes a clade of mtDNA in our analyses is the monotypic *longipes* group.

In *Silurana*, mtDNA of two tetraploids, *X. epitropicalis* and *S. new tetraploid 1* form a clade. MtDNA of the diploid *S. tropicalis* and a putative new tetraploid, *S. new tetraploid 2*, form a clade (Fig. 3).

Topologies recovered from these analyses differ in some respects within Clades A and C, and within *Silurana* (Fig. 3). Within Clade A, the mtDNA relationship of *X. largeni* with respect to (*X. laevis* + *X. gilli*) is not well supported in any analysis. In the MP trees, *X. largeni* is sister to a clade containing *X. laevis*, *X. gilli*, and Clade C, but in ML and Bayesian analysis *X. largeni* is sister to Clade C only (Fig. 3).

In Clade C, MP places (*X. pygmaeus* + *X. cf. fraseri* 1) sister to a clade containing (*X. wittei*, *X. andrei*, *X. cf. fraseri* 2, *X. ruwenzoriensis*, *X. longipes*, *X. amieti*, and *X. boumbaensis*). Within this latter clade, *X. wittei* is sister to (*X. andrei*, *X. cf. fraseri* 2, *X. ruwenzoriensis*, *X. longipes*, *X. amieti*, and *X. boumbaensis*). *X. andrei* is sister to a clade within which *X. cf. fraseri* 2 is sister to (*X. ruwenzoriensis*, *X. longipes*, *X. boumbaensis*, and *X. amieti*).

In *Silurana*, 19 out of 55 MP trees ally *S. cf. tropicalis* with (*S. epitropicalis* + *S. new tetraploid 1*) and the other MP trees have the same topology in *Silurana* as the ML and Bayesian analysis (Fig. 3).

3.2. Divergence times, rates of evolution, and ancestral areas

The most recent common ancestor of *P. pipa* and *P. parva* is thought to be the ancestor of all extant species in this genus (Cannatella and Trueb, 1988a; Trueb and Cannatella, 1986). If this is the case, we estimate that all extant members of the genus *Pipa* diversified in the New World about 55.7 mya with a 95% confidence interval of 46.5–65.7 mya (Table 2).

Extant lineages of clawed frogs, the Xenopodinae, appear to have evolved in eastern and/or central equatorial Africa well after the split of South America from Africa (Fig. 4 and Table 2). Ancestral area reconstruction of *Xenopus* is eastern Africa and of *Silurana* is central Africa. The ancestor of (*X. laevis* + *X. gilli*) probably originated in southern Africa. All octoploids and dodecaploids, except possibly *X. vestitus*, originated in central Africa. Under the assumptions of no extinction of ancestral tetraploids (Fig. 1d) and comprehensive sampling of extant species, all age estimates for octoploid and dodecaploid lineages in this study predate the Pleistocene. Confidence intervals of *X. amieti*, *X. longipes*, *X. boumbaensis*, and *X. wittei* have some overlap with the Pleistocene.

3.3. Hypothesis testing

Our analyses reject the hypothesis that octoploid clawed frogs evolved from only one polyploidization event. A parametric bootstrap test rejected Hypothesis 1 ($P < 0.001$) and the prior and posterior probabilities are near 0.0. A search for the most parsimonious trees consistent with Hypothesis 1 recovered 57 MP trees of 3395 steps in length as compared to the unconstrained tree length of 3355.

A parametric bootstrap test did not reject Hypothesis 2, a single origin of dodecaploids, ($P = 0.955$). A search for the most parsimonious trees consistent with monophyly of dodecaploid mtDNA, recovered 149 MP trees of length 3357 steps, two steps longer than the unconstrained tree. The prior probability of this hypothesis was 0.01025 and the posterior probability was 0.25313.

The hypothesis of a single origin of *Silurana* tetraploids, Hypothesis 3, was rejected ($P < 0.001$) and the prior and posterior probabilities of this hypothesis are near 0.0. A search for the most parsimonious tree consistent with this hypothesis recovered 54 MP trees of length 3372, 17 steps longer than the unconstrained trees.

Under a scenario of two origins of octoploids and one origin of dodecaploids, dodecaploids could have arisen from an octoploid maternal ancestor and have mtDNA that is nested in a clade containing only octoploid mtDNA, or not, and therefore have a clade of mtDNA that is not nested within octoploid mtDNA. Both of these




Fig. 4. Timing, location, and polyploidization of clawed frog lineages. Branch lengths are proportional to divergence time as estimated from the ML topology and a relaxed molecular clock. Letters associated with branches refer to reconstructed ancestral locations in Africa (W, C, E, and/or S), the New World (NW) or, for nodes older than 112 mya, to Gondwana (G1 and G2). Reconstructed ancestral areas in Gondwana do not include southern Africa (G1) or do not include southern and eastern Africa (G2). The large map of Africa depicts the ancestral locations and the distribution of rainforest during the Pleistocene glacial maxima in black and at present in gray (eastern forests not shown). Small maps depict the distribution of *Silurana*, and species in *Xenopus* Clades A and B in Fig. 3. Arrows on the time scale indicate the date that Africa split from South America 112 mya, and the beginning of the Pleistocene, 1.8 mya. Six polyploidization events from 18 to 36 chromosomes, 36 to 72 chromosomes, 36 plus 72 to 108 chromosomes, and 20 to 40 chromosomes are indicated. Two of these events are depicted on the small topology that corresponds with the constraints of Hypothesis 4; this hypothesis was not rejected by parametric bootstrapping. A strict interpretation of larger topology suggests that additional polyploidization events occurred.

scenarios were explored (Hypotheses 4 and 5, respectively). Hypothesis 4 was not rejected after Bonferroni correction ($P = 0.048$, adjusted critical value = 0.0167, $k=9$ tests; Rice, 1989) but Hypothesis 5 was rejected ($P < 0.001$). One hundred and eighty MP trees of length

3359 steps were consistent with Hypothesis 4 and 54 MP trees of length 3370 steps were consistent with Hypothesis 5. These sets of trees are 4 and 15 steps longer than the unconstrained trees, respectively. The prior probabilities of Hypotheses 4 and 5 are about 0.0 and the posterior

probabilities are 0.02644 and 0.0000, respectively. Hypothesis 4 is consistent with a scenario in which octoploids originated twice and dodecaploids originated once from an octoploid maternal (mtDNA contributing) ancestor. It does not support the hypothesis that dodecaploids originated from a tetraploid maternal ancestor (Fig. 1b).

Hypotheses 6–8, monophyly of the *laevis*, *muelleri*, and *vestitus-wittei* groups, were rejected ($P = 0.007$, adjusted critical value = 0.0125, $k = 9$ tests; Rice, 1989; $P < 0.001$; and $P < 0.001$, respectively). The set of most parsimonious trees consistent with these hypotheses were 7, 18, and 36 steps longer, respectively, than the unconstrained tree. The prior probabilities of Hypotheses 6–8 are close to 0.0 and the posterior probabilities are 0.0019, 0.0062, and about 0.0.

A parametric bootstrap test did not reject Hypothesis 9, which postulates monophyly of the *fraseri*-like group ($P = 0.556$). The most parsimonious tree consistent with this hypothesis was 3356 steps (as compared to an unconstrained length of 3355 steps). The prior probability of this hypothesis is close to 0.0 and the posterior probability is 0.0172.

4. Discussion

4.1. Phylogenetic relationships

The monophyly of *Xenopus + Silurana* is extremely well supported by this analysis, in accord with other studies (Cannatella and de Sá, 1993; de Sá and Hillis, 1990; Trueb and Báez, 1997), but not Cannatella and Trueb (1988a), who placed *Silurana* as the sister taxon of Pipinae. This last result was reconsidered by Cannatella and de Sá (1993), and there is consensus that the

morphological data also supports monophyly of *Xenopus + Silurana*.

Monophyly of both *Silurana* and of *Xenopus* is consistent with results of other molecular and morphological studies that include fossil taxa (Cannatella and de Sá, 1993; Kobel et al., 1998; Trueb and Báez, 1997). It should be recognized that given our results, the choice between a taxonomy that recognizes only *Xenopus* for clawed frogs versus *Xenopus* and *Silurana* is a matter of emphasis rather than science. Although recognition of *Silurana* as a separate genus is not mandatory given strong support for monophyly of all clawed frogs, use of both names underscores trenchant biological and historical differences between the two clades. Each of these lineages is older than most families in the order Primates, for instance (Table 2; Yoder and Yang, 2004), and fixed chromosomal and morphological differences are present (Trueb and Báez, 1997).

Topologies recovered in this analysis correspond almost exactly to that detailed by Kobel et al. (1998) based on a smaller portion of mitochondrial rDNA and fewer species. Close relationships between *X. muelleri* and *X. borealis* and between *X. laevis* and *X. gilli* reported in our study agree with results from electrophoretic studies of sperm and serum proteins (Graf and Fischberg, 1986; Mann et al., 1982). MtDNA and nuclear DNA studies also both suggest a close relationship between *X. ruwenzoriensis* and *X. amieti* (Bürki and Fischberg, 1985; Graf and Fischberg, 1986; Mann et al., 1982)—species that have similar male vocalizations (Kobel et al., 1996).

However, few consistencies beyond monophyly of *X. laevis* emerge from comparisons to the unrooted mtDNA networks of Carr et al. (1987), or a reanalysis of those data (Cannatella and de Sá, 1993). To test whether inaccuracy in mapped restriction sites could explain this

Table 3

Possible mtDNA-donating most recent common maternal ancestor (MRCMA) of polyploid clawed frog species, assuming no extinction of ancestral mtDNA lineages

Species	MRCMA (posterior probability)	Contribution of MRCMA to descendant allopolyploid nuclear genome(s)
All <i>Xenopus</i> tetraploids	A diploid ($2N = 18$) MRCMA of all <i>Xenopus</i> (100%)	At least 50%
<i>Silurana</i> epitropicalis + <i>S. new</i> tetraploid 1	A diploid ($2N = 20$) MRCMA of all <i>Silurana</i> (100%).	at least 50%
<i>X. vestitus</i>	A tetraploid MRCMA of <i>X. vestitus</i> , <i>X. largeni</i> , <i>X. laevis</i> , <i>X. gilli</i> , <i>X. pygmaeus</i> , <i>X. cf. fraseri</i> 1 and 2, all other octoploids, and both dodecaploids (76%), or a tetraploid MRCMA of <i>X. vestitus</i> and <i>X. largeni</i> (11%)	At least 50%
Octoploids other than <i>X. vestitus</i>	A tetraploid MRCMA of <i>X. pygmaeus</i> and <i>X. fraseri</i> 1 and 2, both dodecaploids, and all octoploids except <i>X. vestitus</i> (100%)	At least 50%
<i>X. ruwenzoriensis</i>	An octoploid MRCMA of <i>X. ruwenzoriensis</i> , <i>X. longipes</i> , <i>X. amieti</i> , <i>X. boumbaensis</i> (100%), or, possibly more specifically, an octoploid MRCMA of <i>X. ruwenzoriensis</i> and <i>X. longipes</i> only (25%)	66.7%
<i>X. longipes</i>	An octoploid MRCMA of <i>X. longipes</i> , <i>X. ruwenzoriensis</i> , <i>X. amieti</i> , <i>X. boumbaensis</i> (100%), or, possibly more specifically, an octoploid MRCMA of <i>X. amieti</i> , <i>X. boumbaensis</i> , and <i>X. longipes</i> (70%), or an octoploid MRCMA of <i>X. amieti</i> and <i>X. longipes</i> only (67%), or an octoploid MRCMA of <i>X. ruwenzoriensis</i> and <i>X. longipes</i> (25%).	66.7%

difference, we examined whether the location of restriction sites inferred by Carr et al. (1987) corresponds to the locations of these sites as predicted by our sequence data. For the few comparisons that were possible, there was excellent congruence. For example, both datasets suggest two *SacII* sites are fixed across all taxa, and a *PvuII* site is present only in *X. ruwenzoriensis*. Our sequences predict more *EcoRI* positions (AATT) than were mapped by Carr et al. (1987), and suggest that *EcoRI** sites (GAATTC) were actually mapped. Our predictions for *EcoRI** positions relative to the first fixed *SacII* site correspond exactly to the map of Carr et al. (1987). Thus we dismiss differences in data quality as a source of differences in phylogeny estimation, and attribute the increased phylogenetic resolution in our study to increased sampling of taxa and polymorphic characters (Pollock et al., 2002; Zwickl and Hillis, 2002).

This mtDNA phylogeny also is not consistent with the *laevis*, *muelleri*, or *vestitus-wittei* species groups. These discrepancies could stem from the incomplete picture of allopolyploid evolution that is gleaned from the former (Fig. 1). For example, mtDNA paraphyly of each species group could be due to the fact that group members do not share a common maternal ancestor (that contributes mtDNA) but do share a common paternal one. Alternately, the paternal ancestor of some species could be the maternal ancestor of the others. A portion of the recent shared ancestry of some species, however, can be discerned from the mtDNA phylogeny (Table 3), barring recent extinction of ancestral mtDNA lineages (Fig. 1d). Paraphyly of mtDNA in species groups also could occur if characters used to define species groups are symplesiomorphic rather than synapomorphic.

4.2. Time and place of diversification

It is useful to examine the degree to which molecular estimates of the time and place of diversification are consistent with other sources of information from fossils, climatic history, geological history, and other molecules, while also avoiding circularity when some of this information was used to make the molecular estimates. Estimation of divergence times by a molecular clock is confounded by error in phylogenetic estimation and taxonomic identification of fossils, and by error and uncertainty in the age of geological events and fossils (Graur and Martin, 2004). Incomplete information in the fossil record and in the molecular phylogeny (which does not contain fossil taxa) also poses a challenge. Error in estimating ancestral distributions is expected to increase with an increasing age of a clade and this error could be compounded by potentially high dispersability of clawed frogs, some of which have close mtDNA relationships that span large geographic distances. Under the assumption of a correlation between time and mo-

lecular divergence (that is not necessarily clock-like), the wide dispersability of some species is evidenced by, for example, mtDNA of *X. laevis* from Nigeria and Botswana, separated by over 3500 km, that is only 0.5% divergent.

Living pipoid frogs (Rhinophrynidae and Pipidae) are distributed in North, Central, and South America, and Africa, and their ancestors were widely distributed before the breakup of Gondwana. Close relatives of the Pipidae are known from Early Cretaceous deposits of Israel (Trueb, 1999), and Middle Cretaceous strata of Argentina (Báez et al., 2000), and the age of pipoid origin is at least Late Jurassic (150 mya; Henrici, 1998). A proposed sister relationship between (*Hymenochirus* + *Pseudhymenochirus*) and the extinct African species *Pachybatrachus taqueti* (83.5–89.0 mya; Báez and Rage, 1998) is consistent with the hypothesis that Gondwana breakup led to the diversification of Pipinae.

We estimate that the ancestor of extant *Xenopus* originated between 31.8 and 54.6 mya and that all extant clawed frogs originated between 50.5 and 81.3 mya (Table 2). These time intervals predate fossils of the extinct species *Xenopus arabiensis*, a putative sister taxon of *X. muelleri*, which were deposited in the Late Oligocene of Yemen (28.5–23.8 mya; Henrici and Báez, 2001), but are well after the breakup of Gondwana about 112 mya. Other studies that assumed clock-like divergence of mtDNA sequences (Kobel et al., 1998), duplicate genes (Hughes and Hughes, 1993), or serum albumins (Bisbee et al., 1977), estimate that *Xenopus* is even younger. However, Estes (1975b) detailed similarities between an extinct fossil species “*Xenopus*” *romeri* from the late Paleocene of Brazil (61–54.8 mya; Estes et al., 1978) and fossils similar to *Xenopus* (= *Silurana*) *tropicalis* from Niger, implying that the ancestor of Xenopodinae existed before the separation of Africa and South America. Cannatella and de Sá (1993) placed “*Xenopus*” *romeri* as sister to either *Silurana* or Pipinae. A more detailed phylogenetic analysis by Trueb and Báez (1997) placed “*Xenopus*” *romeri* as sister to extant clawed frogs (*Xenopus* + *Silurana*). If “*Xenopus*” *romeri* is not nested within the Xenopodinae, it is plausible that cladogenesis of the most recent common ancestor of these taxa occurred before the breakup of Gondwana. Under the assumption that the breakup of Gondwana did lead to the divergence of *Pipa* from (*Hymenochirus* + *Pseudhymenochirus*), our age estimates (calibrated by this event) cast further doubt on the relationship between South American and African species proposed by Estes (1975a) because the most recent common ancestor of extant clawed frogs is younger than the formation of the Atlantic Ocean.

Similarly, Cannatella and de Sá (1993) addressed the implications of Estes’ (1975a) synonymy of the Argentine fossil *Shelania pascuali* (Paleogene, 47.2–43.4 mya) into *Xenopus*. They showed that this placement intro-

duced an inconsistency, because the split between Argentine and African *Xenopus* would then have to be older than the Atlantic Ocean. The conundrum was resolved by a thorough re-analysis by Trueb and Báez (1997), in which *Shelania pascuali* was shown to be outside of Xenopodinae.

The following integration of postulated dates with geological and paleoclimatological information is necessarily speculative. We do not test hypotheses, but rather point to consistencies between these sources of data. This scenario, coupled with future discoveries, may provide testable predictions.

Reconstruction of ancestral distributions suggests that the most recent common ancestor of extant *Xenopus* is from eastern equatorial Africa and that of *Silurana* is from central equatorial Africa; all clawed frogs may be derived from an ancestor that occupied either one or both of these areas (Fig. 4). Fifty million years ago, Africa was south of its current position and the equatorial rain forest was located on land that is now the Sahara desert (Lovett, 1993). Ancient biogeographic patterns of clawed frogs are thus best conceptualized in terms global location relative to the equator, rather than in terms of a fixed position on the African continent.

The extent of moist and dry habitats also varied on Africa. Clawed frogs may have become widespread in the Middle East and Africa when the Sahara region was more humid (Henrici and Báez, 2001). As Africa drifted northward, the central African plateau was uplifted (Tertiary; 65–23 mya), and the Rift Valley opened first in northern Africa about 30–20 mya, then in eastern Africa by 15–12 mya, and in southern Africa by the Pliocene (5–1.8 mya; Griffiths, 1993). During this time the equatorial rainforest was fragmented by savannah habitat into the expansive western/central rainforest and the patchy eastern rainforest (Lovett, 1993). Some divergent species such as *X. clivii* and *X. largeni* may have had wider ranges before the Ethiopian highlands became isolated from other habitat by desert. A similar explanation might account for the disjunct populations of *X.* new tetraploid in Chad and Nigeria, but the relatively low divergence between mtDNA sequences from these regions (0.9% patristic distance, Fig. 3) suggests more recent vicariance or dispersal is a better explanation.

During the Pleistocene, repeated changes in aridity led to a retraction of rain forest to small refugia and an expansion of savannah and desert habitat (Fig. 4; Nichol, 1999; Stokes et al., 1997). These changes may have affected the distributions of forest-adapted species, such as those in the *fraseri*-like group (Kobel, 1996a). The diverse topology and environmental conditions of the Rift Valley and eastern Congo may have promoted interaction and hybridization among species that are adapted to different habitats and elevation (Tinsley et al., 1996). Interestingly, octoploid and dodecaploid species of clawed frogs are known only from areas in

Central Africa that had refugia of tropical rainforest during the Pleistocene, although these lineages probably originated before this period (Fig. 4 and Table 2). That octoploids and dodecaploids (which are thought to be allopolyploid) are endemic to, but older than, Pleistocene rainforest refugia suggests that these regions acted as “lifeboats” for some species to endure the vagaries of climatic change, and possibly as platforms from which lineages were able to disperse and diversify when suitable habitat became more widespread. This does not support the hypothesis that rainforest refugia acted as “species pumps” that spawned new successful lineages (proposed for the Neotropics, Haffer, 1969; but see Knapp and Mallet, 2003) because the diversity within these areas predates the age of Pleistocene refugia.

We estimate that *X. laevis* originated in Southern Africa about 6.0–19.3 mya (Fig. 4 and Table 2), although the sister relationship between *X. laevis* mtDNA from South Africa and other *X. laevis* mtDNA is not strongly supported (Fig. 3). *Xenopus l. laevis* also has separate mtDNA clades in the southwestern and eastern parts of South Africa (Grohovaz et al., 1996; Measey and Channing, 2003). Regional differences in morphology and vocal characteristics in this species may warrant taxonomic revision (Kobel et al., 1996; Measey and Channing, 2003).

Marine barriers may pose formidable obstacles to dispersal of clawed frogs. For example, populations of *X. gilli* that are separated by only 100 km have 3.6% patristic divergence (Fig. 3). This genetic difference is virtually fixed in each *X. gilli* population (Evans et al., 1997, 1998). Our estimate of the divergence times of *X. gilli* corresponds with the Pliocene (Fig. 4 and Table 2) and implicates a Pliocene marine inundation into the area that is now Cape Town and the Cape Flats (Siesser and Dingle, 1981). Patterns of differentiation similar to *X. gilli* have also been reported in other animals (Mounton, 1986; Stewart, 1997).

4.3. Polyploidization

Studies of HOX gene clusters suggest that complete or partial genome duplication happened twice in all vertebrates and once again in most fish (Holland, 1997; Holland and Garcia-Fernández, 1996) but subsequent episodes of polyploidization are infrequent in animals as compared to plants (Orr, 1990). In frogs, tetraploid (probably mostly autopolyploid) species are known in the Leptodactylidae, Myobatrachidae, Hylidae, Bufonidae, Ranidae, Arthroleptidae, and Microhylidae (Duellman and Trueb, 1994; Martino and Sinsch, 2002), but it is in the Pipidae that polyploids comprise the majority of the species, in which species ploidy reaches one of its highest levels (dodecaploidy; Tymowska, 1991), and in which suspected instances of allopolyploidization are most prevalent.

Genome duplication could be advantageous if novel protein function evolves when one duplicated gene copy is released from stabilizing selection while the other carries out the duties of its single-copy precursor (Hughes, 1999; Ohno, 1970). Benefits might be gleaned from increased levels of gene expression after genome duplication (Clark, 1994). Allopolyploid clawed frogs might also benefit from resistance to both sets of parasites that are specific to each of their ancestors (Jackson and Tinsley, 2003). However, polyploidization could be disadvantageous due to complications associated with the formation of multivalents during cell division, and because of difficulties associated with variation in the stoichiometry and expression levels of genes. Allopolyploid speciation, which by definition must be sympatric speciation because it involves hybridization, also poses challenges to a new allopolyploid species, which potentially could be “reabsorbed” into a sympatric species via further hybridization.

Our analyses suggest polyploidization occurred at least six times in clawed frogs (Fig. 4). In *Xenopus*, polyploidization happened at least four times: from 18 to 36 chromosomes at least once, from 36 to 72 chromosomes at least twice, and from 72 + 36 to 108 chromosomes at least once. In *Silurana*, the mtDNA phylogeny supports the occurrence of two polyploidization events to generate tetraploids, though the karyotype of one tetraploid species (*S. new tetraploid 2*) was not available and was estimated by cloning nuclear DNA paralogs of the RAG-1 gene. A strict interpretation of the mtDNA phylogeny suggests that additional episodes of polyploidization occurred, but parametric bootstrap tests did not reject alternative hypotheses of fewer events (Hypotheses 2 and 4).

Ploidy levels of extant pipoids are higher than 20 (Duellman and Trueb, 1994), but an extinct ancestor of *Xenopus* is inferred with a diploid chromosome number of $2N = 18$. Thus, the most parsimonious explanation is that a pair of homologous chromosomes in a diploid ancestor of all *Xenopus* species fused to yield a lower fundamental number of chromosomes ($2N = 18$). Genome duplication then yielded tetraploid, octoploid, and dodecaploid descendants.

5. Conclusions

This work supports the monophyly of clawed frogs and further division of these animals into two clades that are each descended from an ancestor with a different ploidy level: $2N = 18$ for *Xenopus* and $2N = 20$ for *Silurana*. Within *Xenopus*, two major mtDNA clades exist in partial sympatry and these clades are both partially sympatric with *Silurana*. Mitochondrial DNA phylogeography points to an origin of clawed frogs in eastern and/or central equatorial Africa, with *Xenopus*

possibly diversifying from an ancestor in eastern Africa, and *Silurana* possibly diversifying from an ancestor in central Africa. This study also suggests that at least six independent polyploidization events occurred in clawed frogs; four occurred in *Xenopus*, and two in *Silurana*. Further information on nuclear genes will test whether polyploid species that share a maternal ancestor, as suggested by closely related mtDNA, are also derived from the same paternal ancestor.

Acknowledgments

We are grateful to D. Rungger, J. Montoya-Burgos, C. Thiebaud, A. Solaro, M. Picker, and M. Lebreton, and the Station de Zoologie Expérimentale, Université de Genève, for their hospitality. We also thank P. Chippindale, B. Colombelli, L. Du Pasquier, L. Du Preez, M. Fischberg, D. Foguekem, D. Hillis, M. Klemens, H. Kobel, J. Le Doux Diffo, G. Legrand, C. Lieb, C. Loumont, J. Perret, D. Rungger, M. Ruggles, E. Rungger-Brandle, C. Thiebaud, T. Titus, and J. Tymowska for their fieldwork, without which this study would have been impossible. For loans of genetic samples, we thank the following individuals and their respective institutions (museum abbreviations follow Leviton et al., 1985): J. Montoya-Burgos, S. Fischmuller, and A. Schmitz (MHNG), J. Vindum and A. Leviton (CAS), L. Ford (AMNH), T. LaDuc (TNHC), L. Trueb and J. Simmons (KU), and R. Murphy (ROM). We also thank M. Tobias and two anonymous reviewers for helpful comments. This work was supported by INT-01-07294 and a Columbia University Earth Institute Postdoctoral Fellowship (BJE), DEB-9981631 (DCC), NIH grant NS23684 (DBK), and the Center for Environmental Research and Conservation at Columbia University.

References

- Báez, A.M., 1996. The fossil record of the Pipidae. In: Tinsley, R.C., Kobel, H.R. (Eds.), *The Biology of Xenopus*. Clarendon Press, Oxford, pp. 329–347.
- Báez, A.M., Rage, J., 1998. Pipid frogs from the upper Cretaceous of In Beceten, Niger. *Palaeontology* 41, 669–691.
- Báez, A.M., Trueb, L., Calvo, J.O., 2000. The earliest known pipoid frog from South America: a new genus from the middle Cretaceous of Argentina. *J. Vert. Paleontol.* 20, 490–500.
- Biju, S.D., Bossuyt, F., 2003. New frog family from India reveals an ancient biogeographical link with the Seychelles. *Nature* 425, 711–714.
- Bisbee, C.A., Baker, M.A., Wilson, A.C., Hadji-Azimi, I., Fischberg, M., 1977. Albumin phylogeny for clawed frogs (*Xenopus*). *Science* 195, 785–787.
- Bürki, E., Fischberg, M., 1985. Evolution of globin expression in the genus *Xenopus* (Anura: Pipidae). *Mol. Biol. Evol.* 2, 270–277.
- Cannatella, D.C., de Sá, R.O., 1993. *Xenopus laevis* as a model organism. *Syst. Biol.* 42, 476–507.

- Cannatella, D.C., Trueb, L., 1988a. Evolution of pipoid frogs: intergeneric relationships of the aquatic frog family Pipidae (Anura). *Zool. J. Linn. Soc.* 94, 1–38.
- Cannatella, D.C., Trueb, L., 1988b. Evolution of pipoid frogs: morphology and phylogenetic relationships of *Pseudhymenochirus*. *J. Herpetol.* 22, 439–456.
- Carr, S., Brothers, A., Wilson, A., 1987. Evolutionary inferences from restriction maps of mitochondrial DNA from nine taxa of *Xenopus* frogs. *Evolution* 41, 176–188.
- Clark, A.G., 1994. Invasion and maintenance of a gene duplication. *Proc. Natl. Acad. Sci. USA* 91, 2950–2954.
- de Sá, R.O., Hillis, D.M., 1990. Phylogenetic relationships of the pipid frogs *Xenopus* and *Silurana*: an integration of ribosomal DNA and morphology. *Mol. Biol. Evol.* 7, 365–376.
- Duellman, W.E., Trueb, L., 1994. *Biology of Amphibians*. The Johns Hopkins University Press, Baltimore.
- Eriksson, T., 1999. Autodecay, version 4.0, a program used to calculate branch support values. Distributed by author. Stockholm, Bergius Foundation, Royal Swedish Academy of Sciences.
- Estes, R., 1975a. Fossil *Xenopus* from the Paleocene of South America and the zoogeography of pipoid frogs. *Herpetologica*, 31.
- Estes, R., 1975b. *Xenopus* from the Palaeocene of Brazil and its zoogeographic importance. *Nature* 254, 48–50.
- Estes, R., 1977. Relationships of the South African fossil frog *Eoxenopoides reuningi* (Anura, Pipidae). *Ann. South Afr. Mus.* 73, 49–80.
- Estes, R., Spinar, Z.V., Nevo, E., 1978. Early Cretaceous pipid tadpoles from Israel (Amphibia: Anura). *Herpetologica* 34, 374–393.
- Evans, B.J., Brown, R.M., McGuire, J.A., Supriatna, J., Andayani, N., Diesmos, A., Iskandar, D.T., Melnick, D.J., Cannatella, D.C., 2003. Phylogenetics of Fanged Frogs (Anura; Ranidae; *Limnocyrtetes*): testing biogeographical hypotheses at the Asian–Australian faunal zone interface. *Syst. Biol.* 52, 794–819.
- Evans, B.J., Morales, J.C., Picker, M.D., Kelley, D.B., Melnick, D.J., 1997. Comparative molecular phylogeography of two *Xenopus* species, *X. gilli* and *X. laevis*, in the southwestern Cape Province, South Africa. *Mol. Ecol.* 6, 333–343.
- Evans, B.J., Morales, J.C., Picker, M.D., Kelley, D.B., Melnick, D.J., 1998. Absence of extensive introgression between *Xenopus gilli* and *Xenopus laevis laevis* (Anura: Pipidae), in southwestern Cape Province, South Africa. *Copeia* 1998, 504–509.
- Fischer, W.J., Koch, W.A., Elepfandt, A., 2000. Sympatry and hybridization between the clawed frogs *Xenopus laevis laevis* and *Xenopus muelleri* (Pipidae). *J. Zool. Lond.* 252, 99–107.
- Ford, L.S., Cannatella, D.C., 1993. The major clades of frogs. *Herpetol. Monogr.* 7, 94–117.
- Goldman, N., 1993. Statistical tests of models of DNA substitution. *J. Mol. Evol.* 36, 182–198.
- Goldman, N., Anderson, J.P., Rodrigo, A.G., 2000. Likelihood-based tests of topologies in phylogenetics. *Syst. Biol.* 49, 652–670.
- Graf, J.D., 1996. Molecular approaches to the phylogeny of *Xenopus*. In: Tinsley, R.C., Kobel, H.R. (Eds.), *The Biology of Xenopus*. Clarendon Press, Oxford, pp. 379–389.
- Graf, J.D., Fischberg, M., 1986. Albumin evolution of polyploid species of the genus *Xenopus*. *Biochem. Genet.* 24, 821–837.
- Graur, D., Martin, W., 2004. Reading the entrails of chickens: molecular timescales of evolution and the illusion of precision. *Trends Genet.* 20, 80–86.
- Griffiths, C.J., 1993. The geological evolution of East Africa. In: Lovett, J.C., Wasser, S.K. (Eds.), *Biogeography and Ecology of the Rain Forests of Eastern Africa*. Cambridge University Press, Cambridge, pp. 9–22.
- Grohovaz, G.S., Harley, B., Fabran, B., 1996. Significant mitochondrial DNA sequence divergence in natural populations of *Xenopus laevis* (Pipidae) from South Africa. *Herpetologica* 52, 247–253.
- Haffer, J., 1969. Speciation in Amazonia forest birds. *Science* 165, 131–137.
- Henrici, A.C., 1998. A new pipoid anuran from the Late Jurassic Morrison Formation at Dinosaur National Monument, Utah. *J. Vert. Paleontol.* 18, 321–332.
- Henrici, A.C., Báez, A.M., 2001. First occurrence of *Xenopus* (Anura: Pipidae) on the Arabian Peninsula: a new species from the Late Oligocene of the Republic of Yemen. *J. Palaeontol.* 75, 870–882.
- Hillis, D.M., Mable, B.K., Moritz, C., 1996. Applications of molecular systematics: the state of the field and a look to the future. In: Hillis, D.M., Moritz, C., Mable, B.K. (Eds.), *Molecular Systematics*. Sinauer Associates Inc, Sunderland, pp. 515–543.
- Holland, P.W.H., 1997. Vertebrate evolution: something fishy about *Hox* genes. *Curr. Biol.* 7, R570–R572.
- Holland, P.W.H., Garcia-Fernández, J., 1996. *Hox* genes and chordate evolution. *Dev. Biol.* 173, 382–395.
- Huelsenbeck, J.P., Hillis, D.M., Jones, R., 1996. Parametric bootstrapping in molecular phylogenetics: applications and performance. In: Ferraris, J.D., Palumbi, S.R. (Eds.), *Molecular Zoology: Advances, Strategies, and Protocols*. Wiley-Liss, New York, pp. 19–45.
- Huelsenbeck, J.P., Rannala, B., 1997. Phylogenetic methods come of age: testing hypotheses in an evolutionary context. *Science* 276, 227–232.
- Huelsenbeck, J.P., Rannala, B., Masly, J.P., 2000. Accommodating phylogenetic uncertainty in phylogenetic studies. *Science* 288, 2349–2350.
- Huelsenbeck, J.P., Ronquist, F., 2001. MrBayes: Bayesian inference of phylogenetic trees. *Bioinformatics* 17, 754–755.
- Hughes, A.L., 1999. *Adaptive Evolution of Genes and Genomes*. Oxford Press, New York.
- Hughes, M.K., Hughes, A.L., 1993. Evolution of duplicate genes in a tetraploid animal, *Xenopus laevis*. *Mol. Biol. Evol.* 10, 1360–1369.
- Jackson, J.A., Tinsley, R.C., 2003. Parasite infectivity to hybridising host species: a link between hybrid resistance and allopolyploid speciation? *Int. J. Parasitol.* 33, 137–144.
- Knapp, S., Mallet, J., 2003. Refuting refugia? *Science* 300, 71–72.
- Kobel, H., Barandun, B., Thiebaud, C.H., 1998. Mitochondrial rDNA phylogeny in *Xenopus*. *Herpetol. J.* 8, 13–17.
- Kobel, H.R., 1996a. Allopolyploid speciation. In: Tinsley, R.C., Kobel, H.R. (Eds.), *The Biology of Xenopus*. Clarendon Press, Oxford, pp. 391–401.
- Kobel, H.R., 1996b. Reproductive capacity of experimental *Xenopus gilli* × *X. l. laevis* hybrids. In: Tinsley, R.C., Kobel, H.R. (Eds.), *The Biology of Xenopus*. Oxford University Press, Oxford, pp. 73–80.
- Kobel, H.R., Du Pasquier, L., 1986. Genetics of polyploid *Xenopus*. *Trends Genet.* 2, 310–315.
- Kobel, H.R., Loumont, C., Tinsley, R.C., 1996. The extant species. In: Tinsley, R.C., Kobel, H.R. (Eds.), *The Biology of Xenopus*. Clarendon Press, Oxford, pp. 9–33.
- Leviton, A., Gibbs, R., Heal, E., Dawson, C., 1985. Standards in herpetology and ichthyology: part 1. Standard symbolic codes for institutional resource collections in herpetology and ichthyology. *Copeia* 1, 802–832.
- Lovett, J.C., 1993. Climatic history and forest distribution in eastern Africa. In: Lovett, J.C., Wasser, S.K. (Eds.), *Biogeography and Ecology of the Rain Forests of Eastern Africa*. Cambridge University Press, Cambridge, pp. 23–31.
- Maddison, D.R., Maddison, W.P., 2000. *MacClade*. Sinauer Associates, Sunderland.
- Maglia, A.M., Pügener, L.A., Trueb, L., 2001. Comparative development of anurans: using phylogeny to understand ontogeny. *Am. Zool.* 41, 538–551.
- Maisey, J.G., 2000. Continental break up and the distribution of fishes of Western Gondwana during the Early Cretaceous. *Cretaceous Res.* 21, 281–314.
- Mann, M., Risley, M.S., Eckhardt, R.A., Kasinsky, H.E., 1982. Characterization of spermatid/sperm basic chromosomal proteins in the genus *Xenopus* (Anura, Pipidae). *J. Exp. Zool.* 222, 173–186.

- Martino, A.L., Sinsch, U., 2002. Speciation by polyploidy in *Odonotophrynus americanus*. *J. Zool. Lond.* 257, 67–81.
- McCoid, M.J., Fritts, T.H., 1980. Observations of feral populations of *Xenopus laevis* (Pipidae) in southern California. *Bull. South Cal. Acad. Sci.* 79, 82–86.
- McLoughlin, S., 2001. The breakup history of Gondwana and its impact on pre-Cenozoic floristic provincialism. *Austr. J. Bot.* 49, 271–300.
- Measey, G.J., Channing, A., 2003. Phylogeography of the genus *Xenopus* in southern Africa. *Amphibia–Reptilia* 24, 321–330.
- Mounton, P.L.F.N., 1986. Lizard distributions as palaeoenvironmental indicators in the South-Western Cape. *Palaeoecol. Africa Surround. Islands* 17, 231–236.
- Nichol, J.E., 1999. Geomorphological evidence and Pleistocene refugia in Africa. *Geograph. J.* 165, 79–89.
- Ohno, S., 1970. *Evolution by Gene Duplication*. Springer-Verlag, Berlin.
- Orr, H.A., 1990. ‘Why polyploidy is rarer in animals than in plants’ revisited. *Am. Nat.* 136, 759–770.
- Picker, M.D., Harrison, J.A., Wallace, D., 1996. Natural hybridization between *Xenopus laevis laevis* and *X. gilli* in the southwestern Cape Province, South Africa. In: Tinsley, R.C., Kobel, H.R. (Eds.), *The Biology of Xenopus*. Clarendon Press, Oxford, pp. 61–71.
- Pollock, D.D., Zwickl, D.J., McGuire, J.A., Hillis, D.M., 2002. Increased taxon sampling is advantageous for phylogenetic inference. *Syst. Biol.* 51, 664–671.
- Posada, D., Crandall, K.A., 1998. Modeltest: testing the model of DNA substitution. *Bioinformatics* 14, 817–818.
- Púgner, L.A., Maglia, A.M., Trueb, L., 2003. Revisiting the contribution of larval characters to an analysis of phylogenetic relationships of basal anurans. *Zool. J. Linn. Soc.* 139, 129–155.
- Rambaut, A., Grassly, N.C., 1997. Seq-Gen: an application for the Monte Carlo simulation of DNA sequence evolution along phylogenetic trees. *Comp. Appl. Biosci.* 13, 235–238.
- Rice, W.R., 1989. Analyzing tables of statistical tests. *Evolution* 43, 223–225.
- Ronquist, F., 1996. DIVA. Uppsala University, Uppsala.
- Ronquist, F., 1997. Dispersal–vicariance analysis: a new approach to the quantification of historical biogeography. *Syst. Biol.* 46, 195–203.
- Sanderson, M.J., 1997. A nonparametric approach to estimating divergence times in the absence of rate constancy. *Mol. Biol. Evol.* 19, 101–109.
- Sanderson, M.J., 2002. Estimating absolute rates of molecular evolution and divergence times in the absence of rate consistency. *Mol. Biol. Evol.* 19, 1218–1231.
- Sassoon, D., Kelley, D.B., 1986. The sexually dimorphic larynx of *Xenopus laevis*: development and androgen regulation. *Am. J. Anat.* 177, 457–472.
- Schmid, M., Steinlein, C., 1991. Chromosome banding in Amphibia, XVI. High resolution replication banding patterns in *Xenopus laevis*. *Chromosoma* 101, 123–132.
- Shubin, N.H., Jenkins, F.A., 1995. An early Jurassic jumping frog. *Nature* 377, 49–52.
- Siesser, W.G., Dingle, R.V., 1981. Tertiary sea-level movements around southern Africa. *J. Geol.* 89, 83–96.
- Stewart, B.A., 1997. Morphological and genetic differentiation between populations of river crabs (Decapoda: Potamonautidae) from the Western Cape, South Africa, with a taxonomic re-examination of *Gecarcinautes brincki*. *Zool. J. Linn. Soc.* 119, 1–21.
- Stokes, S., Thomas, D.S.G., Washington, R., 1997. Multiple episodes of aridity in southern Africa since the last interglacial period. *Nature* 388, 154–158.
- Swofford, D.L., 2002. *Phylogenetic Analysis Using Parsimony* (* and other methods). Version 4. Sinauer Associates, Sunderland.
- Thiebaud, C.H., Fischberg, M., 1977. DNA content in the genus *Xenopus*. *Chromosoma* 59, 253–257.
- Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F., Higgins, D.G., 1997. The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 24, 4876–4882.
- Tinsley, R.C., Kobel, H. (Eds.), 1996. *The Biology of Xenopus*. Clarendon Press, Oxford.
- Tinsley, R.C., Loumont, C., Kobel, H.R., 1996. Geographical distribution and ecology. In: Tinsley, R.C., Kobel, H.R. (Eds.), *The Biology of Xenopus*. Clarendon Press, Oxford, pp. 35–59.
- Tinsley, R.C., McCoid, M.J., 1996. Feral populations of *Xenopus* outside Africa. In: Tinsley, R.C., Kobel, H.R. (Eds.), *The Biology of Xenopus*. Clarendon Press, Oxford, pp. 81–94.
- Tobias, M.L., Viswanathan, S.S., Kelley, D.B., 1998. Rapping, a female receptive call, initiates male–female duets in the South African clawed frog. *Proc. Natl. Acad. Sci. USA* 95, 1870–1875.
- Tobias, M.L., Barnard, C., O’Hagan, R., Horng, S.H., Rand, M., Kelley, D.B., 2004. Vocal communication between male *Xenopus laevis*. *Animal Behaviour* 67 (2), 353–365.
- Trueb, L., 1996. Historical constraints and morphological novelties in the evolution of the skeletal system of pipid frogs (Anura: Pipidae). In: Tinsley, R.C., Kobel, H.R. (Eds.), *The Biology of Xenopus*. Clarendon Press, Oxford, pp. 349–377.
- Trueb, L., 1999. The early cretaceous pipoid anuran, *Thoraciliacus*: redescription, reevaluation, and taxonomic status. *Herpetologica* 55, 139–157.
- Trueb, L., Báez, A.M., 1997. Redescription of the Paleogene *Shelania pascuali* from Patagonia and its bearing on the relationships of fossil and Recent pipoid frogs. *Scientific Papers, Natural History Museum, University of Kansas*, 4, pp. 1–41.
- Trueb, L., Cannatella, D.C., 1986. Systematics, morphology, and phylogeny of genus *Pipa* (Anura: Pipidae). *Herpetologica* 42, 412–449.
- Tymowska, J., 1991. Polyploidy and cytogenetic variation in frogs of the genus *Xenopus*. In: Green, D.S., Sessions, S.K. (Eds.), *Amphibian Cytogenetics and Evolution*. Academic Press, San Diego.
- Yager, D.D., 1996. Sound production and acoustic communication in *Xenopus borealis*. In: Tinsley, R.C., Kobel, H.R. (Eds.), *The Biology of Xenopus*. Clarendon Press, Oxford, pp. 121–141.
- Yoder, A.D., Yang, Z., 2004. Divergence dates for Malagasy lemurs estimated from multiple gene loci: geological and evolutionary context. *Mol. Ecol.* 13, 757–773.
- Zwickl, D.J., Hillis, D.M., 2002. Increased taxon sampling greatly reduces phylogenetic error. *Syst. Biol.* 51, 588–598.