

F4 A finely grained gray limestone. The fossils are comparatively much less abundant than in either the preceding or succeeding bed. Coeymans 3 feet

The following is a list of the fossils:

| | |
|--------------------------------|--|
| 7 Favosites sphaericus Hall r | 89 Stropheodonta varistriata? (Conrad) R |
| 12 Lichenalia torta Hall c | 102 Uncinulus sp. |
| 45 Gypidula galeata (Dalman) c | 123 Platyceras sp. R |

F5 A coarsely grained, dark gray limestone made up in great part of large crinoid joints. Most of the grains are surrounded by a coating of limonite. Coeymans..... 4½ feet

The following fossils were obtained here:

| | |
|--------------------------------|--------------------------------|
| 6 Favosites heldbergiae Hall C | 45 Gypidula galeata (Dalman) C |
| 7 F. sphaericus Hall c | G. galeata var. |
| 12 Lichenalia torta Hall r | 88 Stropheodonta becki Hall |

F6 Alternately finely and coarsely grained limestone. Coeymans 20 feet

This is subdivided as follows:

F6a Finely grained similar to F4..... 15 inches
 F6b Coarsely grained similar to F5..... 8 inches
 F6c Finely grained, friable..... 5 inches
 F6d Very coarsely grained. Favosites heldbergiae Hall is common here..... 4½ feet
 F6e Finely grained limestone, at times shaly..... 10 inches
 F6f Very coarsely grained..... 3½ feet

Here were found:

| | |
|----------------------------------|--------------------------------|
| 12 Lichenalia torta Hall r | 45 Gypidula galeata (Dalman) c |
| 40 Dalmanella subcarinata Hall r | |

F6g A rather coarsely grained limestone. The upper foot and a half is at times exceedingly coarse grained, with limonite surrounding each grain where weathered..... 9 feet

The following fossils were found here:

| | |
|-------------------------------|---------------------------------|
| 5 Enterolasma strictum Hall c | 23 Atrypa reticularis (Linné) r |
| 8 Favosites sp. | 44 Gypidula galeata (Dalman) C |