58 Nucleospira ventricosa Hall r 75 Camarotoechia semiplicata (Conrad) r

5 Enterolasma strictum Hall c 47 Leptaena rhomboidalis (Wilck-45 Gypidula galeata (Dalman) c ens) c

F7c Very dense, dark blue limestone with very many chert bands which, when weathered, furnish excellent bryozoa. The chert of the preceding beds even when weathered shows little else than crinoid joints and comparatively few of these. Coeymans.. 4 feet 4 inches

The following fossils were collected here, most of them from the weathered surfaces of the chert bands:

```
5 Enterolasma strictum Hall c
```

- 12 Lichenalia torta Hall r
- 13 Lioclema cellulosum (Hall) c
- 17 Orthopora regularis (Hall) c
- 18 O. rhombifera (Hall) C
- 19 Unitrypa nervia (Hall) r
- 20 U. praecursa (Hall) c
- 33 Coelospira concava Hall r
- 37 Cyrtina sp. R
- 39 Dalmanella perelegans Hall r
- 40 D. subcarinata Hall c
- 41 Delthyris perlamellosa Hall c

- 45 Gypidula galeata (Dalman) c
- 47 Leptaena rhomboidalis (Wilck-ens)
 - Orbiculoidea sp.
- 63 Pholidops ovata Hall c
- 68 Reticularia modesta (Hall) r
- 71 Rhipidomella oblata Hall R
- 73 Camarotoechia altiplicata? Hall R
- 77 Rhynchospira globosa? Hall r
- 88 Stropheodonta becki Hall r
- 136 Dalmanites nasutus? Conrad R

The following fossils were collected:

- 5 Enterolasma strictum Hall C
- 7 Favosites sphaericus Hall r
- 26 Bilobites varicus (Conrad) r
- 40 Dalmanella subcarinata Hall c