

- |  |   |
|--|---|
| 5 Enterolasma strictum <i>Hall</i> r<br>21 Vermipora serpuloides <i>Hall</i> r<br>33 Coelospira concava <i>Hall</i> c<br>40 Dalmanella subcarinata <i>Hall</i> C<br>47 Leptaena rhomboidalis ( <i>Wilckens</i> ) c<br>55 Meristella <i>sp.</i> r | 71 Rhipidomella oblates <i>Hall</i> r<br>82 Spirifer cyclopterus <i>Hall</i> C<br>88 Stropheodonta becki <i>Hall</i> c<br>94 Strophonella punctulifera ( <i>Conrad</i> ) r<br>137 Dalmanites <i>sp.</i><br>140 Phacops <i>cf.</i> <i>logani</i> <i>Hall</i> r |
|--|---|

L<sub>3</sub> Upper New Scotland..... 8½ feet  
 Subdivided as follows:

L<sub>3a</sub> Dark blue limestone in beds from 3 to 6 inches thick with considerable chert in the lower beds..... 3½ feet

L<sub>3b</sub> Very cherty, bluish gray limestone..... 1 foot

L<sub>3c</sub> Coarsely grained to very finely grained limestone usually in a single bed. Chert is developed in its lower portion. At times there are mere pockets of the coarsely grained scattered at intervals through the finely grained. This is very fossiliferous..... 2½ feet

L<sub>3d</sub> A dark bluish gray limestone alternating with half inch bands of an arenaceous limestone, the latter showing light and dark laminae and few or no fossils..... 20 inches

The following is a list of fossils identified from L<sub>3</sub>. The fossils from the different beds were not kept separate.

- |  |   |
|--|---|
| 5 Enterolasma strictum <i>Hall</i> r<br>12 Lichenalia torta <i>Hall</i> c<br>31 Chonostrophia jervisensis <i>Schuchert</i> R<br>40 Dalmanella subcarinata <i>Hall</i> r<br>41 Delthyris perlamellosa <i>Hall</i> c<br>47 Leptaena rhomboidalis ( <i>Wilckens</i> ) c | 52 Meristella laevis ( <i>Vanuxem</i> ) r<br>55 M. <i>sp.</i> c<br>82 Spirifer cyclopterus <i>Hall</i> c<br>88 Stropheodonta becki <i>Hall</i> c<br>95 Trematospira multistriata <i>Hall</i> R<br>97 Uncinulus campbellanus ( <i>Hall</i> ) R<br>101 U. vellicatus <i>Hall</i> R<br>127 Tentaculites elongatus? <i>Hall</i> R |
|--|---|

L<sub>4</sub> Coarsely grained, gray limestone, usually in one bed. Very fossiliferous. Be craft..... 2½ feet

- |  |  |
|--|--|
| 5 Enterolasma strictum <i>Hall</i> r<br>11 Edriocrinus pocilliformis <i>Hall</i> c<br>12 Lichenalia torta <i>Hall</i> r<br>23 Atrypa reticularis ( <i>Linn.</i> ) r<br>46 Gypidula pseudogaleata ( <i>Hall</i> ) C<br>47 Leptaena rhomboidalis ( <i>Wilckens</i> ) C | 52 Meristella laevis ( <i>Vanuxem</i> ) C<br>54 M. princeps <i>Hall</i> r<br>81 Spirifer concinnus <i>Hall</i> c<br>82 S. cyclopterus <i>Hall</i> r<br>92 Strophonella headleyana? <i>Hall</i> R |
|--|--|

L<sub>5</sub> Be craft..... 5 feet, 9 inches

L<sub>5a</sub> Rather finely grained, dark gray limestone. The basal foot