

	Feet
3 Dark bluish gray, somewhat impure limestone, in beds of variable thickness; often packed with <i>Orthis costalis</i> Hall, which occurs with more or less frequency through the whole mass. Other fossils are: <i>Lingula huronensis</i> Bill., <i>Harpes antiquatus</i> Bill., <i>H. ottawaensis</i> Bill.(?), <i>Illaenus arcturus</i> Hall, (<i>I bayfieldi</i> Bill.), <i>Lituities</i> sp.(?)... 110	110
4 Gray, tolerably pure limestone in beds 8 to 20 inches thick, separated by earthy seams, the bedding being uneven. Many layers consist of crinoidal fragments, largely of <i>Paleocystites tenuiradiatus</i> Hall. Near the middle of the mass for a thickness of 10 feet, some of the fragments and small, ovoid masses (<i>Bolboporites americanus</i> Bill.) are of a bright red color: 90	90
Making for the total thickness of A.....	338

Group B (Middle Chazy)

1 Impure, nodular limestone containing <i>Maclurea magna</i> Les.....	25
2 Gray, massive, pure limestone, abounding in crinoidal fragments	20
3 Bluish black, thick bedded limestone, usually weathering so as to show pure nodular masses enveloped in a somewhat impure lighter colored matrix; everywhere characterized by <i>Maclurea magna</i> . Near the middle of this mass for a thickness of about 30 feet, the fossils are silicified and of jet-black color. The more important besides <i>Maclurea</i> , are species of <i>Strophomena</i> , <i>Orthis</i> and <i>Orthoceras</i>	210
4 Dark, compact, fine grained limestone, with obscure bedding, weathering to a light gray. Fossils are infrequent, but at a single locality were collected <i>Orthis perveta</i> Con., <i>O. platys</i> Bill., <i>Leptaena fasciata</i> Hall, <i>Asaphus canalis</i> Con., <i>Cheirurus polydorus</i> Bill., <i>Harpes</i> sp. und., <i>Illaenus incertus</i> Bill., <i>Lichas minganensis</i> Bill., <i>Sphaerexochus parvus</i> Bill., and several undescribed species.....	20