

CHEMISTRY S1403
 PROFESSOR J. MORROW

FIRST EXAM

6/4/99

PRINT NAME, LAST: _____

FIRST: _____

ID.#: _____

MAXIMUM POINT VALUE IS IN PARENTHESES

- | | | |
|---------------|----------------|----------------|
| 1. _____ (6) | 8. _____ (6) | 15. _____ (4) |
| 2. _____ (15) | 9. _____ (12) | 16. _____ (4) |
| 3. _____ (10) | 10. _____ (5) | 17. _____ (4) |
| 4. _____ (12) | 11. _____ (5) | 18. _____ (8) |
| 5. _____ (8) | 12. _____ (10) | 19. _____ (10) |
| 6. _____ (8) | 13. _____ (8) | |
| 7. _____ (8) | 14. _____ (4) | |

COLUMN TOTALS (MAXIMUM):

_____ (67) _____ (50) _____ (30)

EXAM TOTAL (147 pts) _____



 OUT OF 100

NO PARTIAL CREDIT on any question except where indicated by the statement SHOW WORK.
 CHECK FRONT BLACKBOARD FOR CORRECTIONS/CHANGES.
 SUGGESTION: DO THE SIMPLER PROBLEMS FIRST.
 IF ANY PART OF EXAM IS NOT CLEAR - ASK PROCTORS ABOUT IT!


SCRAP WORK SHEETS ARE AT THE END OF EXAM. FEEL FREE TO TEAR THESE PAGES OFF.

USE THE FOLLOWING INFORMATION FOR PROBLEMS 1 AND 2

Given the following 3 reactions:



and



You are given a mixture of iron(II)carbonate (FeCO_3), iron(II)bicarbonate ($\text{Fe}(\text{HCO}_3)_2$) and sodium hydroxide (NaOH). When heated this mixture completely reacts as shown above, forming 17.60 g of $\text{CO}_{2(g)}$, 3.60 g of H_2O , and 0.100 mol Na_2O .

Molar masses: FeCO_3 (115.9), $\text{Fe}(\text{HCO}_3)_2$ (177.9), NaOH (40.0),
 CO_2 (44.0), FeO (71.9), H_2O (18.0), Na_2O (62.0)

1) Calculate the number of moles of H_2O and of CO_2 formed. (6 pts)

ANSWER IS (H_2O): _____

ANSWER IS (CO_2): _____


2) Calculate the number of moles of FeCO_3 , $\text{Fe}(\text{HCO}_3)_2$, and NaOH present initially. SHOW WORK (15 pts - 5 pts each part)

$\underline{n}_{\text{NaOH}}$ IS: _____


$\underline{n}_{\text{Fe}(\text{HCO}_3)_2}$ IS: _____

$\underline{n}_{\text{FeCO}_3}$ IS: _____

3) Given the following 2 reactions:



and




Starting with 1 mole total of FeCO_3 and $\text{Fe}_2(\text{CO}_3)_3$, 1.5 mol of CO_2 are obtained. Calculate the starting number of moles of FeCO_3 .

SHOW WORK (10 pts)

ANSWER IS: _____

- 4) The following gaseous reaction occurs in a vessel of 50.0 L volume at 300 K.



Initially, three moles of N_2H_4 and two moles of F_2 are mixed in this vessel. The reaction then occurs until the reactant in limiting quantity is totally consumed. (12 pts - 4 pts each part)

Molar masses: N_2H_4 (32.0), F_2 (38.0), HF (20.0), NF_3 (71.0)

a) Which reactant is limiting? ANSWER IS: _____


b) How many moles of the reactant in excess remain, when the reaction is complete?

ANSWER IS: _____

c) How many grams of NF_3 are produced?

ANSWER IS: _____


- 5) What volume of CO_2 can be produced from the reaction of 13.1 g of N_2O (as shown below) ? Assume an excess of C_3H_8 , and take the density of CO_2 to be $1.96 \frac{\text{g}}{\text{L}}$. Molar masses: $\text{N}_2\text{O} = \text{CO}_2 = 44.0$ (8 pts)



i) 2.00 L ii) 2.24 L iii) 3.50 L iv) 6.00 L

ANSWER IS: _____

- 6) Given the reaction:



How many liters of CO_2 were recovered by burning 20 L of C_4H_{10} ?
(P and T constant) if the percent yield was 75 % . (8 pts)

- i) 55 L ii) 60 L iii) 70 L iv) 80 L

ANSWER IS: _____

7) An unknown gas has a density of $6.36 \frac{\text{g}}{\text{L}}$ at a pressure of 0.912 atm
and a temperature of 57°C . What is the molar mass of this gas?

$$R = 0.0821 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{deg}} \quad (8 \text{ pts})$$

- i) 28.0 g/mol ii) 114 g/mol iii) 146 g/mol iv) 189 g/mol

ANSWER IS: _____





8) What would be the density of the gas from question 7 if its
temperature and pressure returned to STP? (6 pts)
CREDIT FOR THIS IS BASED UPON YOUR ANSWER FROM QUESTION 7.

ANSWER IS: _____

9) GIVEN: ZrCl_2 is zirconium(II)chloride; ZrCl_4 is; NaVO_3 is sodium
vanadate. For the following: where there is a formula, give it's name;
where there is a name, give it's formula. (12 pts - 2 points each.)

- i) $\text{Zr}(\text{SO}_4)_2$:
ii) SF_6
iii) zirconium(II)vanadate :
iv) CrBr_3 :
v) ferric phosphate :
vi) aluminum chromate

BALANCE THE FOLLOWING EQUATIONS BY INSERTING INTEGERS IN THE
SPACES PRECEDING THE FORMULAS. (5 pts each)



- 12) A compound Q is composed of the elements D, L, and M. For every 2 atoms of D, there are 3 atoms of L and 2 atoms of M. Starting with 0.240×10^{23} atoms of D, and 0.0700 moles of L, exactly 1.20 g of M react. First write the formula of the compound. HINT: This is a limiting quantity problem. (10 pts)

The molar mass (atomic weight) of element M is,

i) $30 \frac{\text{g}}{\text{mol}}$


ii) $45 \frac{\text{g}}{\text{mol}}$

iii) $60 \frac{\text{g}}{\text{mol}}$


iv) $90 \frac{\text{g}}{\text{mol}}$

ANSWER IS: _____

- 13) Give the formal charge of each indicated atom in the thiosulfate ($\text{S}_2\text{O}_3^{2-}$) anion. (8 pts)



- 14) Indicate which reactant species is the Lewis acid and which is the Lewis base in the following reaction. HINT: Think Lewis structure. (4 pts)



15) Give the conjugate acid of $(\text{CH}_3)_2\text{NH}$. (4 pts)

ANSWER IS: _____

16) Give the hydrated form of the anhydrous acid I_2O_5 . (4 pts)

ANSWER IS: _____

17) Give the formula for the anhydrous form of H_2SeO_4 . (4 pts)


ANSWER IS: _____

18) Beaker A contains 0.100 L of an 0.20 M KOH solution; beaker B contains 0.100 L of an 0.40 M HCl solution. The contents of both beakers are thoroughly mixed together in a sufficiently large third beaker. The molarity, M, of the resulting salt solution is; (8 pts)

i) 0.05 M. ii) 0.10 M. iii) 0.20 M. iv) 0.40 M. v) no salt is formed.

ANSWER IS: _____

19) Complete combustion (burning) of two (2) liters of a gaseous hydrocarbon, C_xH_y , to $\text{CO}_2(\text{g})$ and $\text{H}_2\text{O}(\text{g})$ required 14 liters of pure $\text{O}_2(\text{g})$. All volumes were measured at the same temperature and pressure. The TOTAL volume of the products, CO_2 and H_2O , is 18 liters. The unbalanced reaction is,



The most likely molecular formula of the hydrocarbon is; (10 pts)

i) C_6H_{12} ii) C_5H_{12} iii) C_4H_{10} iv) C_5H_8

ANSWER IS: _____

SCRAP WORK PAGE

SCRAP WORK PAGE

SCRAP WORK PAGE