Human Development:

Fertilization through gastrulation

Michael M. Shen, Ph.D.

Departments of Medicine and Genetics & Development

Columbia University Medical Center

Gastrulation movements in the frog embryo



Vegetal view

From blastula to gastrula





The first week of development



- Fertilization
- Cleavage stages
- Blastocyst formation
- Early lineage specification
- Implantation

The second week of development



- Trophoblast differentiation
- Yolk sac formation
- Anterior-posterior axis patterning
- Initiation of gastrulation

The third week of development



- Endoderm and mesoderm ingression
- Mesoderm lineage specification
- Left-right patterning
- Neural plate formation
- Axial midline formation

Pre-implantation mouse development



- Reductive cleavage
- Blastomere potency
- Inside-outside allocation of lineage progenitors
- Compaction
- Blastocyst formation
- Emerging morphological asymmetry

Human embryo development in culture



Fertilization

Cleavages

Compaction

Blastocyst formation

Early cleavages of the mouse embryo



(Bischoff et al. (2008))

Key properties of vertebrate embryogenesis

- Regulative development

  Early blastomeres are totipotent

Regulative development of the vertebrate embryo



(DeRobertis (2006))

Gene expression at pre-implantation stages in the mouse



Zygotic genome activity

Mid-preimplantation genome activity

(Wang and Dey (2006))

Cell types of the blastocyst



Specification of early lineages



(Wang and Dey (2006))

Model for primitive endoderm (hypoblast) specification



(Chazaud *et al.* (2006))

Pluripotency of mouse ES cells

Can contribute to all embryonic cell types in chimeras – including the germ line

Early lineages and stem cells in the mouse embryo



Mouse EpiSC cells resemble human ES cells

Process of implantation



Formation of extraembryonic tissues



Key properties of vertebrate embryogenesis

- Regulative development
- Patterning at a distance by soluble morphogens

Two major signaling pathways regulate early patterning and differentation



Schematic pathway for canonical Wnt/ beta-catenin signaling



Wnt ligand absent *Wnt ligand present*

Schematic pathway for TGF-beta signaling



The Nodal signaling pathway



Key properties of vertebrate embryogenesis

- Regulative development
- Patterning at a distance by soluble morphogens
- Common patterning mechanisms underlie distinct embryo morphologies

Schematic of early mouse development



(Adapted from Hogan *et al.* (1994))

Morphological relationship between mouse and human embryos

Cup-shaped vs discoid

mouse human



(Eakin and Behringer)

Key properties of vertebrate embryogenesis

- Regulative development
- Patterning at a distance by soluble morphogens
- Common patterning mechanisms underlie distinct embryo morphologies
- Antagonism of secreted ligands and inhibitors

Specification of the anterior-posterior axis in the mouse



Nodal and Cripto activity

Nodal inhibitor activity (Lefty, Cerberus)

Movement of the anterior visceral endoderm





View from anterior side

Relationship of blastodisc to implantation site



Formation of the primitive streak



Expression of Brachyury in chick embryo

Early embryogenesis in the chick



Anterior

Posterior

Ingression of nascent endoderm and mesoderm through the streak



- Delamination of epiblast cells
- Movement through the streak
- Initial ingression of endoderm
- Subsequent ingression of mesoderm

Anterior and lateral migration of mesoderm



- *Anterior migration of mesoderm:*
  - *Axial (prechordal)*
  - *Cardiac*
- *Lateral distance from midline determines mesoderm type:*
  - Axial (e.g., notochord)
  - Paraxial (somites)
  - Intermediate (e.g., kidney)
  - Lateral (e.g, limbs)

Regional differentiation of mesoderm





Chick embryo

Anterior-posterior patterning of axial mesoderm



Key properties of vertebrate embryogenesis

- Regulative development
- Patterning at a distance by soluble morphogens
- Common patterning mechanisms underlie distinct embryo morphologies
- Antagonism of secreted ligands and inhibitors
- Instructive inductive interactions

Spemann-Mangold organizer experiment



Blastopore lip transplantation



Induction of secondary axis

(DeRobertis and Kuroda (2004))

Injection of Wnts or Nodal can induce a secondary axis



- *Injection of mRNA into dorsal marginal zone*
  - Wnt8 (complete axis)
  - Nodal (partial axis)

Formation of the neural plate



Macaque embryo
(similar to 20 day human embryo)

Inductive interactions and head formation



AVE	Anterior visceral endoderm
EPI	Epiblast
NE	Neural progenitor
EGO	Early gastrula organizer
PS	Primitive streak

Dorsoventral patterning by axial and paraxial mesoderm





Holoprosencephaly in *Cripto* hypomorphs



Defective forebrain patterning and axial mesoderm formation



Spectrum of human holoprosencephaly

Complex L-R laterality of tissues



(Kosaki and Casey (1998))

Nomenclature for L-R laterality phenotypes



- *Situs solitus:* normal organ position
- *Situs inversus:* complete reversal of organ position
- *Isomerism:* mirror image duplication of tissue morphology
- *Heterotaxia:* discordant and randomized organ position

(Capdevila et al. (2000))

Stages of L-R laterality determination

- Initial symmetry breaking



Nodal flow model

Stages of L-R laterality determination

- Initial symmetry breaking
- Propagation and maintenance of an asymmetric signal
- Specification of tissue-specific laterality



Asymmetric gene expression

Asymmetric expression of *Nodal* and *Lefty*

Nodal

Lefty



(Beddington and Robertson (1998))

Left-right laterality defects in *Cryptic* mutants

Wild-type | *Cryptic*$^{-/-}$ | *Cryptic*$^{-/-}$



Wild-type | *Cryptic*$^{-/-}$



Wild-type | *Cryptic*$^{-/-}$

Cardiac defects in *Cryptic* mutants



Wild-type *Cryptic*$^{-/-}$ *Cryptic*$^{-/-}$

Transposition of the great arteries

Normal



Transposed

Wild-type



Cryptic$^{-/-}$

Morphological changes at early post-gastrulation stages