# Pathophysiology: Heart Failure

Mat Maurer, MD Associate Professor of Clinical Medicine

# **Objectives**

At the conclusion of this seminar, learners will be able to:


- 1. Define heart failure as a clinical syndrome
- 2. Define and employ the terms preload, afterload, contractilty, remodeling, diastolic dysfunction, compliance, stiffness and capacitance.
- 3. Describe the classic pathophysiologic steps in the development of heart failure.
- 4. Delineate four basic mechanisms underlying the development of heart failure
- 5. Interpret pressure volume loops / Starling curves and identify contributing mechanisms for heart failure state.
- 6. Understand the common methods employed for classifying patients with heart failure.
- 7. Employ the classes and stages of heart failure in describing a clinical scenario


## **Heart Failure**


- Not a disease
- A syndrome
  - From "syn" meaning "together" and "dromos" meaning "a running".
  - A group of signs and symptoms that occur together and characterize a particular abnormality.
- Diverse etiologies
- Several mechanisms


### **Heart Failure: Definitions**


- An inability of the heart to pump blood at a sufficient rate to meet the metabolic demands of the body (e.g. oxygen and cell nutrients) at rest and during effort or to do so only if the cardiac filling pressures are abnormally high.
- A complex clinical syndrome characterized by abnormalities in cardiac function and neurohormonal regulation, which are accompanied by effort intolerance, fluid retention and a reduced longevity
- A complex clinical syndrome that can result from any structural or functional cardiac disorder that impairs the ability of the ventricle to fill with or eject blood.

























#### **<u>Ea (Arterial Elastance)</u>**


- If
  - TPR = [MAP CVP] / CO, and
  - CO = SV \* HR
- Substituting the second equation into the first we obtain:
- TPR = [MAP CVP] / (SV\*HR)
- Making two simplifying assumptions.
  - 1. CVP is negligible compared to MAP.
  - 2. MAP is approximately equal to the end-systolic pressure in the ventricle (Pes).
- Then,
- TPR = Pes / (SV\*HR)
- Which can be rearranged to:



































Acute and Chronic Responses – Benefits and Harm		
Response	Short-term Effects (mainly adaptive; hemorrhage, acute heart failure)	Long-term Effects (mainly deleterious; chronic heart failure)
Salt and water retention	Augments preload	Pulmonary congestion, anasarca
Vasoconstriction	Maintains pressure for perfusion of vital organs (brain, heart)	Exacerbates pump dysfunction, increases cardiac energy expenditure
Sympathetic stimulation	Increases heart rate and ejection	Increases energy expenditure
Cytokine activation	Vasodilatation	Skeletal muscle catabolism, deterioration of endothelial function, impaired contraction, LV remodeling.
Hypertrophy	Unloads individual muscle fibers	Deterioration and death of cardiac cells: cardiomyopathy of overload
Increased collagen	May reduce dilatation	Impairs relaxation










# **Pathophysiology of Heart Failure**


#### Four Basic Mechanisms

- 1. Increased Blood Volume (Excessive Preload)
- 2. Increased Resistant to Blood Flow (Excessive Afterload)
- 3. Decreased contractility
- 4. Decreased Filling

















<b>Types of Heart Failure</b>			
	SHF	Diastolic	
Pathophysiology	Impaired Contraction	Impaired filling	
Demographics	All ages	> 60 years	
1° CauseCoronary Artery DiseaseHypertension			




# Systolic Versus Diastolic Failure






# **Decompensated Heart Failure**






# **High vs. Low Output Failure**




- Causes:
  - Anemia
  - Systemic arteriovenous fistulas
  - Hyperthyroidism
  - Beriberi heart disease
  - Paget disease of bone
  - Glomerulonephritis
  - Polycythemia vera
  - Carcinoid syndrome
  - Obesity
  - Anemia
  - Multiple myeloma
    - Pregnancy
  - Cor pulmonale
  - Polycythemia vera



# **Dilated vs. Hypertrophic vs. Restrictive**

Туре	Definition	Sample Etiologies
Dilated	Dilated left/both ventricle(s) with impaired contraction	Ischemic, idiopathic, familial, viral, alcoholic, toxic, valvular
Hypertrophic	Left and/or right ventricular hypertrophy	Familial with autosomal dominant inheritance
Restrictive	Restrictive filling and reduced diastolic filling of one/both ventricles, Normal/near normal systolic function	Idiopathic, amyloidosis, endomyocardial fibrosis



# **Clinical Manifestations**

#### **Symptoms**


- Reduced exercise tolerance
- Shortness of breath
- Congestion
- Fluid retention
- Difficulty in sleeping
- Weight loss

Variable	Sensitivity	Specificity	
Hx of HF	62	94	
Dyspnea	56	53	
Orthopnea	47	88	
Rales	56	80	
S3	20	99	
JVD	39	94	
Edema	67	68	


# **Diagnosis of heart failure**

- Physical examination
- Chest X ray
- EKG
- Echocardiogram
- Blood tests: Na, BUN, Creatinine, BNP
- Exercise test
- MRI
- Cardiac catheterization






NYHA Classification			
		Class	Patient Symptoms
	I	Mild	<ul><li>No limitation of physical activity</li><li>No undue fatigue, palpitation or dyspnea</li></ul>
	Ш	Mild	<ul> <li>Slight limitation of physical activity</li> <li>Comfortable at rest</li> <li>Less than ordinary activity results in fatigue, palpitation, or dyspnea</li> </ul>
	ш	Moderate	<ul> <li>Marked limitation of physical activity</li> <li>Comfortable at rest</li> <li>Less than ordinary activity results in fatigue, palpitation, or dyspnea</li> </ul>
	IV	Severe	<ul> <li>Unable to carry out any physical activity without discomfort</li> <li>Symptoms of cardiac insufficiency at rest</li> <li>Physical activity causes increased discomfort</li> </ul>



<b>ACC/AHA Staging System</b>		
	Stage	Patient Description
A	High risk for developing heart failure	<ul> <li>Hypertension</li> <li>Coronary artery disease</li> <li>Diabetes mellitus</li> <li>Family history of cardiomyopathy</li> </ul>
В	Asymptomatic heart failure	<ul> <li>Previous myocardial infarction</li> <li>Left ventricular systolic dysfunction</li> <li>Asymptomatic valvular disease</li> </ul>
С	Symptomatic heart failure	<ul><li>Known structural heart disease</li><li>Shortness of breath and fatigue</li><li>Reduced exercise tolerance</li></ul>
D	Refractory end-stage heart failure	• Marked symptoms at rest despite maximal medical therapy (e.g., those who are recurrently hospitalized or cannot be safely discharged from the hospital without specialized interventions)




## **Etiologies**

- Ischemic cardiomyopathy
- Valvular cardiomyopathy
- Hypertensive cardiomyopathy.
- Inflammatory cardiomyopathy
- Metabolic cardiomyopathy
- General system disease
- Muscular dystrophies.
- Neuromuscular disorders.
- Sensitivity and toxic reactions.
- Peripartal cardiomyopathy

### **Percipients / Associated Factors**

- Inappropriate reduction in the intensity of treatment, including
  - Dietary sodium restriction,
  - Physical activity reduction,
  - Drug regimen reduction, or,
  - most commonly, a combination of these measures.
- Ischemia
- Hypertension
- Anemia
- Volume Overload
- Increased Metabolic Demand
  - Infection
  - Thyroid Disease
- Arrhythmia
- Asthma/COPD

# **Targets of Treatment**





# Standard Pharmacological Therapy

- ACE inhibitors
- Angiotensin Receptor Blockers
   Beta Blcokers

  - Diuretics
  - Aldosterone Antagonists
    - Statins
    - Vasodilators
      - Inotropes

Treatment		
	Stage	Patient Treatment
A	High risk for developing heart failure	<ul> <li>Optimal pharmacologic therapy (OPT)</li> <li>Aspirin, ACE inhibitors, statins, b-blockers, a-b-blockers (carvedilol) diabetic therapy</li> </ul>
В	Asymptomatic heart failure	<ul> <li>OPT</li> <li>ICD if left ventricular (LV) dysfunction (systolic) present</li> </ul>
С	Symptomatic heart failure	<ul> <li>OPT</li> <li>ICD if LV dysfunction (systolic) present</li> <li>CRT (if QRS wide, LVEF≤35%)</li> </ul>
D	Refractory end-stage heart failure	<ul> <li>OPT</li> <li>Intermittent IV inotropes</li> <li>ICD as a bridge to transplantation</li> <li>CRT</li> <li>Other devices (LVAD, pericardial restraint)</li> </ul>





# Summary

- Complex Clinical Syndrome
- Multiple Etiologies and Classification Systems
- Physiologic Understanding Essential

http://www.columbia.edu/itc/hs/medical/heartsim/

