

Viral Replication

Scott M. Hammer, M.D.

Steps in Viral Replication: Attachment (First Step)

- Surface protein on virus attaches to specific receptor(s) on cell surface
 - May be specialized proteins with limited tissue distribution or more widely distributed
 - Virus specific receptor is necessary but not sufficient for viruses to infect cells and complete replicative cycle

Viral Replication: Basic Concepts

- Viruses are obligate intracellular parasites
- Viruses carry their genome (RNA or DNA) and sometimes functional proteins required for early steps in replication cycle
- Viruses depend on host cell machinery to complete replication cycle and must commandeer that machinery to successfully replicate

Selected Virus Receptors

Adenovirus	CAR
Coxsackievirus	CAR, CD55
Echovirus	Integrin VLA-2, CD55
Epstein-Barr Virus	CD21
HIV-1	CD4, CCR5, CXCR4
Measles virus	CD46
Parvovirus	Erythrocyte P Ag
Poliovirus	PVR
Rhinovirus	ICAM-1

Viral Replication: Basic Concepts

- Replication cycle produces
 - Functional RNA's and proteins
 - Genomic RNA or DNA and structural proteins
- 100's-1,000's new particles produced by each cycle
 - Referred to as burst size
 - Many are defective
 - End of 'eclipse' phase
- Replication may be cytolitic or non-cytolytic

Steps in Viral Replication: Penetration (Second Step)

- Enveloped viruses penetrate cells through fusion of viral envelope with host cell membrane
 - May or may not involve receptor mediated endocytosis
- Non enveloped viruses penetrate by
 - Receptor mediated endocytosis
 - Translocation of the virion across the host cell membrane

Influenza Virus Replication Cycle




FIG. 13. Schematic diagram of the life cycle of influenza virus. See text for details of the model.

From Fields *Virology*

Steps in Viral Replication: Basic Strategies of Transcription and Translation (Fourth and Fifth Steps)

- (+) RNA → Proteins
- (-) RNA → (+) RNA → Proteins
- RNA → DNA → RNA → Proteins
- DNA → RNA → Proteins

Steps in Viral Replication: Uncoating (Third Step)

- Makes viral nucleic acid available for transcription to permit multiplication to proceed
- Mechanism variably understood depending upon the virus

Steps in Viral Replication: Assembly and Release (Sixth and Seventh Steps)

- Process involves bringing together newly formed genomic nucleic acid and structural proteins to form the nucleocapsid of the virus
- Nonenveloped viruses exhibit full maturation in the cytoplasm or nucleus with disintegration of cell

Uncoating of Influenza Virus










FIG. 14. Schematic diagram of the proposed role of the M₂ ion channel activity in virus entry. The M₂ ion channel activity is thought to facilitate the flow of protons from the lumen of the endosome into the virion interior, bringing about dissociation of protein-protein interactions between the M2 cytoplasmic tail and M₂ and lipid and/or RNPs and M₂ from the RNPs.

From Fields *Virology*

Steps in Viral Replication: Assembly and Release (Sixth and Seventh Steps)


- Many enveloped viruses exhibit full maturation as the virion exits the cell
 - Viral proteins are inserted into the host cell membrane
 - Nucleocapsids bind to these regions and bud into the extracellular space
 - Further cleavage and maturation of proteins may occur after viral extrusion
 - Cytolytic activity of these viruses varies



Primary HIV Infection: Pathogenetic Steps


- Virus – dendritic cell interaction
 - Infection is typically with R5 (M-tropic) strains
 - Importance of DC-SIGN
- Delivery of virus to lymph nodes
- Active replication in lymphoid tissue
- High levels of viremia and dissemination
- Downregulation of virus replication by immune response
- Viral set point reached after approximately 6 months

The Variable Course of HIV-1 Infection



Reprinted with permission from Haynes. In: DeVita et al, eds. *AIDS: Etiology, Treatment and Prevention*. 4th ed. Lippincott-Raven Publishers; 1997:89-99.

PHI: Early Seeding of Lymphoid Tissue



Schacker T et al: *J Infect Dis* 2000;181:354-357


Primary HIV Infection: Determinants of Outcome

- Severity of symptoms
- Viral strain
 - SI (X4) vs. NSI (R5) viruses
- Immune response
 - CTL response
 - Non-CTL CD8 responses
 - Humoral responses?
- Viral set point at 6-24 months post-infection
- Other host factors
 - Chemokine receptor and HLA genotype
- Gender and differences in viral diversity?
- Antiviral therapy
 - Near vs. long-term benefit?


Primary HIV Infection: Clinical Characteristics

- 50-90% of infections are symptomatic
- Symptoms generally occur 5-30 days after exposure
- Symptoms and signs
 - Fever, fatigue, myalgias, arthralgias, headache, nausea, vomiting, diarrhea
 - Adenopathy, pharyngitis, rash, weight loss, mucocutaneous ulcerations, aseptic meningitis, occas. oral/vaginal candidiasis
 - Leukopenia, thrombocytopenia, elevated liver enzymes
- Median duration of symptoms: 14 days

Natural History of Untreated HIV-1 Infection



Antiviral Agents for HIV



Mechanism of T20/T1249 Mediated Fusion Inhibition

Modified from Weissenhorn et al., *Nature* 387, 426-430 (1997)
and Furuta et al., *Nature structural biology* 5, 276-279 (1998).

