


Infective Endocarditis






Infective Endocarditis: Definitions

- A microbial infection of a cardiac valve or the endocardium caused by bacteria, fungi, or chlamydia
- Often categorized as acute or subacute based on the rapidity of the clinical course
 - Alternatively described by type of risk factor *e.g.*, nosocomial, prosthetic valve, intravenous drug use associated
- Pathological findings include the presence of friable valvular vegetations containing bacteria, fibrin and inflammatory cells. There is often valvular destruction with extension to adjacent structures.
 - Embolic lesions may demonstrate similar findings



Historical Perspective




Sir William Osler

- ".. A concretion larger than a pigeon's egg; contained in the left auricle." Burns, 1809
- Osler's Gulstonian lectures provided the 1st comprehensive overview of the disease
- Lewis and Grant (1923) were the first to link a transient bacteremia with deformed valves as the two predominant risk factors for infection
- The introduction of penicillin marked the first successful therapy for this otherwise lethal infection

Infective Endocarditis • Definitions and Historical Perspective • Epidemiology • Pathogenesis • Clinical Presentation and Diagnosis • Therapy • Prevention

Epidemiology of Endocarditis

- Incidence the same or slightly increased - 1.7-6.2/100,000 depending on the population
- The age of subjects with endocarditis has increased over the past 60 years (30-40 to 47-69)
 - Among injecting drug users the incidence is as high as 150-2000/100,000 person years
- There has been a major shift in nature of underlying valvular disorders
- There has also been a change in the microbiology of cases – Increasing incidence of staphylococci



- There has been an increasing incidence of nosocomial endocarditis both native and prosthetic valve
- There is an increased risk of IE among injecting drug users, patients on long-term hemodialysis, patients with intravenous catheters, diabetics and HIV-infected patients

Incidence of Underlying Heart Disease in Infective Endocarditis*


Ī	Pre-Antibiotic Era	Present
Rheumatic	++++^	+
Congenital	+	+
Degenerative	+	++
Mitral valve prola	apse +	+++
Normal valves	+	++
*Incidence varies with p ^Relative incidence	population studied	

Endocarditis in Ur	ban Hospitals
<u>Organisms</u>	Percent Cases
Streptococcus spp.	34
Enterococcus spp.	6
Staphylococcus aureus	40
Coagulase-negative staphylococci	5
Gram negative bacilli	6
Fungi	2
Misc. / Polymicrobial	3
Culture negative	4

Risk Factors for Infective Endocarditis


- Dental procedures, poor dental hygiene viridans streptococci, nutritionally variant streptococci, HACEK
- Prosthetic valves
 - Early: coagulase negative staphylococci, S. aureus
 - Late: coagulase negative staphylococci, viridans streptococci
- Gastrointestinal or genitourinary procedures enterococci or *S. bovis* (colon carcinoma)
- Nosocomial *S. aureus* (including MRSA), Gram negatives, Candida species

Brouqui and Raoult, Clin Microbiol Rev, 2001





Infective Endocarditis


- Definitions and Historical Perspective
- Epidemiology
- Pathogenesis
- Clinical Presentation and Diagnosis
- Therapy
- Prevention




- Inoculation of bacteria colonizing a mucosal (*e.g.*, oral mucosa) or peripheral tissue site into the bloodstream
- Transient bacteremia of a serum-resistant pathogen capable of adhering to a cardiac valvular surface
- Turbulent blood flow across the valve
- Bacterial adherence to cardiac valvular surface
- Pathogen host tissue interaction resulting in vegetation formation and local tissue damage
 - Bacterial persistence
- Dissemination of infection to other tissue sites and elicitation of systemic findings












Patient 3: Colonies per cubic centimeter in blood from femoral artery and antecubital vein.					
hainteiler	Femoral	Antecubita			
Time	artery	vein			
<i>p.m.</i>					
1:00	266	228			
1:03	264	251			
1:05	300	285			
1:07	358	256			
1:08	399	397			
1:11	340	358			
1:13	397	344			
1:15	298	318			
1:17	332	342			
1:20	552	533			
1:23	671	534			









Immunologic Manifestations of Infective Endocarditis

- Hypergammaglobulinemia; both antigen specific and polyclonal B cell activation (*e.g.*, rheumatoid factor)
 - May block IgG opsonic response, accelerate microvascular damage or stimulate phagocytosis
- Vasculitis
 - Circulating immune complexes
 - Hypocomplementemia
- Clinical syndromes: "Lumpy-Bumpy" glomerulonephritis with deposition of complexes plus complement, Osler's nodes



Infective Endocarditis

- Definitions and Historical Perspective
- Epidemiology
- Pathogenesis
- Clinical Presentation and Diagnosis
- Therapy
- Prevention




- Valvular destruction and local intracardiac complications
- Bland or septic embolization of vegetations
- Sustained bacteremia
- Immunologic phenomena


Osler, Gulstonian Lectures, Lancet, 1885 Weinstein and Schlesinger NEJM, 1974

Case 1: One Sick Dude

A 33 year-old recreational drug user develops fever, chills and pleuritic chest pain 48h after injecting heroin. He appears acutely toxic with a temperature of 104°, rapid respirations and agitation. His sputum is purulent and bloody. His cardiac exam is unremarkable except for his sinus tachycardia. Examination of his skin reveals a petechial rash and a small soft tissue abscess at his injection site.









Case 1: One Sick Dude

His blood cultures are positive for *S. aureus* within two days. One week into therapy he is feeling better, however he has noticed some increased left upper quadrant pain and on examination you are able to palpate his spleen. He also has developed a soft systolic murmur at his left lower sternal border that is increased with inspiration.



The Sluggish Doc

A 65 year old female physician has noted increasing fatigue and malaise for the past 1-2 months. When she takes her temperature she is surprised to find that it is 100.6°. She also notes some new lesions on her arms and nails. She is too busy to see her own physician. What finally brings her to medical attention is the development of red, painful swelling on the dorsum of her left foot. Her physician notes Roth spots in both eyes and a loud mitral insufficiency murmur. Of note, she has a longstanding history of mitral valve prolapse.











The Sluggish Doc

Multiple blood cultures grow viridans streptococcus. Her laboratory studies reveal a positive rheumatoid factor test. Despite antibiotic therapy, she develops refractory heart failure and requires mitral valve replacement












Transesophageal Echocardiography







Duke Criteria for the Diagnosis of Endocarditis

• Definite:

- Pathologic criteria are histologic or culture confirmation of vegetation or emboli.
- Clinical criteria are 2 major, 1 major plus 3 minor or 5 minor criteria (*e.g.* positive blood cultures or echocardiogram, new regurgitant murmur)
- Possible:
 - 1 major, 1 minor; or 3 minor criteria
- Rejected:
 - Alternate diagnosis, resolution of infection with brief therapy, no pathologic evidence of endocarditis

Durack et al. AJM, 1994; Li et al. Clin Infect Dis, 2000

Mimics of Infective Endocarditis

- Atrial myxoma
- Marantic endocarditis
- Left atrial thrombus
- Acute rheumatic fever with carditis
- Collagen vascular disease (SLE)
- Neoplasms (carcinoid)



- Definitions and Historical Perspective
- Epidemiology
- Pathogenesis
- Clinical Presentation and Diagnosis
- Therapy
- Prevention

Principles of Therapy

- Bactericidal antibiotics must be used
- Prolonged therapy is necessary (weeks)
- Treatment is best started after multiple sets of blood cultures have been taken.
- Urgency in the initiation of therapy is required for acute but not subacute endocarditis.
- Synergistic combinations of antibiotics are used when available.

Infective Endocarditis

- Definitions and Historical Perspective
- Epidemiology
- Pathogenesis
- Clinical Presentation and Diagnosis
- Therapy
- Prevention


Antimicrobial Prophylaxis of Endocarditis – Potential Mechanisms


- Bactericidal activity
- Reduce bacterial adherence
- Reduce bacterial density in the wound at the time of surgery (for prosthetic valves)

Prevention of Infective Endocarditis

- High risk
 - Prosthetic valve
 - Complex congenital heart disease
 - Previous endocarditis
- Moderate risk
 - Acquired valvular dysfunction (*e.g.* rheumatic valve)
 - Mitral valve prolapse with regurgitation
- Negligible risk
 - Mitral valve prolapse without regurgitation
 - Rheumatic fever without valvular dysfunction

AHA Recommendations, JAMA 277:1794, 1997





What Do I Need to Know?

- What are the major epidemiologic features of endocarditis and how have they changed?
- Which bacteria are associated with which types of exposure in the development of endocarditis?
- What is the pathogenesis of endocarditis and what is its correlation with the different clinical presentations?
- How is endocarditis diagnosed?
- What are the basic principles of therapy and prophylaxis?