

VZV, EBV, and HHV-6-8

Anne Gershon

MID 35

Common Features of Herpesviruses

- Morphology
- Basic mode of replication
- Primary infection followed by latency
- Ubiquitous
- Ability to cause **recurrent** infections (reactivation of latent virus), **reinfections** (with a new virus), **persistent** infections (chronic low grade virus multiplication) **immortalizing infections** (EBV only)


8 Human Herpesviruses, 3 categories

- Alpha: short reproductive cycle, variable host range, latent in sensory neurons
 - Herpes simplex virus (HSV 1, 2)
 - Varicella-zoster virus (VZV)
- Beta: long reproductive cycle, narrow host range, latent in lymphoid cells & others (salivary glands, kidney)
 - Cytomegalovirus (CMV)
 - HHV6, HHV 7
- Gamma: narrow host range; latent in lymphoid cells, associated with tumors
 - Epstein Barr Virus (EBV)
 - Kaposi Sarcoma Virus (KSH, HHV8)

Human Herpesviruses

- Replication (lytic infection) occurs in a cascade
 - Latency occurs when the cascade is interrupted
- Transcription of viral genome and protein synthesis (cascade of gene expression), essential and luxury
 - 1. immediate early (IE): regulation of gene expression, DNA binding
 - 2. early (E): more transcription factors, enzymes, DNA polymerase
 - 3. late (L): structural proteins
- Encode targets for antiviral therapy
 - TK, DNA polymerase


The rash of VZV is vesicular.




- Vesicular fluid is highly infectious.
 - Well-formed virions are suspended in it.

MID 35

Human Herpesvirus (VZV)
phospholipid envelope, tegument, icosahedral
capsid, DNA core



VZV is a typical herpesvirus



VZV has a phospholipid envelope.

Contains gps, eg: gB, gC, gE, gH, gI, gK, gL

Tegument

Contains immediate early and early proteins, eg: ORFs 4, 10, 21, 62, 63

Nucleocapsid


ORFp40

DNA core


MID 35

Varicella-zoster virus


- The smallest of the herpesviruses
 - 125,000 base pairs
 - 70 Open reading frames (ORFs)
- Receptors: heparan sulfate, mannose-6 phosphate receptor (MPR), insulin degrading enzyme (IDE)




MPRs sort lysosomal enzymes and target them to endosomes



Steps in the assembly and intracellular transport of VZV







VZV
Receives
Its Final
Envelope
in the
TGN

MID 35


Hypothesis: VZV latency is established by free virions that infect sensory nerve endings



VZV spreads in two ways

MID 35

Pathogenesis of Zoster



Adapted from Hope-Simpson. *The Proceedings of the Royal Society of Medicine*. 1966;58:5.

In the body VZV spreads from cell-to-cell

- In varicella, VZV is transported from the respiratory mucosa to the blood (viremia) in T cells, where virus is not accessible to antibodies.
 - Because cell-to-cell spread is slow, the incubation period of varicella is long (2 weeks).
 - Slow spread prevents host from being overwhelmed before the immune response develops
- T helper (Th1) and cytotoxic T cells are required for host control of virus

MID 35

Natural History of VZV

- Primary infection: varicella
 - Highly contagious (airborne)
 - Complications: bacterial superinfection, encephalitis, pneumonia, congenital syndrome
- Secondary infection: zoster
- Zoster is due to reactivation of latent VZV
 - DNA, RNA, proteins in ganglia at autopsy
 - Zoster in a few vaccinees caused by Oka vaccine
 - From low cell-mediated immunity (CMI) to VZV
- No asymptomatic shedding of VZV as with HSV

MID 35



Varicella is a generalized illness. Infectious virions are produced in the skin vesicles.

35

Zoster is initially localized.



- Limited to 1-3 dermatomes.
- May disseminate in immunocompromised hosts.



Congenital
varicella
syndrome



Fatal neonatal
varicella

MID 35

Zoster in a 3 month old



MID 35

VZV In the Immunocompromised

- **Varicella is likely to be severe**
 - Prevent or modify with pre-formed antibodies just after exposure
 - Virus spreads from cell-cell in body
 - requires CMI (cellular immunity) for host defense
 - Treat most immunocompromised patients immediately with acyclovir
- **The frequency of zoster is increased**
 - Probably related to low CMI response
 - Likely to suffer post-herpetic neuralgia (PHN) (also elderly)

MID 35

Latent Infection with VZV

- Latent infection in dorsal root ganglia (DRG)
- 6 of 68 genes (also RNA and proteins) expressed during latency
- Proteins of regulatory genes are expressed in cell cytoplasm, not nucleus
- Suggests regulatory proteins are blocked from normal action, leading to inhibition of cascade of gene expression preventing lytic infection from occurring (latency)
- Latency is established when cell-free VZV in skin vesicles invades neurons

MID 35

Varicella Vaccine

Only herpesvirus for which there is a vaccine

Live, attenuated, infectious virus (Oka strain)

Licensed for routine use in healthy susceptible individuals in US, in 1995

Recently there has been a marked decrease in varicella, in all age groups

– Indicates herd immunity

- Contraindications: pregnancy, immunocompromised, allergy to vaccine components

MID 35

Varicella Vaccine

Live attenuated; stimulates primary immunity

Major complaint afterwards: mild rash in 5%

- 1 month after vaccination; transmission is rare
- Vaccine is extremely safe

85% completely protected; 15% partial immunity


There is little evidence for waning immunity

Subsequent zoster is rare

Same vaccine (much higher dose) also used successfully to prevent zoster in the elderly (different mechanism of action... stimulates CMI to VZV)


MID 35

Varicella, Antelope Valley, CA



MID 35


The rash of VZV is vesicular.



- Vesicular fluid is highly infectious.
 - Well-formed virions are suspended in it.

MID 35

Indirect immunofluorescence To diagnose VZV, HSV



MID 35

Laboratory Methods for Diagnosis

- Culture (difficult), DFA, PCR, cytology on skin rash (Tzanck)
 - Can distinguish the Oka virus from wild type virus (PCR)
- Antibody titers, IgG (ELISA)
 - Acute serum, early in illness
 - Convalescent serum, 10-14 days after onset
- Antibody titers, IgM
 - False positives and false negatives can be a problem

Acyclovir (ACV) is useful to treat HSV, VZV

- Antiviral activity only in infected cells (TK)
- Sensitivity: HSV1, >HSV2, >VZV (EBV, CMV)
- Toxicity is unusual: gastrointestinal, neurologic (headache, seizures, delirium); anemia, thrombocytopenia, bone marrow suppression
- Resistance is a concern, especially in HIV-infected patients
- Newer drugs: famciclovir, valacyclovir
 - Administered orally and less frequently than ACV because better gastrointestinal absorption

Epstein-Barr Infections (EBV)

- Infectious mononucleosis, nasopharyngeal carcinoma, lymphomas (including Burkitt's), oral hairy leukoplakia (lytic infection), X-linked proliferative disease (males only)
- **B cells are latently infected in mononucleosis**; T cells (atypical lymphocytes) are the host response
- **Latency develops in memory B cells**
- EBV is not related to chronic fatigue syndrome, but rarely severe chronic illness follows mononucleosis
- **In mononucleosis, give steroids if airway obstruction, hemolytic anemia, severe cardiac, neurologic disease (no specific antiviral therapy)**

Diagnosis of Mononucleosis

- Usually occurs in young adults
- **Symptoms, signs:** fever, adenopathy, exudative pharyngitis, rash (ampicillin) hepatosplenomegaly fatigue
- **Positive heterophile antibody** (monospot)
- **EBV specific antibodies**
 - Anti VCA (develops **early**, persists)
 - Anti EBNA (develops **late**, persists)
 - **Positive aby VCA, neg aby EBNA = acute mononucleosis**

Herpesviruses 6, 7

- **Herpesvirus 6 (beta, like CMV)**
 - Roseola in infants (rash, fever, seizures)
 - outcome of latency in CNS not understood
 - Fevers in immunosuppressed
 - Rare mononucleosis syndrome in adults
- **Herpesvirus 7 (beta, like CMV)**
 - Fevers in immunocompromised (HIV)
- **Diagnosis, treatment are not fully developed**
 - Most infections are self-limited

Herpesvirus 8 (KHSV)

- Closely related to EBV
- Encodes for human proteins (piracy)
 - IL-6, Bcl-2 (anti-apoptosis), chemokines
- Infections are rare in children
 - Can cause non-specific fever and rash illness
- Causes Kaposi's Sarcoma
 - Elderly
 - HIV-infected
- Causes primary-effusion lymphoma
- Castleman's disease (lymphoma-like)

Summary: Herpesvirus Infections

- Particularly affect newborns, elderly, immunocompromised
 - Congenital (**CMV**, VZV) vs neonatal (**HSV**, VZV); primary maternal infections high risk
- Primary, latent, recurrent, reinfections
- Best diagnostic tool: PCR
- Antiviral therapy: HSV, VZV, CMV
- EBV and HHV8 cause tumors
- Vaccine now available against to prevent chickenpox (varicella) and zoster (shingles)