

# AIDS at 25

## Epidemiology and Clinical Management

**Regional HIV and AIDS statistics and features, 2006**

	Adults & children living with HIV	Adults & children newly infected with HIV	Adult (15-49) prevalence [%]	Adult & child deaths due to AIDS
Sub-Saharan Africa	24.7 million (21.8 - 27.7 million)	2.8 million (2.4 - 3.2 million)	5.9% (5.2% - 6.6%)	2.1 million (1.8 - 2.4 million)
Middle East & North Africa	460 000 (270 000 - 760 000)	66 000 (41 000 - 220 000)	0.2% (0.1% - 0.4%)	36 000 (20 000 - 60 000)
South and South-East Asia	7.8 million (5.2 - 12.6 million)	860 000 (560 000 - 2.3 million)	0.6% (0.4% - 1.0%)	590 000 (380 000 - 850 000)
East Asia	750 000 (480 000 - 1.2 million)	100 000 (56 000 - 300 000)	0.1% (0.07% - 0.2%)	43 000 (26 000 - 64 000)
Latin America	1.7 million (1.3 - 2.5 million)	140 000 (100 000 - 410 000)	0.5% (0.4% - 0.7%)	65 000 (51 000 - 84 000)
Caribbean	250 000 (190 000 - 320 000)	27 000 (20 000 - 41 000)	1.2% (0.9% - 1.6%)	19 000 (14 000 - 26 000)
Eastern Europe & Central Asia	1.7 million (1.2 - 2.6 million)	270 000 (170 000 - 810 000)	0.9% (0.6% - 1.4%)	84 000 (58 000 - 120 000)
Western & Central Europe	740 000 (580 000 - 970 000)	22 000 (18 000 - 33 000)	0.3% (0.2% - 0.4%)	12 000 (11 000 - 24 000)
North America	1.4 million (880 000 - 2.2 million)	43 000 (34 000 - 60 000)	0.6% (0.5% - 0.8%)	18 000 (11 000 - 29 000)
Oceania	81 000 (50 000 - 170 000)	7 100 (3 400 - 54 000)	0.4% (0.2% - 1.0%)	4 000 (2 000 - 6600)
<b>TOTAL</b>	<b>39.5 million</b> (34.1 - 47.1 million)	<b>4.3 million</b> (3.6 - 6.6 million)	<b>1.0%</b> (0.9% - 1.2%)	<b>2.9 million</b> (2.5 - 3.5 million)

AIDS Epidemic Update, November 2006, Table 1b

250 MMWR June 5, 1981

**Dengue - Continued**

**Editorial Note:** Dengue type 4 frequently occurs in Southeast Asia, the South Pacific, and Africa. How it was introduced onto St. Barthelemy, a small and relatively remote island in the Caribbean, remains unknown. However, French health authorities have reported to CAREC that an outbreak of dengue-like illness has been observed on St. Barthelemy, beginning in February or March, but has since declined. In the absence of reports of an ongoing outbreak of dengue in the Caribbean, the risk that travelers to this area will acquire dengue is probably small.


Dengue types 2 and 3 have been present in the Caribbean at least since the 1960s. Dengue type 1 was first recognized in that area when an outbreak in Jamaica in 1977 was followed by numerous outbreaks on other Caribbean islands and in Central America. All these dengue types, as well as type 4, usually cause an illness that is clinically mild and typically of short duration.

**Pneumocystis Pneumonia - Los Angeles**


In the period October 1980-May 1981, 5 young men, all active homosexuals, were treated for biopsy-confirmed *Pneumocystis carinii* pneumonia at 3 different hospitals in Los Angeles, California. Two of the patients died. All 5 patients had laboratory-confirmed previous or current cytomegalovirus (CMV) infection and candidal mucosal infection. Case reports of these patients follow.







**Patient 1:** A previously healthy 33-year-old man developed *P. carinii* pneumonia and oral mucosal candidiasis in March 1981 after a 2-month history of fever associated with elevated liver enzymes, leukopenia, and CMV viraemia. The serum complement-fixation CMV titer in October 1980 was 256; in May 1981 it was 32.\* The patient's condition deteriorated despite courses of treatment with trimethoprim-sulfamethoxazole (TMP-SMX), pentamidine, and acyclovir. He died May 3, and postmortem examination showed transient alveolar






Reported by MS, Div of Clinical Microbiology, Cedars Mt. Sinai H. Editorial Note: limited to severe immunodeficiency in association with sexual contact in this report has been reported elsewhere.



- ## HIV Transmission
- Blood
    - transfusion
    - injection drug use
  - Sexual Intercourse
    - heterosexual
    - male to male
  - Perinatal
    - intrapartum
    - breast feeding







- ### Frequency of HIV 'Non-Progressors'
- San Francisco City Clinic Cohort
    - 489 HIV+ Gay men with known seroconversion date.
    - 13% developed AIDS by 5 years;
    - 51% developed AIDS by 10 years.
    - 89% had died, developed AIDS or had CD4<500 by 10 years.
- [Rutherford et al. *BMJ.* 1990; 301:1183-8 ]

## Explaining the variability of HIV disease

- Viral Factors
  - Nef deletion
  - Non-clade B subtypes?
- Host Factors
  - Chemokine co-receptors
  - Immune response
  - Gender?
- Environmental Factors
  - Infection, diet?, stress?

## AIDS Restriction Genes

Table 2 Genes that limit AIDS

Gene	Allele	Mode	Effect	Mechanism of action	Reference
HIV entry	CCR5 Δ32	Recessive	Prevent infection	Knockout CCR5 expression	17
	CCR5 Δ32	Dominant	Prevent lymphoma (L)	Decrease available CCR5	90
	CCR5 Δ32	Dominant	Delay AIDS	Decrease available CCR5	17
CCR5	P1	Recessive	Accelerate AIDS (E)	Increase CCR5 expression	34
CCR2	Δ64	Dominant	Delay AIDS	Interact with and reduce CXCR4	38,39
CCL5	in1,1c	Dominant	Accelerate AIDS	Decrease RANTES expression	45
CXCL12	3'A	Recessive	Delay AIDS (L)	Impede CCR5-CXCR4 transition (T)	46
CXCR6	EDK	Dominant	Accelerate PCP (L)	Alter T-cell activations (T)	48
CCL3-CCL7-CCL11	H7	Dominant	Enhance infection	Stimulate immune response (T)	49
Cytokine with HIV	IL10 5'A	Dominant	Limit infection	Decrease IL10 expression	53
	IL10 5'A	Dominant	Accelerate AIDS	Decrease IL10 expression	53
IFNG	-179T	Dominant	Accelerate AIDS (E)		55
Acquired immunity, cell mediated	HLA A,B,C	Homozygous	Accelerate AIDS	Decrease breadth of HLA class I epitope recognition	62,66
	B*27	Codominant	Delay AIDS	Delay HIV-1 escape	9
	B*57	Codominant	Delay AIDS	Delay HIV-1 escape	9
	B*35-Fx	Codominant	Accelerate AIDS	Defect CD8-T cell clearance of HIV-1	60
Acquired immunity, innate	KIR3DS1	Epistatic with HLA-B*57	Delay AIDS	Clear HIV*, HLA* cells (T)	70

S. O'Brien, G. Nelson. *Nature Genetics* 2004;36:565

## HIV Co-receptors

CD4 necessary but not sufficient for infection.  
Beta chemokine receptors act as HIV co-receptors.

CXCR4 (lymphocyte) CCR5 (macrophage)

Homozygous CCR5 deletion found in <1%.

MACS High risk cohort:

No HIV+ among those homozygous for deletion.


3.6% of HIV Negative were homozygous.

Among persistently HIV Neg: up to 33% were homozygous.

## Early indicators of HIV Infection



## Effect of Co-receptor Heterozygosity




## Key features of OIs in AIDS

- HIV causes profound defect mostly restricted to T cell-based immunity (restricted range of pathogens)
- OIs usually reflect reactivation of latent infections.
- Reinfection may occur (eg: tuberculosis)
- Chronic suppression needed after acute treatment.
- Immune reconstitution with anti-retroviral therapy may reverse OI susceptibility (but may also trigger an inflammatory response to active OIs)


## Pneumocystis pneumonia in AIDS

- Commonest life threatening complication of AIDS in U.S.
- Subacute illness (fever, cough, dyspnea).
- Diffuse interstitial infiltrate on x-ray.
- Addition of corticosteroids to antimicrobials cuts mortality in severe disease 50%.
- Fully preventable with trimethoprim-sulfa.



## Cryptococcal disease in AIDS

- Ubiquitous soil fungus.
- Initial asymptomatic pneumonia.
- Reactivation in advanced HIV disease (CD4<100).
- Meningitis commonest presentation but wide dissemination frequent.




## CD4 count predicts risk of PCP

TABLE 1. Cumulative incidence\* of *Pneumocystis carinii* pneumonia (PCP) according to CD4+ count at baseline among the MACS seroprevalent cohort†


CD4+ count at baseline	N	PCP	Percentage with PCP		
			6 mo.	12 mo.	36 mo.
< 200	77	19	8.4	18.4	33.3
201-350	217	47	0.5	4.0	22.9
351-500	389	39	0.0	1.4	9.0
501-700	483	43	0.0	0.4	8.3
> 700	499	20	0.0	0.0	3.8

## CMV disease in AIDS

- Common viral infection (50% adult seroprevalence).
- Reactivation at CD4<50
- Retinitis commonest.
- Other sites: Colon, CNS.




## CNS toxoplasmosis



- Protozoan parasite; cats shed oocysts; farm animals incidental hosts; humans infected from cysts, uncooked meat.
- Commonest cause of focal CNS disease in AIDS.
- Serum IgG antibody reliable marker of past infection.
- Reactivation in AIDS associated with CD4<100.

## Disseminated Mycobacterium-avium complex (MAC) disease in AIDS |


- Common in environment (water).
- Local lung disease known prior to AIDS.
- Widespread visceral dissemination in AIDS.
- Diagnosis by blood culture.
- Absence of inflammation in tissue sites.



### Prophylaxis of Opportunistic Infections

Pathogen	Indication	Regimen
PCP	CD4<200	Trimethoprim-sulfa
Toxo	CD4<100 and IgG+	Trimethoprim-sulfa or Dapsone +Pyrimethamine
MAC	CD4<50	Clarithro/Azithromycin
TB	+PPD (5mm)	INH (9 months)


### MAC IRIS simulating TB or Lung cancer



### OI Guidelines November, 2001 Comparison of Indications to Discontinue Primary and Secondary Prophylaxis

Agent	Recommendation
PCP	1° CD <sub>4</sub> > 200 X 3 months 2° CD <sub>4</sub> > 200 X 3 months
Toxo.	1° CD <sub>4</sub> > 200 X 3 months 2° CD <sub>4</sub> > 200 X 6 months + initial Rx + asymptomatic
MAC	1° CD <sub>4</sub> > 100 X 3 months 2° CD <sub>4</sub> > 100 X 6 months + 12 mo Rx + asymptomatic

### CNS crypto IRIS



### Immune Reconstitution with HIV Therapy

- Focal MAC adenitis
- Inflammatory flare of CMV retinitis
- Worsening of previously stable hepatitis
- Development of cavitary TB