

Transplantation Immunology

Mitchell S. Cairo, MD

Professor of Pediatrics, Medicine and Pathology
Chief, Division, Pediatric Hematology &
Blood & Marrow Transplantation
Children's Hospital New York Presbyterian
Director Leukemia, Lymphoma, Myeloma Program
Herbert Irving Comprehensive Cancer Center
Columbia University
Tel – 212-305-8316
Fax – 212-305-8428
E-mail – mc1310@columbia.edu

Types of Grafts

- Autologous (self)
 - e.g., BM, peripheral blood stem cells, skin, bone
- Syngeneic (identical twin)
- Allogeneic (another human except identical twin)
- Xenogeneic (one species to another)


Rejection

- First Set Rejection
 - Skin graft in mice 7-10 days
- Second Set Rejection
 - Skin graft in mice in 2-3 days


Mechanisms

- Foreign alloantigen recognition
- Memory lymphocytes (adaptive immunity)
- Can be adoptively transferred

MHC Restricted Allograft Rejection




First & Second Allograft Rejection




AlloAntigen Recognition

- Major Histocompatibility Complex (MHC)
 - Class I HLA A, B, C bind to TCR on CD8 T-Cell
 - Class II DR, DP, DQ bind to TCR on CD4 T-Cell
 - Most polymorphic genes in human genome
 - Co-dominantly expressed
- Direct presentation (Donor APC)
 - Unprocessed allogeneic MHC
- Indirect presentation (Host APC)
 - Processed peptide of allogeneic MHC


Map of Human MHC




T-Cell Recognition of Peptide-MHC Complex




Direct and Indirect AlloAntigen Recognition



T-Cell Anergy vs T-Cell Activation



Antigen Recognition & Immunological Synapse



Mixed Lymphocyte Reaction (MLR)


• Definition & Mechanism

- *In vitro* test of T-cell regulation of allogeneic MHC
- Stimulators (donor-irradiated mononuclear cells)
- Responders (recipient mononuclear cells)
- Measure proliferative response of responders (tritiated thymidine incorporation)

• Requirements

- Can be adoptively transferred
- Require co-stimulation
- Require MHC
- Require Class I differences for CD8 T-cell response
- Require Class II differences for CD4 T-cell response

Mixed Lymphocyte Reaction (MLR)



Pathological Mechanism of Rejection


Solid Organ

- Hyperacute
 - Minutes to hours
 - Preexisting antibodies (IgG)
 - Intravascular thrombosis
 - Hx of blood transfusion, transplantation or multiple pregnancies
- Acute Rejection
 - Few days to weeks
 - CD4 + CD8 T-Cells
 - Humoral antibody response
 - Parenchymal damage & Inflammation
- Chronic Rejection
 - Chronic fibrosis
 - Accelerated arteriosclerosis
 - 6 months to yrs
 - CD4, CD8, (Th2)
 - Macrophages


Bone Marrow/PBSC

- Not Applicable
- Primary Graft Failure
 - 10 – 30 Days
 - Host NK Cells
 - Lysis of donor stem cells
- Secondary Graft Failure
 - 30 days – 6 months
 - Autologous T-Cells
 - CD4 + CD8
 - Lysis of donor stem cells

Immune Mechanisms of Solid Organ Allograft Rejection



Hyperacute, Acute, Chronic Kidney Allograft Rejection




Hyperacute

Acute

Acute

Chronic

Mechanisms of Acute Allograft Rejection




Prevention & Treatment of Allograft Rejection

- ABO Compatible
 - Prevent hyperacute rejection in solid organs
 - Prevent transfusion reaction in BM/PBSC
- MHC allele closely matched
- Calcineurin inhibitors
 - Cyclosporine binds to Cyclophilin
 - Tacrolimus (FK506) binds to FK Binding Proteins (FKBP)
 - Calcineurin activates Nuclear Factor of Activated T-Cells (NFAT)
 - NFAT promotes expression of IL-2
- IMPDH Inhibitors (Inosine Monophosphate Dehydrogenase)
 - Mycophenolate Mofetil (MMF)
 - Inhibits guanine nucleotide synthesis
 - Active metabolite is Mycophenolic acid (MPA)


Prevention & Treatment of Allograft Rejection

- Inhibition of mTOR
 - Rapamycin binds to FKBP
 - Inhibits mTOR
 - Inhibits IL-2 signaling
- Antibodies to T-Cells
 - OKT3 (Anti-CD3)
 - Daclizumab (Anti-CD25)
- Corticosteroids
 - Prednisone/Solmedrol
 - Inhibits Macrophage Cytokine Secretion
- Anti-inflammatory
 - Infliximab (Anti-TNF- α Antibody)
- Blocks B7 Co-Stimulation
 - CTLA-4-Ig
 - Inhibits T-cell Activation
 - Induces Tolerance
- Block CD40 Ligand Binding
 - Anti CD40 Ligand
 - Inhibits Macrophage & Endothelial Activation

Incidence of Renal Allograft Survival in Influenced by HLA Matching




Mechanism of T-Cell Activation vs Tolerance




Immunological Tolerance

- Immunological specific recognition of self antigen by specific lymphocytes
- Central tolerance (Thymus-derived)
 - Negative selection of autoreactive T-Cells
 - Regulation of T-Cell development
- Peripheral Tolerance
 - Clonal anergy (Inadequate co-stimulation)
 - Deletion (Activation-induced cell death)
 - Regulatory / Suppressor Cells (Inhibit T-Cell activation / proliferation)


Central T-Cell Tolerance Mechanisms (Deletion and Regulatory T-Cells)



Mechanism of T-Cell Inactivation (CTLA-4/B7 Interaction)



Mechanism of T-Cell Inhibition (Regulatory T-Cells)



General Indications of Blood and Marrow Transplantation

- Dose intensity for malignant tumor (DI)
- Graft vs Tumor (GVT)
- Gene replacement
- Graft vs Autoimmune (GVHI)
- Gene therapy
- Marrow failure

Specific Indications (Pediatric)

Malignant

- Leukemia
- Solid Tumors
- Lymphomas

Specific Indications (Pediatric)

Non-Malignant

Marrow Failure	Metabolic Disorders
Hemoglobinopathy	Histiocytic
Immunodeficiency	Autoimmune

Conditioning Therapy

Myeloablative – TBI Based

Myeloablative - Non TBI Based

Non-Myeloablative

Engraftment

- **Myeloid** Absolute neutrophil count $\geq 500/\text{mm}^3$ x 2 days after nadir
- **Platelet** Platelets $\geq 20 \text{ k}/\text{mm}^3$ x 7 days untransfused after nadir

Chimerism (Allogeneic)

- **Fluorescence *in situ* Hybridization (FISH)** (Sex mismatch)
- **VNTR** (Molecular)

Complications (Acute)

- Graft failure (GF)
- Hemorrhagic cystitis
- Graft vs Host Disease (GVHD)
- Infections
- Mucositis
- Persistent and/or recurrent disease
- Veno-occlusive disease (VOD)

Essential Components Required for GVHD

- Immuno-incompetent host
- Infusion of competent donor T-cells
- HLA disparity between host and donor

Graft vs Host Disease

- Hyperacute Day 0 – 7
- Acute Day 7 – 100
- Chronic Day 100 ≥

Acute Graft vs Host Disease

- **Dermal (Skin) :** Maculopapular
 Palms / Soles
 Pruritic ±
 Cheeks/ Ears/ Neck / Trunk
 Necrosis / Bullae
- **Hepatic :** Hyperbilirubinemia
 Transaminemia
- **Gastrointestinal :** Diarrhea
 Abdominal pain
 Vomiting
 Nausea


Risk Factors of GVHD

- HLA disparity 6/6 > 5/6 > 4/6
- Allo stem cell source MRD > UCB > UBM
- Donor Age
- Sex incompatibility
- CMV incompatibility
- Immune suppression

Common Prophylactic Immune Suppressants

- Methotrexate (MTX)
- Cyclosporine (CSP)
- Prednisone (PDN)
- Tacrolimus (FK506)
- Mycophenolate Mofetil (MMF)
- Anti Thymocyte Globulin (ATG)
- Alemtuzamab (Campath)
- T-Cell Depletion

Risk of Acute GVHD and HLA Disparity



Beatty et al *NEJM*: 313; 765, 1985

Chronic GVHD

- **Skin:** Rash (lichenoid, sclerodermatous, hyper/hypo pigmented, flaky), Alopecia
- **Joints:** Arthralgia, arthritis, contractures
- **Oral/Ocular :** Sjogren's Syndrome
- **Hepatic:** Transaminemia, hyperbilirubinemia, cirrhosis
- **GI:** Dysphagia, pain, vomiting, diarrhea, abdominal pain
- **Pulmonary:** Bronchiolitis obliterans (BO), Bronchiolitis obliterans Organizing Pneumonia (BOOP)
- **Hematologic/Immune:** Cytopenias, dysfunction
- **Serositis :** Pericardial, pleural

Summary

- First set donor tissue rejection from a non-identical MHC recipient is a primary adaptive immune response
- Second set donor tissue rejection for a non-identical MHC recipient involves memory antigen host T & B cells
- Alloantigen antigen direct and indirect presentation involves donor and host APC, respectively

Summary

- T-cell activation & proliferation requires immunological synapse with TCR/MHC and co-simulating ligands & receptors
- Tissue rejection maybe hyperacute (preexisting Ab) acute (days to weeks) and/or chronic (months to years)
- Allogenic stem cell transplantation may result in hyperacute (1-7d), acute (7-10d) and/or chronic (100d – 5yr) GVHD.