


The Innate Immune Response is Conserved
Throughout Evolution and is
Triggered by Pattern Recognition

Lipopolysaccharide = Lipid + Polysaccharide



From: Beutler and Rietschel, *Nature Reviews Immunology* 3: 169-176 (2003)

Diversity of "Pathogen-associated Molecular Patterns" (PAMPs)

QuickTime™ and a
TIFF (Uncompressed) decompressor
are needed to see this picture.

From: Akira et al., *Cell* 124:783, 2006

Innate Immune Receptors for PAMPs

- Toll-like receptors (TLRs)
- Complement
- Collectins (e.g., Surfactant Protein-A)
- Scavenger receptors
- Pentraxins (e.g., CRP)
- Lectins (e.g., Dectin-1)
- CD14
- NOD-like receptors (NLRs)
- RIG-1-like receptors

Collectins and Innate Immune Recognition

QuickTime™ and a TIFF (Uncompressed) decompressor are needed to see this picture.

QuickTime™ and a TIFF (Uncompressed) decompressor are needed to see this picture.

SP-A SP-D
Surfactant proteins

C1q


Globular heads bind to:

- Ag/Ab complexes
- Apoptotic cells
- Pathogens
- C-reactive protein

Collagen-like tails bind to:


- C1q receptors on phagocytes (to promote phagocytosis)
- C1r/C1s (to initiate the classical pathway)

Collectins Can Serve as Opsonins




Modified from: Wright, *Nature Rev. Immunol.* 5: 58, 2005



The Scavenger Receptor Superfamily Recognizes PAMPs



Innate Immune Receptors Also Trigger a Systemic Response to Infection




Innate Immune Functions of C Reactive Protein (CRP), an Acute Phase Protein Synthesized by Hepatocytes



CRP acts as a bridge between phosphocholine on bacterial targets and C1q

Adapted from: Black et al., *J. Biol. Chem.* 47:48487, 2004

Co-localization of CRP and Activated Complement in Infarcted Human Myocardium



CRP

C4d

J = jeopardized; N = normal


From: Nijmeijer et al., *Am. J. Pathol.* 163:269, 2003

Treatment of Experimental Myocardial Infarction with a CRP-binding Analog of Phosphocholine Limits Infarct Size


QuickTime™ and a TIFF (Uncompressed) decompressor are needed to see this picture.

From: Kitsis and Jialal, *New Engl. J. Med.* 355:513, 2006

History of Endotoxin Research



Modified from: Beutler and Rietschel, *Nature Reviews Immunology* 3; 169-176 (2003)



Cell
Volume 99 Number 6 September 20, 1996

Antifungal Defense in Drosophila

letters to nature

A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity

Ruslan Medzhitov, Paula Preston-Hurlburt & Charles A. Janeway Jr

*Section of Immunobiology, Yale University School of Medicine, and *Howard Hughes Medical Institute, New Haven, Connecticut 06510-3033, USA*

Induction of the adaptive immune response depends on the expression of co-stimulatory molecules and cytokines by antigen-presenting cells. The mechanisms that control the initial induction of these signals upon infection are poorly understood. It has been proposed that their expression is controlled by the non-clonal, or innate, component of immunity that preceded in evolution the development of an adaptive immune system in vertebrates. We report here the cloning and characterization of a human homologue of the *Drosophila* toll protein (Toll) which has been shown to induce the innate immune response in adult *Drosophila*. Like *Drosophila* Toll, human Toll is a type I transmembrane protein with an extracellular domain consisting of a leucine-rich repeat (LRR) domain, and a cytoplasmic domain homologous to the cytoplasmic domain of the human interleukin-1 (IL-1) receptor. Both *Drosophila* Toll and the IL-1 receptor are known to signal through the NF- κ B pathway. We show that a constitutively active mutant of human Toll (transfected into human cell lines) can induce the activation of NF- κ B and the expression of NF- κ B-controlled genes for the inflammatory cytokines IL-1, IL-6 and IL-8, as well as the expression of the co-stimulatory molecule B7.1, which is required for the activation of naive T cells.

The Toll protein controls signal-ventral patterning in *Drosophila* embryos and activates the transcription factor Dorsal upon binding to its ligand Spatzle. In adult *Drosophila*, the Toll/Dorsal signaling pathway participates in an anti-fungal immune response. Signaling through Toll initiates the signaling pathway induced by the IL-1 receptor (IL-1R) in mammalian cells. IL-1R signals through the NF- κ B pathway and Dorsal and its substrate Gata3 are homologous to NF- κ B and I- κ B proteins, respectively. Moreover, the cytoplasmic domain of *Drosophila* Toll is homologous to the cytoplasmic domain of IL-1R (ref. 9). Remarkably, the rubisco virus resistance gene that encodes N-protein is also similar to Toll in that it contains both a Toll signaling domain and an LRR domain. It thus appears that the immune-response system mediated by Toll represents an ancient host defence mechanism (Fig. 1). To investigate the possibility that this pathway has been retained in the immune system of vertebrates, we used sequence and pattern searches of the expressed sequence tag (EST) database at the National Center for Biotechnology Information (NCBI). A search with a sequence profile of the Toll/IL-1R signaling domain identified a matching sequence in the EST database derived from human fetal liver/placenta library (Genebank accession number U48462), corresponding to clone 320857 from the IMAGE consortium. Sequencing of this clone showed that it corresponds to the 3' untranslated region (UTR) and the C' terminus of the coding region of the messenger RNA. A fragment of the clone amplified by using the polymerase chain reaction (PCR) was used to screen a human spleen complementary DNA library by hybridization, and the 5' end of the cDNA was cloned using the 5' RACE technique as described^{11,12}. The full-length, 4.874 base pair (bp) cDNA clone contained an open reading frame of 2,523 bp by which encoded an

841-amino-acid protein chain (Fig. 2a), as well as a LINE-1 reverse transcription pseudogene in the 3'-UTR. Alignment of the sequences of the human and *Drosophila* Toll proteins shows that there is homology over the entire length of the protein chains (Fig. 2b). Notably, the similarity between the cytoplasmic domains is higher than between the human proteins Toll and IL-1R (not shown). The extracellular domain of human Toll (hToll) consists of 21 tandemly repeated leucine-rich motifs separated by a non-LRR region, similar to *Drosophila* Toll (dToll). At the N-terminal end of the LRR domain there is a 31-amino-acid long N-flanking region that is also present in several other LRR-containing proteins, for example BPI26, Decoyr and Biglycan. The C-terminal end of the LRR domain is flanked by a cysteine-rich domain which is also present in dToll and some other transmembrane proteins¹³.

To examine the expression pattern of hToll, a 720 bp cDNA fragment was used to probe Northern blots containing poly(A⁺) RNA from several organs. Most organs expressed two mRNA species one of ~5 kilobases (kb) was predominant in most tissues except peripheral blood leukocytes (PBL), and corresponded to the length of the cDNA that we cloned. The lower band was ~4 kb long and this band was undetectable in the PBL. The 4 kb band was not detectable in *Drosophila* and liver did not contain any mRNA at all (Fig. 3). We then hybridized different mouse and human cell lines for expression of hToll mRNA by using PCR with reverse transcription (RT-PCR). We found mRNA for hToll in monocytes, macrophages, dendritic cells, v8 T cells, Th1 and Th2 v8 T cells, a small intestinal epithelial cell line, and a B-cell line (data not shown). The hToll gene is expressed most strongly in spleen and PBL (Fig. 3); its expression in other tissues may be due to the presence of macrophages and dendritic cells, in which it could act as an early warning system for infection. Alternatively, hToll may be widely expressed because hToll signals through the conserved NF- κ B pathway (see below) and NF- κ B is a ubiquitous transcription factor.

To characterize hToll functions and see whether it can induce transcription of immune response genes like dToll, we generated a dominant positive mutant of hToll because the natural ligand of hToll is unknown. To produce a constitutively active mutant hToll, we made use of genetic information from dToll: analysis of ventralizing mutants in *Drosophila* embryos had identified the function of the cytoplasmic C'-flanking cysteine-rich region in dToll¹⁴ as controlling the activity of dToll in signal transduction. In three dominant





Figure 1 Ancient immune defence systems of plants, insects and vertebrates. A homologous immune response system based on the Toll signalling domain is used in plants, insects and vertebrates. Immune response genes required for the activation of both an innate and an adaptive immune response (see text). The figures are modified from ref. 4. Diagram illustrating responses to Toll signaling domain, striped dots, leucine-rich repeat domains, black rectangles, C-terminal cysteine-rich domains; white ovals, immunoglobulin domains.


A Re-interpretation of the Endotoxin Research Timeline



Modified from: Beutler and Rietschel, *Nature Reviews Immunology* 3: 169-176 (2003)

Primitive Specificity in Target Recognition by the Innate Immune System

Recruitment of TLR2 to Yeast Phagosomes




From: Underhill et al., *Nature* 401:811, 1999

Ligand Specificity of TLRs

QuickTime™ and a
PDF decompressor
are needed to see this picture.


Adaptor proteins, kinases

Specificity of TLR Transcriptional Programs



From: Luster, *Curr. Opin. Immunol.* 14:129, 2002

TLR Signaling: Two Major Pathways




Cartoon of major signal transduction pathways following engagement of TLRs. **TLR4** is the major sensor of LPS. **TLR3** recognizes dsRNA and is important in the anti-viral response. The **IRF** pathway leading to production of **Type I IFNs** (i.e., IFN- α/β) is particularly prominent in a minor subset of dendritic cells (called "plasmacytoid DCs") that are the major source of these IFNs in response to viral infections.

Do not memorize this cascade but rather appreciate that it consists of two parallel pathways, one that activates NF- κ B, leading to production of most pro-inflammatory proteins, and one that activates the IRF pathway, leading to production of Type I IFNs.

From: Moynagh, *Trends Immunol.* 26:469, 2005


TLRs Sense Microbial Pathogens and Trigger Expression of Pro-inflammatory Cytokines and Chemokines



Adapted from: Creagh and O'Neill, *Trends Immunol.* 27:352, 2006

Newly Recognized Components of the Innate Immune System

NOD Proteins: Intracellular Peptidoglycan Sensors



Polymorphisms in *Nod-2* are associated with up to 30-40% of cases of Crohn's disease (an inflammatory bowel disease)

CARD, caspase-recruitment domain; LRR, leucine-rich repeat; RICK, a CARD-containing protein kinase

Cytosolic Bacterial Recognition Systems and "the Inflammasome"

CARD = caspase-recruitment domain
LRR = leucine-rich repeat
NOD = NOD2-like domain
MDP = muramyl dipeptide

Cytoplasm

QuickTime™ and a
TIFF (Uncompressed) decompressor
are needed to see this picture.

From: Akira et al., *Cell* 124:783, 2006

Mutations in Pyrin, Another CARD-containing Innate Immune-like Protein, is Responsible for Familial Mediterranean Fever




Contrast-enhanced abdominal CT from a 31 year-old patient with Familial Mediterranean Fever suffering an acute attack of abdominal pain, nausea, vomiting, and arthritis. Note mesenteric vessel with thickened mesenteric fold (*white arrow*). Histopathology demonstrated neutrophilic infiltrate and associated vasculitis. Treatment with an IL-1 receptor antagonist (Anakinra) resulted in prompt cessation of symptoms.

Another Disease Associated with Activation of the Inflammasome




Pathogenesis of Gout Uncovered in 2006: Monosodium Urate Crystals Activate the Inflammasome



From: Martinon and Glimcher *J. Clin. Invest.* 116:2073, 2006

Nod-like Receptors (NLRs) Sense Microbial Products, Activate the "Inflammasome," and Trigger Maturation of IL-1




Adapted from: Creagh and O'Neill, *Trends Immunol.* 27:352, 2006

The Dendritic Cell and Development of The Primary Immune Response:

Wisdom Through Maturity


Dendritic Cell Maturation



From: Mellman & Steinman, *Cell* 106:255, 2001


Question: What Triggers Maturation of DCs?

The Innate Immune Response Orchestrates DC Trafficking to Secondary Lymphoid Organs




From: Luster, *Curr. Opin. Immunol.* 14:129, 2002


Functional Differences Between Immature and Mature DCs



The (Primary) Acquired Immune Response is Initiated by Innate Immune Recognition





Chemokines Direct Trafficking of Immune Cells




From: Luster, *Curr. Opin. Immunol.* 14:129, 2002

The Early Antiviral Response and the Innate Immune System



NK cells are a major source of a rapidly mobilizable pool of pro-inflammatory cytokines

Innate Immune Receptors for dsRNA Cooperate to Initiate the Immune Response to RNA Viruses




Double-stranded RNA products of virus infection bind to RIG-I or MDA5, which in turn bind to IPS-1 via CARD domain interactions. This complex then signals the activation of IKK- ϵ and TBK1 or other kinases to phosphorylate IRF-3, possibly through direct recruitment of signaling effectors, leading to IRF-3 dimerization, nuclear translocation and assembly onto the IFN- β enhancer. IPS-1 might also signal the activation of the IKK complex via direct binding of IKK components or through recruitment of RIP-1, FADD and/or TRAF6, causing the phosphorylation of I κ B, the inhibitor of NF- κ B. Phosphorylated I κ B is then ubiquitinated and targeted to the proteasome for degradation, releasing the active NF- κ B complex to translocate to the nucleus. During virus infection, dsRNA products can signal through TLR3 to activate IRF-3 and NF- κ B by the actions of the TRIF adaptor protein and RIP-1, respectively.

N.B.: Do not memorize this cartoon, but appreciate how cytosolic dsRNA receptors (RIG-1, MDA5) and plasma membrane-associated dsRNA receptors (TLR3) cooperate to activate IRF- and NF- κ B-dependent gene expression.

From: Johnson and Gale, *Trends Immunol.* 27:1, 2006

The Antiviral Response: a Cascade of Transcriptional Events




Some targets of IRFs

Gene	Function
p21	Cell cycle arrest
IL-15	NK cell maturation
FasL	Cell death
IL-12	Th1 immune response

Multiphasic induction of murine type I IFN genes can be divided into three phases. (a) The immediate early phase. Virus infection stimulates a phosphorylation cascade, leading to the activation of at least three families of transcription factors, including NF- κ B, AP-1 and IRF3. Activation of the IFN- α promoter requires all three transcription factors. (b) IRF7 induction phase. Secretion of early IFN produces an autocrine response through stimulation of the JAK-STAT pathway. Among the pathway's target genes is IRF7, itself. (c) Delayed early (amplification) phase. Many members of the IFN- α gene family possess promoter binding sites for activated IRF7 and become transcriptionally active.

RIG-1-like Receptors (RLRs) Sense Viral Products, Activate the IRF Pathway, and Trigger Production of Antiviral Proteins



Adapted from: Creagh and O'Neill, *Trends Immunol.* 27:352, 2006

Summary

1. Innate immunity is conserved throughout evolution and is triggered by recognition of "pathogen-associated molecular patterns" (e.g., LPS) by "pattern recognition receptors."
2. Collectins (e.g., SP-A, C1q, MBP) recognize carbohydrates on pathogen surfaces and perform multiple anti-microbial functions (e.g., opsonization). Collectins are essential for innate immunity, but also help clear apoptotic debris.
3. Members of the Scavenger Receptor superfamily recognize bacteria as well as glucose-modified proteins and oxidized lipoproteins. They are implicated in the response to infection as well as atherosclerosis and other degenerative diseases.
4. TLR4 is the major LPS receptor in mammalian cells. TLR4 triggers activation of NF- κ B (leading to production of TNF- α , for example). Other TLRs recognize additional microbial products. NOD-like receptors (NLRs) are intracellular sensors of bacterial products that activate the "inflammasome," triggering caspase-dependent maturation of IL-1.
5. Dendritic cells undergo a maturation program: immature DCs, which traffic to the periphery, capture antigen, and mature DCs, which traffic to the lymph node, present antigen. Innate immune stimuli trigger DC maturation, which upregulates co-stimulatory molecules and facilitates antigen presentation. Thus, the innate immune response ushers in the acquired immune response.
6. NK cells, a component of innate immunity, especially to viruses, represent an early source of IFN- γ and serve to stimulate macrophages and DCs in inflammatory sites. Additional components of the antiviral response include intracellular dsRNA sensors (RIG-like proteins) that activate the IRF pathway to signal antiviral gene expression.