


# The Patient with Visual Loss: Localization of Neuropathologic Disease and Select Diseases of Neuropathologic Interest


Steven A. Kane, M.D., Ph.D.  
The Edward S. Harkness Eye Institute

## Shared embryology



- Eye and brain develop from neuro-ectoderm
- Their functions and responses to disease are related
- Blood ocular/brain barriers
- The eye is a window into the brain and systemic disease

## Ocular anatomy




- Unique example of structure supporting function
- Optics
- Neuro-transduction
- Neuro-transmission

## Normal left ocular fundus




- Optic disc
- Retinal vessels
- Transparent retina
- Macula
- Retinal pigment epithelium
- Choroid

## Retinal nerve fiber layer anatomy



- Papillomacular bundle begins the macular-cortical projection
- Ganglion cells and axons respect the horizontal raphe

## Retro-bulbar visual anatomy




- Optic nerves carry information from each eye
- Axons from the nasal retinas cross at the optic chiasm
- Optic tracts carry right and left sided visual information
- Thalamus
- Optic radiations

## Localization and characterization of impaired vision

- Pattern of visual loss may identify the lesion site
- Disease course and accompanying symptoms may clarify its nature


## Patterns of visual loss



- Scotomas
- Central vision
- Peripheral vision
- Symmetry/congruity change as information nears cortex

# Ocular causes of impaired vision



- Refractive error
- Media opacity
- Retinal disease
- Optic nerve disease




# Retinoblastom

a

- Most common intraocular malignancy in childhood
- Leukocoria and strabismus
- 13 q14 mutation
- Spreads along the optic nerve into the brain



## Flexner-Wintersteiner rosettes



## Retinal causes of impaired vision







- Symptoms
- Age-related macular degeneration is the most common cause of visual loss > 65 years
- Diabetic retinopathy is the most common cause of visual loss < 65 years

## Symptoms and signs of optic nerve disease

- Blurred vision
- Dimming of vision with decreased color perception
- Decreased pupillary response to light
- Centrocecal, and arcuate scotomata

# Centrocecal scotomas





## Bilateral optic atrophy with centrocoecal scotoma

- Hereditary (dominant, Leber's)
- Toxic (medications, methanol, heavy metals)
- Nutritional (folate, B12)
- Demyelinating (optic neuritis, multiple sclerosis)

# Unilateral optic nerve disorders

- Ischemic (anterior ischemic optic neuropathy, retinal occlusive disease)
- Compressive (orbital, anterior fossa)
- Inflammatory (demyelinating, infectious, rheumatologic)



## AION

- Patients usually > 50
- Sudden, usually stable visual loss
- Altitudinal scotoma
- Optic atrophy in 4-6 wk
- Causes
  - Idiopathic (anatomic)
  - Giant cell arteritis




## Giant cell arteritis

- Senior citizens
- Subacute, granulomatous, stenosing arterial disease
- Headache, amaurosis fugax, arthralgia, myalgia, weight loss
- Brain, cardiac, eye, skin, muscle end artery damage




## Compressive optic neuropathy

- Progressive scotoma
- Initially normal disc
- Signs of atrophy
  - Decrease in color
  - Decrease in vessels
  - Decrease in NFL



# Inflammatory optic neuropathy


- Children and younger adults
- Centrocecal, arcuate, and hemianopic scotomas
- Subacute, often painful
- Retrobulbar neuritis or papillitis



# Papillitis and retrobulbar neuritis


Childhood

Adult






## Other causes of optic atrophy

- Glaucoma
- Secondary to retinal degeneration
- Central retinal artery obstruction
- Post-papilledema
- Congenital anomalies: hypoplasia, coloboma




## Glaucoma

- Common, usually bilateral, often asymmetric optic neuropathy
- Initial selective damage to branching axons





# Retinal degeneration

- Photoreceptor and/or retinal pigment epithelium disturbance
- Vascular narrowing is earliest sign
- Pigment released from damaged RPE cells clumps or migrates into the retina
- Many causes






# Central retinal artery obstruction



# Papilledema

- versus other disc swelling
- Intracranial mass
- Pseudotumor cerebri
- Hydrocephalus
- Intracranial hemorrhage
- Venous thrombosis
- Meningitis




## Papilledema in a 12 year old with idiopathic intracranial hypertension



## Other causes of disc swelling


- Optic neuritis
- Anterior ischemic optic neuropathy
- Central retinal vein occlusion
- Diabetic papillopathy
- Infiltrative disorders
- Hypertension
- Pseudopapilledema





## Lesions of the chiasm

- Usually compressive
- Pediatric
- Hypothalamic glioma
- Craniopharyngioma
- Adult
- Pituitary adenoma
- Meningioma
- Craniopharyngioma
- Aneurysm




# Retrochiasmal lesions

- Hemianopic scotoma
- Grossly incongruous field defects
- Small afferent defect
- Children: neoplasm > vascular > trauma
- Adults: vascular > neoplasm > trauma



# Retrogeniculate lesions

- Normal pupils, nerves unless perinatal
- Superior hemianopia: temporal lobe
- Inferior hemianopia: parietal lobe
- More posterior, more congruity




## Select Neuro-ophthalmic manifestations of systemic diseases




## Cherry red spots

- Tay Sachs & Sandoff's
- Niemann Pick type A
- Metachromatic leukodystrophy
- Sialidosis
- Farber disease



## Retinal pigmentary degeneration

- Mucopolysaccharidoses, Gaucher's, Refsum,
- Neuronal ceroid lipofuscinosis, cystinuria
- Abetalipoproteinemia, Kearns-Sayre
- Hallervorden Spatz, Spinocerebellar ataxias
- Usher, Cockayne




# Optic atrophy

- Krabbe, Metachromatic leukodystrophy
- Adrenoleukodystrophy, Alexander
- Spinocerebellar ataxia type I
- Friedreich's ataxia, Canavan's,
- Pelizaeus-Merzbacher, Alper's

## Ocular manifestations of diabetes


- Clinical background and pre-proliferative disease
- Proliferative disease
- Diabetic papillitis
- Neovascular glaucoma
- Cataract



# Ocular manifestations of hypertension


Narrowed arterioles

- Hypertensive retinopathy
- Hypertensive choroidopathy
- Hypertensive optic neuropathy





# Neurofibromatosis

- Dominant with complete penetrance and variable expressivity
- Skin, brain, eye, bone, visceral
- Ocular signs: Lisch nodules, optic nerve glioma, choroidal hamartoma




# Tuberous sclerosis

- Hamartomas: skin, kidney, eye, brain, heart
- Dominant and new mutations
- Symptoms: seizures, MR, facial angiofibromas, hydrocephalus
- Cortical hamartoma = tuber
- Retinal astrocytic hamartoma





# Sturge-Weber syndrome

- Port wine stain
- Glaucoma
- Leptomeningeal angioma and seizures




# von Hippel-Lindau disease

- Dominant
  - incomplete penetrance
  - variable expressivity
- VHL1 on 3p
- Retina, CNS, viscera: angiomas and capillary hemangioblastoma



# Wyburn-Mason syndrome

- AVMS
- Locations
  - Retina
  - Orbit
  - Nasopharynx
  - Midbrain



# Summary

- Visual loss can be understood when knowledge of neuropathophysiology is combined with knowledge of ocular embryology and anatomy
- The pattern of visual loss can localize and identify neuropathologic disease
- The number of systemic diseases having neuro-ophthalmic manifestation is legion