

Acute Respiratory Failure

Phil Factor, D.O.
Associate Professor of Medicine
Pulmonary, Allergy, and Critical Care Medicine
Director, Medical Intensive Care Unit
Columbia University Medical Center

Respiratory Failure

Physiologic Definition:

**Inability of the lungs to meet the
metabolic demands of the body**

Can't take in enough O_2
or
Can't eliminate CO_2 fast enough to keep up
with production

Respiratory Failure

- Failure of Oxygenation: $P_aO_2 < 60$ mmHg
- Failure of Ventilation*: $P_aCO_2 > 50$ mmHg

* P_aCO_2 is directly proportional to alveolar minute ventilation

Acute Respiratory Failure

Physiologic Classification

	Type 1 Hypoxemic	Type 2 Ventilatory	Type 3 Post-op	Type 4 Shock
Mechanism	Shunt	↓ V_a	Atelectasis	↓ Cardiac Output
Etiology	Airspace Flooding	Increased Respiratory load, Decreased ventilatory drive	Decreased FRC and increased Closing Volume	Decreased FRC and increased Closing Volume
Clinical Setting	Water, Blood or Pus filling alveoli	CNS depression, Bronchospasm, Stiff respiratory system, respiratory muscle failure	Abdominal surgery, poor insp effort, obesity	Sepsis, MI, acute hemorrhage

Ventilatory Failure




Inbalance between load on the lungs and the ability of bellows to compensate

Type 1 Respiratory Failure

Acute Hypoxemic Respiratory Failure

- Shunt disease - intracardiac or intrapulmonary
- Severe V/Q mismatch - asthma, PE
- Venous admixture due to low cardiac output states, severe anemia coupled with shunt and/or V/Q mismatch



Acute Respiratory Distress Syndrome (ARDS)

American-European Consensus Definition:*


- Refractory hypoxemia
 $P_aO_2/F_{I}O_2$ (P/F ratio)
<300 for ALI,
<200 for ARDS
- A disease process likely to be associated with ARDS
- No evidence of elevated left atrial pressure elevation (by clinical exam, echo or PA catheter)
- Bilateral airspace filling disease

Report of the American-European Consensus conference on acute respiratory distress syndrome: definitions, mechanisms, relevant outcomes, and clinical trial coordination. Consensus Committee.

Acute Respiratory Distress Syndrome

Each year in the U.S.:

75,000-150,000 cases




Causes of ARDS

DIRECT LUNG INJURY

- Pneumonia
- Aspiration of gastric contents
- Pulmonary contusion
- Near-drowning
- Inhalation injury (Cl⁻, smoke)
- Reperfusion pulmonary edema after lung transplantation or pulmonary embolectomy

INDIRECT LUNG INJURY

- Non-pulmonary sepsis/SIRS
- Severe trauma with shock
- Cardiopulmonary bypass
- Drug overdose (Narcotics)
- Acute pancreatitis
- Massive transfusion (TRALI)
- Drug reaction (ARA-C, nitrofurantoin)
- fat/air/amniotic fluid embolism, bypass







ARDS

Fundamental Pathophysiology:


Increased alveolar permeability due to direct neutrophil-mediated injury to the alveolar epithelium

Not a distinct disease - rather a sequelae of activation of lung and systemic inflammatory pathways





Shunt




Severe Hypoxemia

Therapeutic Goals

Maintain reasonable oxygen delivery

Find & fix the primary cause





"Baby Lungs"



FRC can be reduced by 80% or more in ARDS

Gattinoni, et. al. *Anesthesiology*, 74:15-23, 1991.


ARDS Network Trial

Day 1 Ventilatory Characteristics

	Low V_t Group n=432	Traditional V_t Group n=429
V_t :	6.2 ± 0.9	11.8 ± 0.8
PEEP:	9.4 ± 3.6	8.6 ± 3.6
F_{iO_2} :	0.56 ± 0.19	0.51 ± 0.17
P_{plat} :	25.7 ± 7	33 ± 9
P_{peak} :	32.8 ± 8	39 ± 10
P_aO_2 / F_{iO_2} :	158 ± 73	176 ± 76
P_aCO_2 :	40 ± 10	35 ± 8
pH:	7.38 ± 0.08	7.41 ± 0.07


NEJM 342:1301-1308, 2000

ARDS Network Trial: Oxygenation



Ventilation with low tidal was associated lower P/F ratios on days 1 & 3

ARDS Network Trial



Mortality: 39.8 % in traditional tidal volume group, 31% in low tidal volume group (P=0.007)

Also: @ 28 days: more ventilator free days (12 vs. 10), more days without organ failure (15 vs 12), higher rate of liberation from ventilation rate (65.7% vs 55%)

NEJM 342:1301-1308, 2000

Permissive Hypercapnea

Low tidal volume mechanical ventilation are usually associated with low alveolar minute ventilation which results in high P_aCO_2

- High P_aCO_2 is well tolerated*
- P_aCO_2 should be allowed to rise slowly
- Contraindications: increased ICP, active myocardial ischemia/severe RV or LV failure, severe metabolic acidosis
- Bicarb infusion may be used if $pH < 7.10$

High P_aCO_2 is preferable to flogging the lungs to "normalize gases"

*Mullu, Schwartz, Factor, Crit Care Med, 30:477-480, 2002
*Tuxen, Am J Resp Crit Care 150:870-90, 1994

Pulmonary Edema Accumulation

Starling's Equation


$$Q_E = K_F \left[(P_{mv} - P_{is}) - \sigma (\pi_{mv} - \pi_{is}) \right]$$

Hydrostatic Gradient pushing fluid out of the capillary

~~Oncotic Forces pulling fluid back into the capillary~~

Fluid movement across a semipermeable membrane is governed by opposing hydrostatic and oncotic pressure gradients

Pulmonary Edema Clearance Correlates with Outcome in Patients with Acute Lung Injury




Patients with intact active Na^+ transport had better survival (20% vs 62%) and shorter duration of mechanical ventilation

Ware, Am J Resp Crit Care Med. 163:1376-1383, 2001

Pulmonary Edema

Not just due to fluid
leaking into alveoli

It's also a problem
with fluid clearance




Reduce further edema accumulation

Reduce pulmonary vascular hydrostatic pressures to limit edema accumulation

"Keep-em-Dry"

*Diuresis, Phlebotomy,
Venodilators*

What happens to alveoli in ARDS?




What happens to alveoli in ARDS?

*Edema accumulates
in alveoli*


*Diluting & disaggregating
surfactant*

Surface tension increases

Alveoli collapse



Alveolar collapse decreases FRC and contributes to hypoxemia



Positive End-Expiratory Pressure (PEEP)

- Beneficial Effects
 - Increases FRC, CI, P_aO_2
 - Recruits Atelectatic Units
 - Decreases Q_s/Q_t
 - Allows Reduction in F_iO_2
- Detrimental Effects
 - Barotrauma - Volutrauma
 - Alveolar Overdistention
 - Hemodynamic Derangements

PEEP

Oxygen is:

- A) good for you
- B) bad for you
- C) all of the above

$F_iO_2 > 0.6$ for 24 hours or more may cause lung injury

PEEP recruits collapsed alveoli,
improves FRC and
improves oxygenation

An essential therapy for patients with ARDS

ARDS Network Trial

Assist Control
 V_T 6 cc/kg ideal body weight
PEEP of \approx 8-10

Generally not due to
respiratory failure

Does Mechanical Ventilation Contribute to MSOF?


Ranieri, et al.*: randomized prospective study of the effects of mechanical ventilation on bronchoalveolar lavage fluid and plasma cytokines in patients with ARDS (primarily non-pulmonary causes).

Controls (n=19): Rate 10-15 bpm, V_T targeted to maintain $PaCO_2$ 35-40 mmHg (mean: 11 ml/kg), PEEP titrated to SaO_2 (mean: 6.5), P_{plat} maintained <35 cmH₂O

Lung protective ventilation (n=18): Rate 10-15 bpm, V_T targeted to keep P_{plat} less than upper inflexion point (mean: 7 ml/kg), PEEP 2-3 cmH₂O above LIP (mean: 14.8)

Plasma and BALF levels of IL-1 β , IL-6, IL-8, TNF α , TNF α -sr 55, TNF α -sr 75, IL-1ra, measured within 8 hrs of intubation and again @24-30 hours & 36-40 hours after entry

*Ranieri, et al. Effect of mechanical ventilation on inflammatory mediators in patients with acute respiratory distress syndrome: a randomized controlled trial. JAMA 282:54-61, 1999.



The lung is not just an innocent bystander - it functions as an immunomodulatory organ that may participate in the systemic inflammatory response that leads to multiple organ system dysfunction syndrome

Biotrauma

Goals for Management of ARDS

The American-European Consensus Conference on ARDS, Part 2

- Ensure appropriate O₂ delivery to vital organs
- Minimize oxygen toxicity/tolerate mediocre ABG's
- Reduce edema accumulation
- Minimize airway pressures
- Prevent atelectasis/Recruit alveoli
- Use sedation and paralysis judiciously

Am J Resp Crit Care Med 157:1332-47, 1998.

Survival from "pure" ARDS

1979: 20-50%

2002: 50-90%