


Pulmonary Function Tests



PFT Interpretation

The interpretation of lung function tests involves two tasks:

- 1) the classification of the derived values with respect to a reference population and assessment of the reliability of the data; and
- 2) the integration of the obtained values into the diagnosis, therapy and prognosis for an individual patient.

ATS/ERS TASK FORCE: STANDARDISATION OF LUNG FUNCTION TESTING" Eur Respir J 2005; 26: 153-161

60 yo with progressive SOB


- FVC: 1.39 L (37%)
- FEV1: 0.54 L (19%)
- FEV1/FVC: 39%
- VC: 1.82 L (49%)
- TLC (PL): 7.42 L (122%)
- VA (He): 2.34 L
- DLCO: 40% of predicted

Pulmonary Function: Tests

- "Dynamic function": obstructive defects
- "Static function": restrictive defects
- Diffusion abnormalities (gas exchange)


60 yo with progressive SOB

- Asthma
- Interstitial Pulmonary Fibrosis
- Emphysema
- Primary pulmonary hypertension
- Amyotrophic lateral sclerosis






Spirometry and Maximal Expiratory and Inspiratory Flow Volume Curves

- “Dynamic function”



Spirometry



Obstructive Ventilation:
Expiratory

- Decrease in expiratory airflow (volume and/or rate of flow)
- FEV₁ decreased
- FVC normal or decreased
- FEV₁/FVC decreased*
- FEF₂₅₋₇₅ decreased


*definition of obstructive defect

Types of Airflow Obstruction

- Bronchoconstriction
- Dynamic airway compression (FVC vs SVC).
Emphysema: FVC < slow or inspiratory VC, and plethysmographic volumes greater than gas dilution volumes
- Upper Airway
- Small Airways
- "Mixed"

PFT Question #1


- FEV₁/FVC=obstructive ventilatory defect:
- Why is FEV₁ itself NOT diagnostic of an obstructive defect?






Patient: [redacted]	Gender: Male	Location: Out-Pt	Date: [redacted]
Age: 65	Height(in): 70	Temp: 29	PBar: 7
Weight(lb): 204	(cm): 179	Physician: [redacted]	
	(kg): 92.5	Technician: GD	

Spirometry		Ref	Pre Meas	Pre % Ref	Post Meas	Post % Ref
FVC	Liters	4.70	1.93	41	2.71	58
FEV1	Liters	3.63	0.54	15	0.60	17
FEV1/FVC	%	77	28		22	
FEF25-75%	L/sec	2.88	0.25	9	0.24	8
FEF25%	L/sec	7.80	0.27	3	0.29	4
FEF50%	L/sec	4.32	0.18	4	0.19	4
FEF75%	L/sec	1.57	0.10	6	0.09	6
PEF	L/sec	8.44	2.27	27	2.96	35
MVV	L/min	134			26	19
PIF	L/sec	3.67				
FIF50%	L/sec	4.59				
FET100%	Sec		13.02		19.70	

Lung Volumes				
VC	Liters	4.49	2.85	63
TLC	Liters	6.59	8.66	132
RV	Liters	2.46	5.81	236
RV/TLC	%	39	67	
FRC PL	Liters	3.52	7.02	199
FRC He	Liters	3.52		
Vtg	Liters		6.94	



Upper Airway Obstruction



Gas Equilibration Lung Volumes

- “Wash in:” Helium (insoluble gas) breathed from a reservoir of known VOLUME and CONCENTRATION, thus diluting its concentration by the volume of the lungs
- $V_{FRC} = V_{reservoir} \times \frac{Conc_{INIT} - Conc_{FINAL}}{Conc_{FINAL}}$

Lung Volumes

- “Static function”
- Gas Equilibration (“wash in” and “wash out”)
- Body plethysmography

Gas Equilibration Lung Volumes

- “Wash out:” Lung gas (N₂) washed out during breathing of 100% O₂
- Initial N₂ concentration known (atmospheric); volume and N₂ concentration of expired gas measured
- $V_{FRC} = V_{EXP} \times \frac{Conc_{EXP}}{.79 - Conc_{ALV} (final)}$

Plethysmographic Lung Volumes

- $P_1V_1=P_2V_2$ in a closed system at same temperature
- Lungs and airway closed system when occluded
- Panting at FRC: inhalation=decreased intrathoracic pressure, increased volume

Measurement of Alveolar Volume (V_A) in Emphysema

$$V_{A \text{ pleth}} > V_{A \text{ He rebreath}} > V_{A \text{ He single breath}}$$

$$V_{A \text{ He rebreath}} > V_{A \text{ single breath}} \text{ correlated with decreased FEV}_1/\text{FVC, increased RV/TLC}$$

Plethysmographic Lung Volumes

So, in inspiration:

$$P_{\text{FRC}} \times V_{\text{FRC}} = (P_{\text{FRC}} - \Delta P)(V_{\text{FRC}} + \Delta V)$$

- $V_{\text{FRC}} = \Delta V / \Delta P (P_{\text{FRC}} - \Delta P)$ where ΔP is negligible c/w P_{FRC}
- $V_{\text{FRC}} = \Delta V / \Delta P (P_{\text{FRC}})$
- ΔP obtained from change in mouth pressure against occluded valve
- ΔV obtained from change in pressure in the plethysmograph as air in the box is compressed by increase in lung volume
- P_{FRC} = alveolar pressure = atmospheric pressure with zero flow against occluded airway

Restrictive Ventilation

- A decrease in lung expansion
- FEV_1 decreased
- FVC decreased
- FEV_1/FVC normal or increased
- Total Lung Capacity (TLC) decreased*

* Definition of restrictive ventilatory defect

PFT Question #2

- With airways disease/dysfunction (e.g., emphysema), if gas dilution is not complete, how will lung volume measurement be affected?


PFT Questions #3 and #4

- Why is FVC itself NOT diagnostic of a restrictive ventilatory defect?
- Why is VC itself not diagnostic of a restrictive ventilatory defect?

Types of Restrictive Defects

- Parenchymal removal/destruction
- Parenchymal infiltration
- Extrapulmonary deformity
- Reduced force generation

Diffusing Capacity (Transfer Factor)



Restrictive patterns

- Diffuse parenchymal disease, thoracic cage restriction: symmetric decrease in TLC, VC, FRC, RV
- Neuromuscular weakness: IC mainly decreased; TLC and VC decreased and FRC and RV spared

Diffusing Capacity for CO (DL_{CO})

- $DL_{CO} = CO \text{ rate of uptake (ml/min)} / \Delta PCO \text{ (mmHg)}$
- O₂ and CO combine with Hgb; therefore reflect properties of alveolar-capillary membrane, and its uptake therefore limited by resistance across this interface
- Soluble gases limited by pulmonary blood flow
- 2 major resistances therefore: membrane properties (*D_m*), and "reactive" conductance (molecular conformation/rate of reaction properties of Hgb binding x pulmonary capillary blood volume (*V_c*)).

New York Presbyterian Hospital
Columbia Presbyterian Medical Center
622 West 168th Street New York, NY 10032

Adult Pulmonary Diagnostic Unit

52427-E

Patient: ID: Date:
 Age: Gender: Female Location:
 Height: 65 in (164 cm) Weight: Physician:
 Body Mass Index: Technician:

	Ref	Pre	Pre	Post	Post	Post
		Meas	% Ref	Meas	% Ref	% Chg
Spirometry						
FVC	Liters	3.51	2.00	57		
FEV1	Liters	2.91	1.52	52		
FEV1/VC %		82	76	38		
FEF25-75%L/Sec		3.15	1.21	38		
FEF50%	L/Sec	5.79	3.65	97		
FEF75%	L/Sec	1.55	0.39	25		
PEF	L/Sec	6.25	4.32	69		
MVV	L/min	104				
PiE	L/Sec	4.31	4.31	100		
PIE50%	L/Sec	3.95	4.29	105		
FEF100%	L/Sec		0.50			
FEF100%	Sec		7.50			
Lung Volumes						
VC	Liters	3.51	1.87	53		
TLC	Liters	5.14	2.67	52		
RV	Liters	1.90	0.80	47		
RV/TLC	%	33	30			
FRC FL	Liters	2.85				
FRC H2	Liters	2.85	1.23	43		
FRC He	Liters	2.85				
Vg2	Liters					
Diffusion						
DLCO	mL/min/Hg/mm	27.3	13.8	51		
DL Adj	mL/min/Hg/mm	27.3	13.8	51		
VA	Liters	2.85	2.58			
DLCO/VA	mL/min/L	5.48	5.37	98		

Diffusing Capacity for CO (DL_{CO})

Determinants:
Gas gradient, solubility, hemoglobin and rate of gas uptake by Hgb, membrane thickness, surface area, capillary blood volume

Single breath Diffusing Capacity for CO (DL_{CO SB})

- Inspirate 0.25% CO, 10% inert gas, 21%O₂, balance N₂
- Expire to RV; inhale rapidly to TLC; hold for remainder of 10 seconds of breath hold time
- Expire; discard anatomic dead space gas; sample 500-1000 ml alveolar gas

“Diffusion Capacity” vs Diffusion

- Note that: decreased diffusing capacity/gas transfer abnormality can result from numerous abnormalities not having anything to do with diffusion block itself
- So when we say diffusion abnormality=cause of hypoxemia, we mean those abnormalities which involve some form of diffusion block, or other inability to transfer gas completely (eg, low PIO₂+ increased circulatory time) so that insufficient transfer of alveolar PO₂ occur
- Low alveolar volume, low Hgb, may result in low diffusing capacity as measured by transfer of CO, and low O₂ content, but not low PaO₂

Measured Diffusing Capacity

- Increased in alveolar hemorrhage, erythrocytosis, obesity, asthma??, altitude?? (since CO and O₂ in competition, altitude decreases PIO₂ and increases DLCO; but less PO₂ gradient as well), supine, L-R shunt
- Decreased in emphysema (destruction and/or non-equilibration), ? restrictive disorders (all:why??), pulmonary vascular disorders, anemia, abnormal Hgb

New York Presbyterian Hospital
Columbia Presbyterian Medical Center
622 West 168th Street New York, NY 10032

Adult Pulmonary Diagnostic Unit ID: 52427-E Date:

Patient: Age: Gender: Female Location: Height: 65 in (164 cm) Weight: Body Mass Index: Physician: Technician: Post Meas % Ref Post % Chg

Spirometry	Ref	Pre Meas	Pre % Ref	Post Meas	Post % Ref	Post % Chg
FVC	Liters	3.51	2.00	57		
FEV1	Liters	2.91	1.52	52		
FEV1/FVC	%	82	76			
FEF25-75%L/Sec		3.18	1.21	38		
FEF25%	L/Sec	5.79	3.85	67		
FEF50%	L/Sec	3.77	2.14	57		
FEF75%	L/Sec	1.55	0.39	25		
PEF	L/Sec	6.25	4.32	69		
MVV	L/min	104				
PIF	L/Sec	4.31	4.31	100		
FEF50%	L/Sec	3.65	4.29	103		
FEF75%	L/Sec	0.50				
FET100%	Sec	7.50				
Lung Volumes						
VC	Liters	3.51	1.87	53		
TLC	Liters	5.14	2.67	52		
RV	Liters	1.60	0.80	47		
RV/TLC	%	33	30			
FRC PL	Liters	2.85				
FRC N2	Liters	2.85	1.23	43		
FRC Hc	Liters	2.85				
Vg	Liters					
Diffusion						
DLCO	ml/min/Hg/min	27.3	13.8	51		
DL Adj	ml/min/Hg/min	27.3	13.8	51		
VA	Liters		2.55			
DLCO/VA	ml/min/Hg/ml	5.48	5.37	96		

Diffusing Capacity

- Single breath (10 sec) vs steady state/rebreathe techniques: SB may UNDERESTIMATE true diffusing capacity in emphysema if it underestimates gas dilution V_A since DLCO = (kcO_xVA)/(P_b-PH₂O)

New York Presbyterian Hospital
Columbia Presbyterian Medical Center
622 West 168th Street New York, NY 10032

Adult Pulmonary Diagnostic Unit ID: Physician: Technician: Date: PBar:

Patient: Age: Gender: Male Location: Height: 67 in (168 cm) Weight: Body Mass Index: Physician: Technician: Post Meas % Ref Post % Chg

Spirometry	Ref	Pre Meas	Pre % Ref	Post Meas	Post % Ref	Post % Chg
FVC	Liters	4.06	2.73	67	2.52	72
FEV1	Liters	3.17	1.22	38	1.21	38
FEV1/FVC	%	78	45		41	
FEF25-75%L/Sec		3.03	0.32	10	0.27	9
FEF25%	L/Sec	7.41	1.27	17	1.06	14
FEF50%	L/Sec	3.37	0.44	12	0.36	11
FEF75%	L/Sec	1.22	0.11	9	0.09	7
PEF	L/Sec	7.85	4.52	66	4.11	54
MVV	L/min	121			52	43
PIF	L/Sec	3.54	3.64	103	4.35	123
FEF50%	L/Sec	4.54	3.34	74	3.93	87
FET100%	Sec		15.58		19.81	27
Lung Volumes						
VC	Liters	4.06	3.10	76		
TLC	Liters	6.32	5.22	83		
RV	Liters	2.20	3.12	142		
RV/TLC	%	35	50			
FRC PL	Liters	3.29	3.72	113		
FRC N2	Liters	3.29				
FRC Hc	Liters	3.29				
Vg	Liters			3.68		
Diffusion						
DLCO	ml/min/Hg/min	26.3		11.5	39	
DL Adj	ml/min/Hg/min	26.3		11.5	39	
VA	Liters			3.75		
DLCO/VA	ml/min/Hg/ml	4.80		3.06	64	

DLCO Pearl

- Isolated DLCO decrease (normal spirometry and volumes): suspect pulmonary vascular disorder
- Or, interstitial disorder not yet, or no longer, affecting parenchymal volume
- Or, abnormality of Hgb (eg, anemia, carboxyHgb, methHgb)

PFT Summary

- Obstructive ventilatory defect: decreased FEV₁/FVC
- Restrictive ventilatory defect: decreased TLC
- Low DLCO: abnormal uptake of gas by Hgb across alveolar capillary membrane: Diffusion determinants= Gas gradient, solubility, hemoglobin, membrane thickness, surface area, alveolar volume, rate of circulatory flow
- Disorders with airway dysequilibrium (eg emphysema): single breath gas dilution will underestimate lung volumes (and ? DLCO)

Pre-operative Pulmonary Assessment: PFTs


- Complications: highest for thoracic and upper abdominal (ie, near the diaphragm)
- All having lung resection, orthopaedic and lower abdominal with lung disease, or smoking
- Age > 60 years

Series "ATS/ERS TASK FORCE: STANDARDISATION OF LUNG FUNCTION TESTING" Edited by V. Brusasco, R. Crapo and G. Viegi.
General considerations for lung function testing

Eur Respir J 2005; 26: 153–161

Postoperative Pulmonary Risks

- Spirometry: FEV₁ or FVC < 70%, FEV₁/FVC < 65%
- PaCO₂ > 45 mmHg, DLCO < 40% in COPD
- None contraindicate
- Lung resection: FEV₁ best for pulmonary reserve and post op complications; post op FEV₁ < 30% predicted = increased long term mortality and immediate post op problems



Diffusing Capacity for CO (DL_{CO}):

- DL_{CO} (if "transfer factor", TLCO) calculated as the product of the rate constant for CO uptake (also known as "permeability factor" and called **kCO, the Krogh coefficient**) and alveolar volume, divided by effective gas pressure (**PB-PH20**), expressed as units of conductance (eg, ml CO/min/mmHg);
- Thus, $DLCO = (kCO \times V_A) / (P_b - P_{H_2O})$.
- This assumes what the conductance would be if 100% of alveolar volume was filled with CO (that is the V_A component is the volume of distribution)
- Influenced by altitude since CO and O_2 in competition for Hgb; thus high altitude decreases PIO_2 and therefore increases DLCO (on the other hand, lower PO_2 gradient for transfer of oxygen)
- **Note that properties of CO uptake are different than those of oxygen (in fact, CO=diffusion limited, O_2 more perfusion limited)**