

Disjoint Set Code

Make-Set(x)

- 1 $p[x] \leftarrow x$
- 2 $rank[x] \leftarrow 0$

Union(x, y)

- 1 LINK(FIND-SET(x), FIND-SET(y))

Link(x, y)

- 1 if $rank[x] > rank[y]$
- 2 then $p[y] \leftarrow x$
- 3 else $p[x] \leftarrow y$
- 4 if $rank[x] = rank[y]$
- 5 then $rank[y] \leftarrow rank[y] + 1$

Find-Set(x)

- 1 if $x \neq p[x]$
- 2 then $p[x] \leftarrow \text{FIND-SET}(p[x])$
- 3 return $p[x]$

Ackerman's Function

$$A_k(j) = \begin{cases} j + 1 & \text{if } k = 0 \\ A_{k-1}^{(j+1)}(j) & \text{if } k \geq 1 \end{cases}$$

$$\alpha(n) = \min\{k : A_k(1) \geq n\}$$

$$\begin{aligned} A_0(j) &= j + 1 \\ A_1(j) &= A_0^{(j+1)}(j) \\ &= 2j + 1 \\ A_2(j) &= A_1^{(j+1)}(j) \\ &= 2(2(\cdots(2j + 1)\cdots) + 1) + 1 \\ &= 2^{j+1}(j + 1) - 1 \end{aligned}$$

Ackerman

$$\begin{aligned} A_3(1) &= A_2^{(2)}(1) \\ &= A_2(A_2(1)) \\ &= A_2(7) \\ &= 2^8 \cdot 8 - 1 \\ &= 2^{11} - 1 \\ &= 2047 \end{aligned}$$

$$\begin{aligned} A_4(1) &= A_3^{(2)}(1) \\ &= A_3(A_3(1)) \\ &= A_3(2047) \\ &= A_2^{(2048)}(2047) \\ &\gg A_2(2047) \\ &= 2^{2048} \cdot 2048 - 1 \\ &> 2^{2048} \\ &= (2^4)^{512} \\ &= 16^{512} \\ &\gg 10^{80}, \end{aligned}$$