

A comprehensive molecular phylogeny of the starlings (Aves: Sturnidae) and mockingbirds (Aves: Mimidae): Congruent mtDNA and nuclear trees for a cosmopolitan avian radiation

Irby J. Lovette^{a,*}, Dustin R. Rubenstein^{a,b,1}

^a Fuller Evolutionary Biology Program, Laboratory of Ornithology, Cornell University, Ithaca, NY 14950, USA

^b Department of Neurobiology and Behavior, Cornell University, Ithaca, NY 14850, USA

Received 27 September 2006; revised 15 March 2007; accepted 19 March 2007

Available online 5 April 2007

Abstract

We generated a comprehensive phylogeny for the avian families Sturnidae (starlings, mynas, *Rhabdornis*, oxpeckers, and allies) and Mimidae (mockingbirds, thrashers, and allies) to explore patterns of morphological and behavioral diversification. Reconstructions were based on mitochondrial DNA sequences from five coding genes (4108 bp), and nuclear intron sequences from four loci (2974 bp), for most taxa, supplemented with NDII gene sequences (1041 bp) derived from museum skin specimens from additional taxa; together the 117 sampled taxa comprise 78% of the 151 species in these families and include representatives of all currently or recently recognized genera. Phylogenetic analyses consistently identified nine major clades. The basal lineage is comprised of the two *Buphagus* oxpeckers, which are presently confined to Africa where they are obligately associated with large mammals. Some species in nearly all of the other major clades also feed on or around large vertebrates, and this association may be an ancestral trait that fostered the world-wide dispersal of this group. The remaining taxa divide into sister clades representing the New-World Mimidae and Old-World Sturnidae. The Mimidae are divided into two subclades, a group of Central American and West Indian catbirds and thrashers, and a pan-American clade of mockingbirds and thrashers. The Sturnidae are subdivided into six clades. The Phillipine endemic *Rhabdornis* are the sister lineage to a larger and substantially more recent radiation of South Asian and Pacific island starlings and mynas. A clade of largely migratory or nomadic Eurasian starlings (within which the basal lineage is the model taxon *Sturnus vulgaris*) is allied to three groups of largely African species. These reconstructions confirm that *Buphagus* should not be included in the Sturnidae, and identify many genera that are not monophyletic. They also highlight the substantial diversity among the major Sturnidae subclades in rates of species accumulation, morphological differentiation, and behavioral variation.

© 2007 Elsevier Inc. All rights reserved.

Keywords: Sturnidae; Mimidae; Buphagidae; Phylogeny; Biogeography

1. Introduction

The Mimidae and Sturnidae are avian sister radiations that show striking parallels in a number of ecological and behavioral traits. The Sturnidae (Starlings and Mynas) are restricted to the Old World (except for human-medi-

ated introductions elsewhere), with centers of diversity in Southeast Asia and Africa. The Mimidae have diversified in southwestern North America, the West Indies, and Central/South America and its satellite islands. Although taxa in both groups continue to serve as models in studies of behavioral (e.g., Derrickson, 1988; Kroodsma and Byers, 1991; Pinxten et al., 2002; Duffy and Ball, 2002; Gentner and Margoliash, 2003; Polo et al., 2004; Rubenstein, 2007a) and life history trait evolution (e.g., Ricklefs and Williams, 1984; Cordero et al., 2001; Christians et al., 2001; Komdeur et al., 2002. Rubenstein, 2007b), neither

* Corresponding author. Fax: +1 607 254 2486.

E-mail address: IJL2@cornell.edu (I.J. Lovette).

¹ Present address: Department of Integrative Biology and Museum of Vertebrate Zoology, University of California, Berkeley, CA 94720, USA.

group has previously been the subject of phylogenetic analysis with robust taxonomic sampling. Here, we use a combination of mtDNA and nuclear DNA sequences to explore the phylogenetic relationships of all genera and most species within this cosmopolitan avian radiation.

Despite earlier evidence from studies of jaw musculature and cranial osteology (Beecher, 1953), and seriology (Stallcup, 1961), the sister relationship of the Sturnidae and Mimidae was not broadly recognized until it was featured in the DNA–DNA hybridization studies of Sibley and Ahlquist (1980, 1990). Although this finding initially met with controversy, subsequent phylogenetic studies based on DNA–DNA hybridization (Sheldon and Gill, 1996), physiological traits (Malcarney et al., 1994), and various mitochondrial and nuclear DNA sequence loci (Voelker and Spellman, 2004; Ericson and Johansson, 2003; Cibois and Cracraft, 2004; Barker et al., 2004; Zuccon et al., 2006) have been completely concordant in grouping the Mimidae and Sturnidae as sister clades, usually with very strong topological support.

The reliable characterization of the full set of taxa that fall within a monophyletic Sturnidae/Mimidae group has been strengthened by recent phylogenetic surveys of related passerine songbird groups, particularly an intensively sampled study (Cibois and Cracraft, 2004) of the deeper Muscicapoida radiation within which the Sturnidae and Mimidae are nested. Cibois and Cracraft (2004) included many taxa that had not been sampled previously in any molecular phylogenetic analysis, and thereby helped confirm that all major lineages within the Sturnidae/Mimidae clade have been assigned correctly to this group. Their most surprising finding involving the Sturnidae/Mimidae was the recognition that *Rhabdornis*, a genus endemic to the Philippines with previously uncertain family-level affinities, is a morphologically aberrant member of the Sturnidae. A phylogenetic enigma involving a second morphologically unusual genus, the *Buphagus* oxpeckers of Africa, remains somewhat less well resolved. *Buphagus* has been variously treated as its own family (Buphagidae), or more commonly included within the family Sturnidae. All molecular phylogenies that have included the *Buphagus* lineage have placed it as a long branch at the base of the Sturnidae/Mimidae clade (e.g., Cibois and Cracraft, 2004; Zuccon et al., 2006), but with low support for distinguishing whether it is the basal lineage in this entire group, or alternatively the sister lineage to either the Sturnidae or Mimidae.

Less is known about relationships within and among the major subclades of the Sturnidae/Mimidae radiation. Although most of the species, as well as all of the genera, of Mimidae have been included in previous DNA-based phylogeographic or phylogenetic studies (Sibley and Ahlquist, 1990; Zink et al., 1997, 1999, 2001; Zink and Blackwell-Rago, 2000; Hunt et al., 2001; Sgariglia and Burns, 2003; Barber et al., 2004; Cibois and Cracraft, 2004; Arbogast et al., 2006), these previous reconstructions have each primarily addressed relationships among sets of closely

allied species and no single reconstruction has included a complete sample of Mimidae genera. The few previous DNA-based studies of relationships within the Sturnidae have similarly been taxonomically circumscribed. For example, the most inclusive survey of Sturnidae (Zuccon et al., 2006) sampled only 30 (of 117) Sturnidae species along with 6 (of 34) Mimidae species, and did not include many genus-level lineages with long-debated affinities.

Here, we use a combination of mitochondrial (mtDNA) and nuclear DNA sequences to reconstruct the phylogenetic relationships of all well-differentiated lineages within the Sturnidae/Mimidae. By using both modern, high-quality blood and tissue samples and skin-snips taken from dried museum specimens, we included 117 of 151 (78%) taxa representing all extant genera recognized by any recent taxonomic revision, and multiple species from most polytypic genera. From the high-quality samples, we obtained substantial mitochondrial DNA (4108 bp of protein-coding gene sequence) and nuclear intron (4 loci, 2974 aligned bp) sequences, to which we added shorter (NDII only, 1041 bp) sequences from samples derived from museum skin source materials. The majority of nodes in the resulting phylogenetic reconstructions have high topological support and provide strong evidence for the historical pattern of diversification in this world-wide avian radiation.

2. Materials and methods

2.1. Taxon sampling

We designed our taxonomic sampling strategy to include at least one representative of all morphologically or biogeographically distinctive lineages in the Sturnidae and Mimidae, including representatives of all genera recognized by any of the five most influential taxonomic treatments of the Sturnidae of the past half-century (Table 1). Here we employ the nomenclature of the most recent “Howard and Moore” checklist (Dickinson, 2003), which recognizes 26 genera and 114 extant species of Sturnidae inclusive of the two species in the “Rhabdornithidae,” which Cibois and Cracraft (2004) showed to fall well within the Sturnidae, and not including the Mascarene starling *Necropsar leguati*—a monotypic genus first described by Forbes (1898)—that recent investigations have shown to be based on fraudulently labeled specimens correctly assignable to the Mimidae genus *Cinlocerthia* (Olson et al., 2005). Three additional known species in the large genus *Aplonis* and the monotypic *Fregilupus* are extinct. There has been substantial recent volatility in the genus- and species-level taxonomy of the Sturnidae. Table 1 compares the Sturnidae genera and species of Dickinson (2003) against those recognized by Amadon (1962), Wolters (1982), Sibley and Monroe (1990), and Feare and Craig (1999). In total these authors have recognized 43 genera of extant Sturnidae (Table 1). We sampled 91 species (80% of the Sturnidae)

Table 1

Genera and species of extant Sturnidae recognized in five recent taxonomic treatments, and those included in this study

Dickinson (2003)		In this study	Amadon (1962)	Wolters (1982)	Sibley and Monroe (1990)	Feare and Craig (1999)
Genus	Species					
<i>Rhabdornis</i> ^a	<i>mysticalis</i>	×	—	—	—	NC
<i>Rhabdornis</i> ^a	<i>inornatus</i>	×	—	—	—	NC
<i>Aplonis</i>	<i>metallica</i>	×	—	<i>Lamprocorax</i>	—	—
<i>Aplonis</i>	<i>mystacea</i>	—	—	<i>Rhinopsar</i>	—	—
<i>Aplonis</i>	<i>cantoroides</i>	×	—	—	—	—
<i>Aplonis</i>	<i>crassa</i>	—	—	—	—	—
<i>Aplonis</i>	<i>feandensis</i>	—	—	—	—	—
<i>Aplonis</i>	<i>insularis</i>	×	—	—	—	—
<i>Aplonis</i>	<i>magna</i>	—	—	<i>Lamprocorax</i>	—	—
<i>Aplonis</i>	<i>brunneicapillus</i>	×	—	<i>Rhinopsar</i>	—	—
<i>Aplonis</i>	<i>grandis</i>	×	—	<i>Lamprocorax</i>	—	—
<i>Aplonis</i>	<i>dichroa</i>	—	—	<i>Lamprocorax</i>	—	—
<i>Aplonis</i>	<i>zelandica</i>	—	—	—	—	—
<i>Aplonis</i>	<i>striata</i>	—	—	—	—	—
<i>Aplonis</i>	<i>atronitens</i> ^b	—	—	ssp <i>striata</i>	ssp <i>striata</i>	ssp <i>striata</i>
<i>Aplonis</i>	<i>santovestris</i>	—	—	—	—	—
<i>Aplonis</i>	<i>panayensis</i>	×	—	<i>Lamprocorax</i>	—	—
<i>Aplonis</i>	<i>mysolensis</i>	—	—	<i>Lamprocorax</i>	—	—
<i>Aplonis</i>	<i>minor</i>	×	—	<i>Lamprocorax</i>	—	—
<i>Aplonis</i>	<i>opaca</i>	—	—	—	—	—
<i>Aplonis</i>	<i>pelzelni</i>	×	—	—	—	—
<i>Aplonis</i>	<i>tabuensis</i>	×	—	—	—	—
<i>Aplonis</i>	<i>atrifusca</i>	—	—	—	—	—
<i>Aplonis</i>	<i>cinerascens</i>	×	—	—	—	—
<i>Mino</i>	<i>dumontii</i>	—	—	—	—	—
<i>Mino</i>	<i>krefftii</i>	×	ssp <i>dumontii</i>	ssp <i>dumontii</i>	ssp <i>dumontii</i>	—
<i>Mino</i>	<i>anaïs</i>	×	—	—	—	—
<i>Basilornis</i>	<i>celebensis</i>	×	—	—	—	—
<i>Basilornis</i>	<i>galeatus</i>	—	—	—	—	—
<i>Basilornis</i>	<i>corythaix</i>	—	—	—	—	—
<i>Basilornis</i>	<i>miranda</i>	×	—	<i>Goodfellowia</i>	—	—
<i>Sarcops</i>	<i>calvus</i>	×	—	—	—	—
<i>Streptocitta</i>	<i>albicollis</i>	×	—	—	—	—
<i>Streptocitta</i>	<i>albertinae</i>	—	—	—	—	—
<i>Enodes</i>	<i>erythrophris</i>	×	—	—	—	—
<i>Scissirostrum</i>	<i>dubium</i>	×	—	—	—	—
<i>Saroglossa</i>	<i>spiloptera</i>	×	—	—	—	—
<i>Saroglossa</i>	<i>aurata</i>	×	—	<i>Hartlaubius</i>	—	—
<i>Ampeliceps</i>	<i>coronatus</i>	×	—	—	—	—
<i>Gracula</i>	<i>ptilogenys</i>	×	—	—	—	—
<i>Gracula</i>	<i>religiosa</i>	×	—	—	—	—
<i>Gracula</i>	<i>indica</i> ^b	×	ssp <i>religiosa</i>	—	ssp <i>religiosa</i>	—
<i>Gracula</i>	<i>robusta</i> ^b	—	ssp <i>religiosa</i>	ssp <i>religiosa</i>	ssp <i>religiosa</i>	—
<i>Gracula</i>	<i>enganensis</i> ^b	—	ssp <i>religiosa</i>	ssp <i>religiosa</i>	ssp <i>religiosa</i>	—
<i>Acridotheres</i>	<i>grandis</i>	×	—	<i>Aethiopsar</i>	—	—
<i>Acridotheres</i>	<i>crisatellus</i>	×	—	<i>Aethiopsar</i>	—	—
<i>Acridotheres</i>	<i>javanicus</i>	×	ssp <i>fuscus</i>	ssp <i>fuscus</i>	—	—
<i>Acridotheres</i>	<i>cinereus</i>	—	<i>Sturnus</i>	<i>Aethiopsar</i>	ssp <i>fuscus</i>	—
<i>Acridotheres</i>	<i>fuscus</i>	×	—	<i>Aethiopsar</i>	—	—
<i>Acridotheres</i>	<i>albocinctus</i>	—	—	<i>Aethiopsar</i>	—	—
<i>Acridotheres</i>	<i>ginginianus</i>	×	—	—	—	—
<i>Acridotheres</i>	<i>tristis</i>	×	—	—	—	—
<i>Acridotheres</i>	<i>melanopterus</i>	—	<i>Sturnus</i>	<i>Leucopsar</i>	<i>Sturnus</i>	—
<i>Leucopsar</i>	<i>rothschildi</i>	×	—	—	—	—
<i>Sturnus</i>	<i>burmannicus</i>	—	—	<i>Leucopsar</i>	—	<i>Acridotheres</i>
<i>Sturnus</i>	<i>nigricollis</i>	×	—	<i>Gracupica</i>	—	<i>Gracupica</i>
<i>Sturnus</i>	<i>contra</i>	×	—	<i>Sturnopastor</i>	—	<i>Gracupica</i>
<i>Sturnus</i>	<i>sturninus</i>	—	—	<i>Agropsar</i>	—	<i>Sturnia</i>
<i>Sturnus</i>	<i>philippensis</i>	×	—	<i>Agropsar</i>	—	<i>Sturnia</i>
<i>Sturnus</i>	<i>sinensis</i>	×	—	<i>Sturnia</i>	—	<i>Sturnia</i>
<i>Sturnus</i>	<i>malabaricus</i>	×	—	<i>Temenuchus</i>	—	<i>Sturnia</i>
<i>Sturnus</i>	<i>erithropygius</i>	—	—	<i>Temenuchus</i>	—	<i>Sturnia</i>

(continued on next page)

Table 1 (continued)

Dickinson (2003)		In this study	Amadon (1962)	Wolters (1982)	Sibley and Monroe (1990)	Feare and Craig (1999)
Genus	Species					
<i>Sturnus</i>	<i>albofrontatus</i>	×	—	<i>Temenuchus</i>	—	<i>Sturnia</i>
<i>Sturnus</i>	<i>pagodarum</i>	×	—	<i>Temenuchus</i>	—	<i>Temenuchus</i>
<i>Sturnus</i>	<i>roseus</i>	×	—	<i>Pastor</i>	—	<i>Pastor</i>
<i>Sturnus</i>	<i>sericeus</i>	×	—	<i>Sturnopastor</i>	—	—
<i>Sturnus</i>	<i>cineraceus</i>	×	—	<i>Sturnopastor</i>	—	—
<i>Sturnus</i>	<i>vulgaris</i>	×	—	—	—	—
<i>Sturnus</i>	<i>unicolor</i>	×	—	—	—	—
<i>Creatophora</i>	<i>cinerea</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>nitens</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>chalybaeus</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>chloropterus</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>elizabeth^b</i>	×	ssp <i>chlorop.</i>	—	—	ssp <i>chlorop.</i>
<i>Lamprotornis</i>	<i>chalcurus</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>splendidus</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>ornatus</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>iris</i>	×	—	<i>Coccycolius</i>	<i>Coccycolius</i>	—
<i>Lamprotornis</i>	<i>purpureus</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>purpuroptera</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>caudatus</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>regius</i>	×	<i>Cosmopsarus</i>	<i>Cosmopsarus</i>	<i>Cosmopsarus</i>	—
<i>Lamprotornis</i>	<i>mavesii</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>australis</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>acuticaudus</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>corruscus</i>	×	—	—	—	—
<i>Lamprotornis</i>	<i>superbus</i>	×	<i>Spreo</i>	<i>Lamprospreo</i>	—	—
<i>Lamprotornis</i>	<i>hildebrandti</i>	×	<i>Spreo</i>	<i>Lamprospreo</i>	—	—
<i>Lamprotornis</i>	<i>shelleyi</i>	×	ssp <i>hildebrandti</i>	<i>Lamprospreo</i>	—	—
<i>Lamprotornis</i>	<i>pulcher</i>	×	<i>Spreo</i>	<i>Lamprospreo</i>	—	—
<i>Lamprotornis</i>	<i>purpureiceps^c</i>	×	—	—	—	<i>Hylopsar</i>
<i>Lamprotornis</i>	<i>cupreocauda^c</i>	×	—	—	—	<i>Hylopsar</i>
<i>Lamprotornis</i>	<i>unicolor</i>	×	<i>Cosmopsarus</i>	<i>Cosmopsarus</i>	<i>Cosmopsarus</i>	<i>Spreo</i>
<i>Lamprotornis</i>	<i>fischeri</i>	×	—	<i>Lamprospreo</i>	<i>Spreo</i>	<i>Spreo</i>
<i>Cinnyricinclus</i>	<i>femoralis^d</i>	×	<i>Cinnyricinclus</i>	<i>Arizelopsar</i>	<i>Cinnyricinclus</i>	<i>Poeoptera</i>
<i>Cinnyricinclus</i>	<i>leucogaster</i>	×	—	—	—	—
<i>Spreo</i>	<i>bicolor</i>	×	—	—	—	—
<i>Spreo</i>	<i>albicapillus</i>	×	—	<i>Poneropsar</i>	—	—
<i>Onychognathus</i>	<i>morio</i>	×	—	—	—	—
<i>Onychognathus</i>	<i>tenuirostris</i>	×	—	—	—	—
<i>Onychognathus</i>	<i>fulgidus</i>	×	—	—	—	—
<i>Onychognathus</i>	<i>walleri</i>	×	—	—	—	—
<i>Onychognathus</i>	<i>blythii</i>	×	—	—	—	—
<i>Onychognathus</i>	<i>frater</i>	×	—	—	—	—
<i>Onychognathus</i>	<i>tristamii</i>	×	—	—	—	—
<i>Onychognathus</i>	<i>nabouroup</i>	×	—	—	—	—
<i>Onychognathus</i>	<i>salvadorii</i>	×	—	—	—	—
<i>Onychognathus</i>	<i>albirostris</i>	×	—	—	—	—
<i>Onychognathus</i>	<i>neumanni</i>	×	ssp <i>morio</i>	ssp <i>morio</i>	ssp <i>morio</i>	—
<i>Poeoptera</i>	<i>stuhlmanni</i>	×	—	<i>Onychognathus</i>	—	—
<i>Poeoptera</i>	<i>kenricki</i>	×	—	<i>Onychognathus</i>	—	—
<i>Poeoptera</i>	<i>lugubris</i>	×	—	<i>Onychognathus</i>	—	—
<i>Pholia</i>	<i>sharpii</i>	×	<i>Cinnyricinclus</i>	—	<i>Cinnyricinclus</i>	<i>Pholia</i>
<i>Grafisia</i>	<i>torquata</i>	×	—	—	—	—
<i>Speculipastor</i>	<i>bicolor</i>	×	—	—	—	—
<i>Neocichla</i>	<i>gutturalis</i>	×	—	—	—	—
<i>Buphagus</i>	<i>erythrorhynchus^e</i>	×	—	—	—	—
<i>Buphagus</i>	<i>africanus^e</i>	×	—	—	—	—

— indicates congruence with “Howard and Moore checklist” taxonomy of Dickinson (2003); ssp: treated as a subspecies of the species named thereafter; NC: not considered.

^a *Rhabdornis* was only recently recognized as a member of the Sturnidae (Cibois and Cracraft, 2004).

^b Taxa ranked as subspecies in Dickinson (2003) but as full species in one or more of the previous treatments.

^c Also assigned to *Hylopsar* in some additional regional treatments (e.g., Craig, 1997; Fry et al., 2000).

^d Also assigned to *Pholia* in some additional regional treatments (e.g., Craig, 1997; Fry et al., 2000).

^e Sometimes treated as a separate family, Buphagidae.

representing all 43 genera. All Sturnidae DNA sequences analyzed here were generated in our laboratory.

Dickinson (2003) recognizes 12 genera and 34 species of Mimidae, all of which are extant (with the possible exception of the highly endangered Cozumel Thrasher, *Toxostoma guttatum*). Although the genus-level taxonomy of the Mimidae has been fairly stable over the past half-century (Davis and Miller, 1960; Sibley and Monroe, 1990; AOU, 1998; Brewer, 2001; Cody, 2005), recent molecular evidence (Hunt et al., 2001; Banks et al., 2002) supported the re-separation of *Allenia* from *Margarops*, two Caribbean genera that had been merged in some earlier treatments. *Donacobius*, a monotypic genus of long-debated affinities, has sometimes been assigned to the Mimidae in the past, but molecular phylogenetic studies have now shown it to be a member of the Sylviidae and not closely related to the Mimidae or Sturnidae (Barker, 2004; Alström et al., 2006), and we did not include it here. We obtained sequences from 25 species (74% of the Mimidae) representing all recently recognized genera. Some or all DNA sequences from 16 Mimidae taxa (Appendix A) were derived from previous studies (Hunt et al., 2001; Barber et al., 2004; Arbogast et al., 2006).

We employed three outgroup taxa based on recent phylogenetic surveys that have addressed relationships between the Sturnidae/Mimidae clade and related passerine groups. These reconstructions show that the waxwing *Bombycilla* is the most basal outgroup lineage in our study, and that the thrushes *Catharus* and *Myadestes* represent the two early and well differentiated lineages (Klicka et al., 2005) within a large clade that is either sister to the Sturnidae/Mimidae (Barker et al., 2004; Cibois and Cracraft, 2004; Zuccon et al., 2006), or sister to a clade that contains the dippers (Cinclidae) and the Sturnidae/Mimidae (Ericson and Johansson, 2003; Voelker and Spellman, 2004).

The majority of samples that served as sources of DNA for our study were frozen tissues associated with traditionally vouchered specimens, either from existing museum collections or from our own collecting activities (Appendix A). We added additional taxa based on DNAs extracted from the toe-pads of traditionally prepared museum skin specimens. From some taxa, high-quality tissue samples were available only from non-vouchered tissue, blood, or feather samples provided by field researchers or taken from live captive birds in zoological or private avicultural collections. In these cases where otherwise high-quality DNA materials were not associated with voucher specimens, we sequenced the NDII gene (see below) from conspecific museum-skin specimens. This allowed us to generate much greater amounts of sequence from the robust but non-vouchered samples, while confirming their phylogenetic affinities by comparison to the NDII sequences from conspecific vouchered samples. We likewise replicated nearly all sequences derived from skin-snips, as well as a number of samples from vouchered frozen tissues, using samples from different conspecific individuals. In all but two cases, these replicated NDII sequences were identical or nearly identi-

cal matches to their conspecific counterparts; investigations of the two mismatches (one each from two different museum-based frozen tissue collections) showed them to be vouchered specimens misidentified by the field preparators, and consequently accessioned incorrectly. To reduce computation times we included only one representative per species in the reconstructions reported here; however, the replicated sequences used only in preliminary analyses are also archived in GenBank (Appendix A).

2.2. Laboratory methods

DNA was extracted from muscle tissue and feather samples using DNAeasy kits (Qiagen) and from blood samples using Perfect gDNA Blood Mini kits (Eppendorf). To amplify the mitochondrial NDII gene, we used primers METb and TRPc (Eberhard and Bermingham, 2004). To amplify the mitochondrial region spanning the COI and ATPase6 genes, we used various combinations of primers COIf and COIa (Kessing et al., 1989); GQL and HMH (Hunt et al., 2001); IL6591L (Lovette, 2004); and IL7513h (ATGGATAGCATGGCTCATACTATTCC), sturnCOIf2 (GACACCTACTACGTWGTAGCYCACTTCC), sturnCOIa2 (GGAAACCGARTTGTGAGTGGT TGG), IL8232I (ATGTTGGTTTCAAGCCAACCGC), and ILLYSh (CCTCTTTCTCCAGCTTAAAAGGCT AG). Amplification of β -fibrinogen introns 5 and 7 used primer pairs Fib-5 (CGCCATACAGAGTATACTGTGACA) and Fib-6 (GCCATCCTGGCGATTCTGAA) provided by F.K. Barker (personal communication), and FIB-B17U and FIB-B17L (Prychitko and Moore, 1997), respectively. To amplify rhodopsin intron 1, we used Rho-IIF and Rho-IIR (Primmer et al., 2002). To amplify intron 5 of transforming growth factor β -2, we used primers TGFB2-I5F and TGFB2-I5R (Primmer et al., 2002).

All 10 μ L PCR amplifications included 1 μ L of undiluted genomic DNA (concentration: 10–50 ng/ μ L), 10 μ M Tris-HCl (pH 8), 50 μ M KCl, variable $MgCl_2$ (range: 1.5–4 mM), 0.25 mM of each nucleotide, 0.25 mM of each primer, and 0.025 U Jumpstart Taq polymerase (Sigma). All sets of PCR reactions included negative controls with no DNA template. Thermal cycling and subsequent cycle sequencing was conducted in PTC-220 Dyad Thermal Cyclers (MJ Research). Thermal cycling profiles varied among primer sets, but most commonly employed an initial denaturing at 95 °C for 4 min 30 s; 30–35 cycles of denaturing at 95 °C for 45 s, annealing at a variable temperature for 45 s, and extension at 72 °C for 1–2 min 20 s; and a final extension at 72 °C for 5 min.

PCR products were electrophoresed in 1.5% agarose TAE gels to confirm amplification and fragment sizes. To digest unincorporated nucleotides and primers, 0.5 U of Exonuclease (USB) and 0.5 U of Shrimp Alkaline Phosphatase (USB) were added to each remaining 7 μ L of PCR product and incubated for 30 min at 37 °C, then for 10 min at 90 °C. Cycle sequencing was conducted using the amplification primers and many additional internal

primers, some of which were designed for particular subclades or individual species. Cycle sequencing reactions employed the BigDye 3.1 (Applied Biosystems) chemistry and the recommended cycling conditions, and sequences were read using Applied Biosystems model 3100 or 3730 automated DNA sequencers. Nearly all fragments were confirmed by sequencing both DNA strands. Sequences were checked and overlapping fragments concatenated using Sequencer 4.5 (Genecodes). In nuclear DNA sequences, heterozygous nucleotides were assigned the corresponding IUPAC ambiguity codes. All sequence alignments were resolved readily by eye, except for several intron sites with single-nucleotide repeats that had high levels of indel mutation homoplasy.

2.3. Processing of museum-skin samples

Because of the degradation of the template DNAs in dried museum skin samples and the concomitantly higher risk of PCR template contamination, we employed a number of specialized protocols for the DNA extraction and PCR assembly of NDII sequences derived from museum skin toe-pads. All toe-pad DNA extractions and PCR set-ups were conducted in a custom-designed and equipped laboratory dedicated solely to work with degraded DNA. Protocols included many precautions that are recommended for ancient DNA studies on older materials (Willerslev and Cooper, 2005), including: physical and air-handling isolation of the degraded-DNA laboratory; prohibition of travel of personnel, reagents, samples, or equipment from the standard molecular markers lab to the degraded-DNA lab; set-up of all reactions within a laminar-flow clean bench with ISO class V air filtering; frequent sterilization of surfaces and equipment with intense 254 nm UV irradiation and 10% sodium hypochlorite (chlorine bleach); and interspersions of negative control reactions at both the extraction (1 control:1 tissue-containing extraction) and PCR (also 1:1) stages. PCR controls differed from the adjacent PCR reactions only in the absence of extracted DNA template. Despite the several hundred sets of PCR reactions required to complete these sequences, we saw no evidence of PCR amplification in any of the negative control PCR reactions, nor in the reactions to which we added solutions from the extraction negative controls.

Conspecific replicates were extracted and amplified from different skin specimens with planned temporal breaks of at least several weeks (and usually 2–3 months), over which interval many skin-snip samples of other taxa were processed using the same laboratory facility, equipment, primers, and reagents. The later comparison of these independently processed conspecific sequences helped ensure that any contaminant fragments would be identified as such. We identified no situations where a single PCR fragment or a concatenated NDII sequence differed by an unexpected magnitude from its conspecific replicate.

All skin-snip DNA extractions were performed with DNAeasy kits (Qiagen). Degraded DNA PCR amplifica-

tions targeted short (100–500 bp), overlapping regions of the NDII gene. We employed the NDII-flanking primers METb and TRPc as well as several dozen additional primers within binding sites within the NDII coding region, many designed for particular subclades or individual taxa. Sealed PCR reaction tubes were transported to the standard laboratory for the thermal cycling through automated sequencing steps described above.

2.4. Phylogenetic analysis

To reconstruct phylogenies, we used Bayesian methods as implemented in MrBayes 3.1 (Huelsenbeck and Ronquist, 2001; Ronquist and Huelsenbeck, 2003), and maximum parsimony as implemented in Paup* 4.0b10 (Swofford, 2002). Because of the heterogeneous composition of the sequence obtained from different taxa, we ran phylogenetic reconstructions on four datasets: (1) mtDNA sequences from all taxa surveyed, including 87 taxa from which we obtained 4108 nucleotides of mtDNA protein-coding sequence and 30 additional taxa from which we obtained only the 1041-nucleotide NDII gene; (2) mtDNA sequences from only the 87 more robustly sampled taxa; (3) four-locus combined intron sequences totaling 2974 aligned nucleotides from those same 87 taxa; and (4) all mtDNA and intron data combined for the complete set of 117 taxa.

Bayesian MCMC chains were sampled every 100 generations, with two independent sets of three heated and one unheated chains. All analyses employed the default flat Dirichlet priors. Stationarity was evaluated graphically for all parameters, and by monitoring the convergence of the standard deviation of split frequencies in the two independent sets of chains. Chains were run for 2×10^6 generations after the average standard deviation of split frequencies fell below 0.01 (0.02 for runs involving four or five data partitions); samples from earlier generations were discarded. Congruence between independent runs based on identical datasets was assessed by comparing parameter estimates, tree topologies, and posterior probability scores for individual branches. Parameters were estimated separately for 1–5 data partitions, depending on the loci included in a particular Bayesian search. All mitochondrial coding sequences were grouped in a single partition. In analyses that included nuclear loci, each intron was represented by a separate partition.

Parsimony analyses were conducted via full heuristic searches with 100 stepwise addition replicates. In analyses that included only mitochondrial coding sequences, transitional substitutions at third-position codon sites were given one-fifth the weight of third-position transversions and all changes at first- and second-position sites. In analyses that included only nuclear intron sequences, all sites were weighted equally. In analyses that included both mtDNA and intron sequences, mtDNA third-position transitions were downweighted as in the mtDNA-only searches, with all other sites equally weighted. Support for individual nodes in parsimony reconstructions was assessed via

heuristic bootstrap searches with five addition sequence replicates and 100 bootstrap replicates. In MP analyses of the nuclear intron sequences alone, the high similarity of many congeneric sequences resulted in many thousands of equally parsimonious shortest trees, and consequently, extremely long search times; these analyses were therefore run for a maximum of 1×10^7 TBR rearrangements per addition sequence replicate.

In all reconstructions, deletions were treated as missing data, except that the intron alignments contained a small number of indel sites (see Section 3) that could not be reliably aligned, and these sites were excluded entirely. Indel mutations that could be reliably aligned were later mapped onto the reconstructed topologies using MacClade 4.08 (Maddison and Maddison, 2005). In MP searches weighted by codon position, the frame-shifted 10-bp overlap between the ATPase6 and ATPase8 mtDNA genes was excluded, as each of these bases occupies two codon positions. All reconstructions were rooted to the outgroup taxa *Bombycilla*, *Catharus*, and *Myadestes*, but these are not shown in the trees.

3. Results

3.1. Sequence characteristics

Sequence alignments were straightforward except for one 6–12 bp region in each of the Fib-5, Fib-7, and Rho-1 intron comparisons, each of which involved single-nucleotide repeats of variable length that had unusually high insertion/deletion mutation rates leading to substantial homoplasy. Excluding these short regions of questionable alignment, we found a total of 66 indels among the ingroup taxa, of which 43 were present in more than one taxon and therefore represent potential synapomorphies.

All mitochondrial coding sequences were of identical length, except that the two NDII replicates from different *Cinnyricinclus leucogaster* individuals shared an unusual 1-codon (alanine) insertion just before the stop codon, making the gene 1044 nucleotides in length in this one taxon. The NDII amino acid sequence of this species was otherwise typical. With this one-taxon insertion excluded, the total aligned length of mtDNA coding sequence was 4118 nucleotides. In comparisons among the 84 ingroup taxa for which we had all five mtDNA gene sequences,

1906 mtDNA nucleotide sites were variable, of which 1723 were potentially phylogenetically-informative; an additional 222 sites varied in comparisons that included the three outgroup taxa.

In comparisons among these 84 ingroup and 3 outgroup taxa, the total aligned length of combined intron sequence was 2974 nucleotides, of which 1298 were variable and 612 parsimony informative. Considered by locus, the β -fibrinogen intron 5 alignment was 542 nucleotides, including 242 variable and 105 informative sites; β -fibrinogen 7 was 880 nucleotides, including 395 variable and 198 informative sites; TGFB2-5 was 562 nucleotides, including 234 variable and 106 informative sites; and rhodopsin 1 was 990 nucleotides, including 427 variable and 203 informative sites. All intron alignment lengths reported here exclude insertion sites present only in single taxa, as well as sites with questionable alignment.

Table 2 summarizes the post-burn-in means of parameters estimated for each data partition in the combined analysis of the 87 taxa for which we had the complete set of mitochondrial and nuclear sequences. As expected, all loci showed a predominance of transitional substitutions, but this bias was substantially greater for the mtDNA partition. As the mtDNA sequences were all protein-coding with relatively high amino acid conservation, they had a greater proportion of sites estimated as invariant than did the four intron sequences. Likewise, the shape parameter (α) of the gamma distribution of rate variation among sites was higher for all nuclear intron loci than for the mtDNA dataset, and much greater for the two β -fibrinogen introns than for the TGFB or rhodopsin introns; these high α values indicate that these loci had lower coefficients of variation in rate among sites.

To compare relative rates of nucleotide evolution among loci, we calculated pairwise ML distances in Paup* using the parameters summarized in Table 2, and plotted pairwise divergences for each intron locus against the corresponding mtDNA divergence calculated from the combined NDII, COI, COII, ATPase6, and ATPase8 genes (Fig. 1). Overall rates of ML divergence were approximately tenfold lower for all nuclear loci relative to the mitochondrial divergence. Relative rates were similar across all four intron loci; ML distances were nearly identical among the Fib-5, Fib-7, and Rho-1 introns, and about 25% greater at the TGFB2-5 locus (Fig. 1).

Table 2
Estimated model parameters for each of five loci under the HKY + G + I model of sequence evolution

Locus	Relative substitution rates						Base frequencies				α	p(I)
	A-C	A-G	A-T	C-G	C-T	G-T	A	C	G	T		
mtDNA	0.29	12.65	0.52	0.21	6.71	1.00	0.37	0.40	0.08	0.15	0.82	0.47
Fib-5	1.71	6.80	0.92	1.81	4.26	1.00	0.31	0.16	0.19	0.34	12.14	0.10
Fib-7	1.17	4.15	0.55	1.80	3.71	1.00	0.32	0.17	0.18	0.33	57.83	0.14
TGFB2-4	0.92	4.12	0.91	1.62	2.58	1.00	0.24	0.23	0.21	0.32	2.23	0.16
Rho-1	1.41	4.52	0.98	1.64	5.05	1.00	0.24	0.23	0.25	0.28	2.03	0.09




Fig. 1. Relative sequence divergence at protein-coding mtDNA genes versus four nuclear intron loci based on pairwise distances among 84 Sturnidae and Mimidae species and three outgroup taxa. (A) β -Fibrinogen intron 5, (B) β -fibrinogen intron 7, (C) rhodopsin intron 1, and (D) transforming growth factor β -2 intron 5. All comparisons were based on maximum-likelihood distances calculated using the locus-specific parameters given in Table 2. The dashed horizontal line is a visual reference to facilitate comparisons among nuclear loci, which indicate a $\sim 25\%$ faster rate of divergence at the TGFB2 intron 5 locus (D) relative to the other three intron loci (A–C). Mitochondrial distances were based on 4116 nucleotides of aligned sequence from five genes.

3.2. Phylogenetic reconstructions

We found high congruence among phylogenetic reconstructions based on different subsets of the combined nuclear and mitochondrial datasets, and among topologies based on Bayesian versus parsimony reconstruction methods (Figs. 2–4). In all reconstructions, the majority of nodes received substantial posterior probability ($>90\%$) and bootstrap ($>70\%$) support, and in no case did one of these moderately to highly supported nodes in a given topology conflict with a similarly well-supported node in an alternative reconstruction. The highest level of resolution was found in the combined-data reconstructions that included both mitochondrial and nuclear loci (Fig. 4). All reconstructions were consistent in the identification of nine major clades within this combined radiation; the composition of these clades and the relationships among them are reviewed in the Discussion below.

A gene-tree based only on mitochondrial sequences is depicted in Fig. 2. This analysis, which involved a mixed

dataset of 4108 nucleotides for 87 taxa and 1041 nucleotides for the 30 remaining taxa, provided high resolution among most groups of allied species and genera, with somewhat more modest support for relationships among some of the more basal Sturnidae clades. In separate analyses (not shown) we investigated potential biases of including mitochondrial sequences of heterogeneous lengths in a single analysis by analyses of only the NDII region from taxa for which we had the full set of sequences; in no case did the position of these taxa differ in the reconstructions based on truncated data. We likewise confirmed the identification of many taxa, most importantly those based on degraded DNA samples derived from museum skin-snips, using the replicate NDII sequences listed in Appendix A. In all cases these replicate samples from different conspecific individuals had very low divergence and grouped together in tree-based reconstructions.

Gene-trees generated from single nuclear introns generally had low resolution, and we compare here the Bayesian tree based on the four intron loci combined with the




Fig. 3. Comparison of Bayesian 90% consensus trees for 82 Sturnidae and Mimidae taxa based on nucleotide substitutions in five mtDNA coding genes (left; 4116 bp) and four nuclear intron loci (right; 2974 bp). All resolved internodes lacking posterior probability scores were supported at 100%. Box symbols along branches indicate insertion (open boxes) and deletion (filled boxes) mutations in the four intron loci mapped onto these topologies via maximum parsimony. The 58 indel mutations indicated on the left-hand topology each mapped with only a single step; the 5 indels (A–F) indicated on the right-hand tree each required 2–3 steps. Trees were rooted to outgroups *Bombycilla*, *Myadestes*, and *Catharus* (not shown).




Fig. 4. Total evidence Bayesian consensus tree from analysis of combined mtDNA and nuclear intron sequences for 120 taxa. Numbers above branches indicate posterior probability values ≥ 90 ; numbers below branches indicate maximum parsimony bootstrap scores ≥ 70 ; missing values indicate scores below these thresholds. Tree was rooted to outgroups *Bombycilla*, *Myadestes*, and *Catharus* (not shown).




Fig. 5. Schematic representation of the relationships among, and species diversities within, the nine well-supported major clades of starlings, mockingbirds, and oxpeckers. The height of the triangle denoting each group is proportional to its species diversity, whereas the width approximates the relative divergence of the earliest bifurcation within the clade.

both likely represent old lineages with no close extant allies. Their sister relationship could therefore be driven by long-branch attraction (Felsenstein, 1978; Bergsten, 2005), although we note that these species group together in all reconstructions, and that they share a synapomorphic intron deletion.

Several relatively old and well differentiated lineages are also present within the large African clade, and some of these are included here only on the basis of NDII sequences; more extensive sequence data from *Grafisia*, *Neocichla*, *Saroglossa spiloptera*, *Lamprotornis purpureiceps*, and *Lamprotornis cupreocauda* would likely help resolve their placements in these reconstructions. Finally, the six clades with moderate to high species diversities each

include some subclades of species that radiated recently and rapidly, but within which resolution is weak.

4. Discussion

4.1. Major clades

All reconstructions were consistent in defining three major clades within the combined Sturnidae/Mimidae radiation (summarized in Fig. 5). These three clades correspond taxonomically to the families Buphagidae, Mimidae, and Sturnidae. They can be further subdivided into two major subclades within the Mimidae and six within the Sturnidae.

4.1.1. *Buphagidae* (oxpeckers)

All reconstructions were consistent in placing the two *Buphagus* oxpeckers together as the basal lineage within this entire radiation. With only three outgroup taxa, our present study was not designed to test the monophyly of *Buphagus* + Mimidae + Sturnidae, as previous studies with robust sampling of allied avian groups have provided universally strong support for this clade that forms the ingroup here (Cibois and Cracraft, 2004; Voelker and Spellman, 2004; Ericson and Johansson, 2003; Barker et al., 2004; Zuccon et al., 2006). The placement of *Buphagus* in our reconstructions confirms relationships seen in several recent studies, in which *Buphagus* has appeared in the same position relative to the Mimidae/Sturnidae (Cibois and Cracraft, 2004; Zuccon et al., 2006). In those previous studies the nodes defining the basal position of *Buphagus* were poorly supported, but with our more substantial taxon and nucleotide sampling, support for the basal position of the *Buphagus* lineage was high in both our mitochondrial-only and nuclear-only reconstructions (Figs. 2 and 3), and in the analyses of combined data where the defining internode had a 93% Bayesian posterior probability and a 96% MP bootstrap score (Fig. 4).

Over the past century, *Buphagus* has either been placed in the monotypic family Buphagidae (Fry et al., 2000), or lumped into the Sturnidae, with most recent taxonomic treatments following this latter classification (e.g., Sibley and Monroe, 1990; Feare and Craig, 1999), often with the Buphaginae ranked as a subfamily (Amadon, 1943, 1956, 1962; Dickinson, 2003). Our strong and independent nuclear and mitochondrial evidence for the basal placement of *Buphagus*, in conjunction with the identical, but more tenuous, results from other recent phylogenetic studies, argues for the recognition of the family Buphagidae in any classification that treats the Mimidae and Sturnidae as separate families, as the inclusion of *Buphagus* renders the family Sturnidae paraphyletic.

4.1.2. *Mimidae* (mockingbirds and thrashers)

In all reconstructions (Figs. 2–5), the traditional Mimidae form a well supported clade with two subclades, a division that has also been found in previous studies that included subsets of these Mimidae taxa (Arbogast et al., 2006; Cibois and Cracraft, 2004; Hunt et al., 2001; Barber et al., 2004; Zuccon et al., 2006). The most diverse Mimidae subclade includes the *Toxostoma* and *Oreoscoptes* thrashers, and all mockingbirds in the genera *Mimus*, *Nesomimus*, and *Mimodes*. Our sampling included only three of the ten *Toxostoma* species, but previous molecular phylogenies of this genus indicate that it is monophyletic (Zink et al., 1999). The monotypic Sage Thrasher *Oreoscoptes* is the sister taxon to the *Mimus* mockingbird group. Our sampling of *Mimus* and its close allies is congruent with recent studies showing that *Mimodes* and *Nesomimus* fall within *Mimus* (Barber et al., 2004; Arbogast et al., 2006), but some species' relationships within this combined mockingbird clade remain unclear, in part probably because several

mockingbird taxa (including all *Nesomimus* and *Mimodes*) were included in our trees only on the basis of NDII sequences. The low resolution at some of the corresponding internodes leaves open the identity of the continental sister lineage of the Galápagos mockingbirds (*Nesomimus*), an important question given the historical prominence of the *Nesomimus* radiation in Charles Darwin's conceptualization of evolutionary modification following from island colonization (Arbogast et al., 2006).

The second Mimidae subclade includes three genera of North and Central American catbirds (*Melanotis*, *Melanoptila*, and *Dumatella*), and four genera of thrashers endemic to the islands of the West Indies (*Ramphocinclus*, *Margarops*, *Allenia*, and *Cinclocerthia*). The two species of *Melanotis* are basal sister species within this subclade. *Melanoptila*, a monotypic genus endemic to the Yucatan Peninsula, is sister to a clade comprised of the four endemic Caribbean genera plus the Grey Catbird *Dumatella*, a long-distance migrant that breeds in continental North America and that over-winters throughout the Greater Antilles and broadly in North and Central America around the Caribbean basin. These relationships are largely congruent with those found previously in molecular phylogenies of this group (Hunt et al., 2001; Barber et al., 2004), except that our reconstructions resolve the placement of *Melanoptila* as basal to the Caribbean endemics + *Dumatella* group. This finding that a Yucatan endemic is basal to a largely Caribbean clade is suggestive of a pathway of colonization into the Antilles from Central America via Cuba as proposed by Hunt et al. (2001), but the placement of *Dumatella* within the Caribbean clade adds a complication to this simple colonization scenario, because it suggests alternatively that either (1) the West Indian taxa are derived from one or more colonization events by a previously migratory ancestor, or that (2) the *Dumatella* lineage has a West Indian origin and has re-evolved the trait of long-distance migration.

4.1.3. *Eurasian, South Asian, and Pacific Island Sturnidae*

The traditional Sturnidae form a monophyletic group when *Buphagus* is excluded and *Rhabdornis* is included (Fig. 5), with support for this Sturnidae clade high in both mtDNA- and intron-only reconstructions, as well as in the analyses of combined dataset. This group is further defined by an indel synapomorphy, a one-nucleotide deletion in the β -fibrinogen intron 7 locus (Fig. 3). The basal division within the Sturnidae separates a clade of largely South Asian and Pacific Island starlings from a group of largely Eurasian and African starlings (Fig. 5). The South Asian/Pacific Island group is further divided into two subclades, the smaller of which includes only the Phillipine endemic *Rhabdornis*. The larger Eurasian/African group is subdivided into four subclades (Fig. 5).

The affinities of the *Rhabdornis* "Phillipine creepers" were uncertain until molecular phylogenetic reconstructions based on the RAG-1 gene placed *Rhabdornis* clearly within the Sturnidae (Cibois and Cracraft, 2004).

Subsequent analyses of these RAG-1 sequences in conjunction with mtDNA and intron sequences clarified the position of *Rhabdornis* as the sister lineage to the South Asian/Pacific Island starling group (Zuccon et al., 2006), a finding mirrored in our reconstructions (Figs. 2–5).

The *Rhabdornis* lineage is sister to a large clade of Australasian mynas and starlings that includes a few lineages found on the Indian subcontinent and in continental south-east Asia, as well as several groups that have diversified more extensively on various archipelagos within the Indopacific region. This clade comprises species in the genera *Aplonis*, *Gracula*, *Mino*, *Basilornis*, and *Streptocitta*, as well as in the monotypic genera *Scissirostrum*, *Enodes*, *Ampeliceps*, and *Sarcops*. The phylogenetic affinities of these genera have been previously suspected based on shared morphological traits and geographical proximity (Feare and Craig, 1999). As viewed best in the ultrametric tree based on mtDNA sequences (Fig. 2), this entire group shares a recent common ancestor, and its remarkable species and morphological diversities result from a correspondingly recent period of rapid diversification. The *Aplonis* clade, which has explosively radiated across much of the Indopacific, is especially notable in this regard, as even within this recent broader group, *Aplonis* is defined by a long basal internode leading to a cluster of species with very low mitochondrial divergence (Fig. 2). Although our sample includes only 10 of the 22 *Aplonis* species, it includes representatives of most of the morphologically and geographically distinctive *Aplonis* subgroups, and thus suggests that this genus is monophyletic. Other island radiations within this broader clade include: (1) the monotypic and morphologically unusual genera *Enodes* and *Scissirostrum*, which are both endemic to Sulawesi and which are likely sister taxa; and (2) the *Basilornis*/*Sarcops*/*Streptocitta* mynas, which are distributed from the Philippines south to Sulawesi.

The remaining Sturnidae include one clade of ‘Eurasian starlings’ and three clades of African taxa. A deep division within the Eurasian clade separates the well-known European Starling *Sturnus vulgaris* (and its close relative *S. unicolor*, which is often treated as a subspecies of *vulgaris*; Feare, 1984; de la Cruz-Cardiel et al., 1997) from the remaining taxa, which comprise lineages that radiated relatively recently (Fig. 2). This group includes species placed in the genera *Acridotheres*, *Leucopsar*, *Creatophora*, and *Sturnus* by Dickinson (2003), but divided among as many as 7 (Feare and Craig, 1999) to 11 (Wolters, 1982) genera in other recent treatments. One monotypic genus within this clade, *Creatophora*, is notable for its entirely African distribution, but its affinities to this otherwise Eurasian group have long been suspected (Amadon, 1956; Feare and Craig, 1999). Although we are still lacking molecular phylogenetic information for 6 of the 26 species in this group, the available evidence suggests that *Acridotheres* (sensu Dickinson, 2003) is monophyletic, but that the lineages assigned to the three remaining genera have a more complicated history than reflected in any previous classifi-

cation (Table 1). The apparently rapid diversification of these lineages helps explain why the relationships within this group have been difficult to discern from morphological evidence.

4.1.4. African Sturnidae

The three African clades include one that is comprised of a pair that are likely sister taxa, the Amethyst Starling *C. leucogaster* and the Madagascar Starling *S. aurata*. These species are highly divergent from one another at both mtDNA and nuclear intron loci (Figs. 2–5), suggesting that they both represent relatively old relictual lineages. They are both found largely in forested ecotones, and both move nomadically in flocks of conspecifics, a trait that may have facilitated the colonization of Madagascar by *S. aurata* and of most of sub-Saharan Africa by *C. leucogaster*.

A second well-supported clade is comprised of all species of ‘red-winged’ starlings in the genus *Onychognathus*. Nine of the 11 species in this genus are found in sub-Saharan continental Africa, with one species (*frater*) endemic to Socotra Island off the Horn of Africa, and one species (*tristramii*) distributed along the Arabian coast of the Red Sea and north into the Sinai Peninsula and Israel. Our species sampling of this group is complete, save for one species from central western Africa (*neumannii*) that was only recently split from the more widespread *O. morio* (Craig, 1998). Our results confirm the monophyly of *Onychognathus*, which has long been suspected on the basis of the morphological similarities of its constituent species (Fry et al., 2000). At the species level, our trees are in substantial conflict with some previous hypotheses of relationship within *Onychognathus*, including the suggestion that the longer tails and more pronounced sexual dimorphism seen in *morio*, *fulgidus*, *blythii*, and *tristramii* indicate their superspecies-level affinity (Hall and Moreau, 1970), and a phylogenetic analysis of morphological and ecological characters (Craig and Hulley, 1992) that suggested that *alirrostris*/*blythii*/*salvadorii*/*neumannii*, *morio*/*tenuirostris*, and *fulgidus*/*walleri* each form clades. Our results show little support for these hypotheses and suggest that further work on this clade is warranted.

The largest African clade contains 34 species with a volatile genus-level history of classification (Table 1; also Fry et al., 2000). Our species sampling of this group is complete. Our trees support the monophyly and close genetic affinities of the three canopy forest *Poeoptera* species, which group into a deeper clade along with *Cinnyricinclus femoralis* and *Pholia sharpii*; these latter two species have frequently been considered congeneric in alternative classifications (Table 1). This well-supported group of five species is subsequently nested within a less strongly supported clade that also contains four deeply rooted, single-species lineages (*Speculipastor*, *Grafisia*, *Neocichla*, and *S. spiloptera*). The placement of *S. spiloptera*, the Spot-winged Starling, within the African starling group is unanticipated, as this species breeds in the Himalayan foothills and migrates nomadically east to the Indochinese

Peninsula, and it is the only member of this entire larger African group to occur outside of Africa. *S. spiloptera* is not closely allied to the one other *Saroglossa* species, the Madagascar Starling *S. aurata*.

We found two separate clades of species usually assigned to *Lamprotornis*. The smaller of these clades comprises two closely related species (*cupreocauda* and *purpureiceps*) previously recognized as a superspecies (Hall and Moreau, 1970; Fry et al., 2000) and placed together in *Hylopsar* by Feare and Craig (1999) on the basis of their unusual feather pigment structures. In all reconstructions that include these taxa, their sister lineage is the single species *Lamprotornis corruscus*, but this sister relationship between *cupreocauda/purpureiceps* and *corruscus* is not highly supported. Accordingly, the *corruscus* lineage appears to have originated early in the African radiation, much like *Neocichla*, *Speculipastor*, and *Grafisia*.

The larger *Lamprotornis* group comprises 22 species from sub-Saharan Africa. Support for the monophyly of this group is high in the sequence-based reconstructions (Figs. 2–4), and further supported by one indel synapomorphy (Fig. 3). Within this clade are four well-supported subclades, the most basal of which unites *hildebrandti* and *shellyi*, taxa that were formerly often considered conspecific (e.g., Amadon, 1962). Five species form a second clade notable for the extreme elongation of their tails, with the basal species (*australis*) showing an intermediate tail length. The long-tailed Ashy Starling *L. unicolor* is a member of this clade, rather than a member of the “*Spreo*” sub-group of *Lamprotornis* as often proposed (e.g., Feare and Craig, 1999). However, the long-tailed Golden-breasted Starling *Lamprotornis regius* is not a member of this “long-tailed” sub-group. Seven shorter-tailed and ground-foraging ‘glossy’ starling form a third clade. Finally, eight species form a group that various authors have separated, in many combinations of species and little consistency among recent classifications, into the genus *Spreo* (Table 1; also Fry et al., 2000). Our results provide strong evidence that all previous treatments of *Spreo* render *Lamprotornis* paraphyletic. Aside from this issue of classification, the general grouping of these eight species is consistent with many previous suggestions that subsets of these taxa are closely allied to one another (Hall and Moreau, 1970; Feare and Craig, 1999; Fry et al., 2000). These eight species have notably high variation in plumage coloration, tail length, habitat affinities, and mating systems (Feare and Craig, 1999).

Phylogenetic relationships among most genera of African starlings have been assessed previously based on cladistic analysis of external color, feather ultrastructure (Craig and Hartley, 1985), body shape, skeletal, and behavioral characters (Craig, 1997), although this latter analysis was characterized by the author as preliminary owing to the lack of data for many species. We note that few nodes are shared between the DNA-based trees reported here and the previous non-molecular phylogenies, possibly because the radiation of the African Sturnidae has involved

high rates of morphological change and concomitant morphological homoplasy.

4.2. Global biogeography and comparison with previous studies

Previous phylogenies for the Sturnidae/Mimidae clade provided the basis for biogeographic scenarios advanced by Sibley and Ahlquist (1990) and Zuccon et al. (2006). Our taxonomically more comprehensive and better resolved reconstructions allow us to evaluate some aspects of these prior historical hypotheses and suggest several modifications and alternatives.

Sibley and Ahlquist (1984, 1990) proposed that the common ancestor of the Sturnidae/Mimidae had a widespread distribution across the Northern Hemisphere during the Miocene, and that the subsequent long period of global cooling severed this range and fostered the diversification of the two families at more southern latitudes. This scenario was logical given their simple, but sparsely sampled, DNA–DNA hybridization-based trees, which did not include *Buphagus* and which divided the Mimidae cleanly from the Sturnidae with little phylogenetic structuring within either family. The temporal dating of this basal division was based on their assumption that the separation of these families was caused by early Pliocene climate change; because the relevant DNA–DNA hybridization distances were not congruent with dates derived from their previous (and chronologically much older) calibration points, Sibley and Ahlquist (1990) used their speculative late Miocene split between the Sturnidae/Mimidae to justify a twofold faster rate of genetic divergence for all birds with short generation times.

With better information on the pattern of diversification within these groups and a more robust calibration point for the Mimidae/Sturnidae split derived from Barker et al. (2004), Zuccon et al. (2006) examined several possible biogeographic histories for the Mimidae and Sturnidae, but favored a scenario much like that of Sibley and Ahlquist (1990) in which a forest-inhabiting Eurasian ancestral taxon gave rise first to the *Buphagus* lineage after colonizing Africa, and then to the respective Mimidae and Sturnidae clades after colonizing North America. The Old World Sturnidae then split into two clades, one that gave rise to *Rhabdornis* and then diversified in Wallacea, and the other which dispersed into Africa. They point out that the notable morphological diversity of this first “Wallacean” clade (which corresponds to our “South Asian/Pacific clade”; we prefer this alternate geographic descriptor because the majority of species in this clade occur outside of Wallacea proper) results from this group retaining the remnant lineages of an old radiation. In contrast, we found that both their trees and ours suggest instead that this largely island-inhabiting group has undergone a very recent period of explosive diversification and correspondingly shares a more recent common ancestor than any other speciose clade within the entire radiation. The remarkable diversity of this

group is therefore more likely a result of ecological release than an example of the relic retention of ancient differentiation.

In the Zuccon et al. (2006) scenario, the clade corresponding to our “Eurasian Starlings” represents a secondary colonization out of Africa, with the Madagascar Starling *S. aurata* also representing a colonization from Africa. Because the relationships among the Eurasian and various African clades are not well resolved, this out-of-Africa hypothesis for the Eurasian group is equally parsimonious with a number of alternative scenarios. The small number of early lineages within the Buphagidae/Mimidae/Sturnidae radiation and their simple pattern of geographical separation make it difficult to polarize biogeographic hypotheses with confidence, but we offer several further observations about the traits likely associated with the diversification of the group and suggest that the history of this group is more complicated than previously recognized. First, the present-day restriction of the basal lineage, the Buphagidae, to Africa could readily result from recent environmental changes elsewhere: the oxpeckers are obligately associated with large herbivores, and the presence of diverse megafauna communities across Eurasia and the Americas prior to the very late Pleistocene could have supported a geographically widespread oxpecker relative. The rapid extinction of the Pleistocene megafauna could have led to a parallel extinction of ancestral Buphagidae throughout North America and Eurasia. Unfortunately, it is not possible to reconstruct the geographic distribution of this common ancestor by reference to the distributions of outgroup taxa, as the two candidate sister clades both have very broad distributions across all (in the Turdidae + allies) or most (Cinclidae) continents. Moreover, as far as we know, there are no known fossils of the Buphagidae or its ancestral relatives.

Second, we suggest that the tendency to associate with large mammals is likely ancestral to the entire radiation, not a derived condition restricted to the Buphagidae: species in all of the major Sturnidae clades (except the forest-dwelling *Rhabdornis* creepers) regularly forage around the feet of large ungulates, and many frequently perch on these animals while foraging (Feare and Craig, 1999). Large mammal associations are more unusual in the Mimidae, but several populations of Galápagos mockingbirds are well known for their oxpecker-like behavior of eating ectoparasites, drinking blood, and picking at wounds on marine iguanas, seabirds, and sea lions (Curry and Anderson, 1987).

Third, we note that several clades contain lineages that have dispersed substantially beyond their current biogeographic centers of diversity. This is most notable in the Mimidae, in which the *Mimus* group appears to have colonized South America relatively recently; in the Asian taxon *S. piloptera*, which is nested within the large African group; and in the African *Creatophora cinerea*, which is nested with the Eurasian group. Many additional species of Sturnidae move nomadically in large flocks during at

least part of the year, often tracking fruit resources (Feare and Craig, 1999). This tendency for flocks of birds to disperse together likely facilitated their colonizations of new regions (Clegg et al., 2002), including the many Sturnidae and Mimidae populations now present on remote islands.

Finally, we found that several of the most diverse clades within the radiation are defined by long basal internodes (shown schematically in Fig. 5), such that the extant diversity of these groups results from a lineage that persisted for substantial periods before diversifying into the presently extant lineages. It is likely that at least some now-extinct lineages were contemporaneous with these now-basal lineages, but we know nothing about their diversity or geographic distributions. Factoring these patterns into even the most simple biogeographic scenarios adds substantial complexity. For example, it is possible that the early split between the Mimidae and Sturnidae resulted from the colonization of the New World by the ancestral mimid as suggested by both Sibley and Ahlquist (1990) and Zuccon et al. (2006), but given the much more recent basal split within the Mimidae clade (Fig. 5), it is equally possible that the ancestral mimid lineage persisted in the Old World for a substantial time and that the radiation of extant Mimidae occurred when one member of this group colonized the New World at a much later point.

4.3. Taxonomic recommendations

Our taxonomic recommendations are based on several conservative criteria: (A) the assignment of names at and above the genus level to monophyletic groups; (B) applying taxonomic revisions only when newly understood relationships are well supported by independent lines of evidence, such as robust and congruent mtDNA and nuclear gene trees; and (C) holding off on revisions in situations where information on some relevant taxa is missing, and where the inclusion of those missing lineages might alter the preferred classification.

At the family level, we recommend the recognition of the Buphagidae, Mimidae, and Sturnidae, as summarized in Fig. 5.

In classifications that include major divisions within families (e.g., subfamilies or tribes), we recommend the further subdivision of the Mimidae into two groups, and the Sturnidae into six groups, also as summarized in Fig. 5.

At the genus level, our results identify a number of genera that are not monophyletic, including *Mimus*, *Basilornis*, *Sturnus*, *Saroglossa*, *Cinnyricinclus*, *Lamprotonis*, and *Spreo*. We currently lack phylogenetic information on only a single species of *Mimus*, and we concur with recent suggestions (e.g., Barber et al., 2004) to merge *Nesomimus* and *Mimodes* into *Mimus*. We lack potentially important species of both *Basilornis* and *Sturnus*, and we therefore do not yet recommend an alternative classification of these genera, although we anticipate generic changes when the requisite data become available. As the two *Saroglossa* species are not sister-taxa (nor apparently otherwise closely

related), we recommend referring *aurata* to *Hartlaubius* Bonaparte 1853. Similarly we recommend moving *Cinnyricinclus femoralis* (but not *C. leucogaster*) and *P. sharpii* to *Poeoptera* Bonaparte 1854.

We follow Feare and Craig (1999) in recognizing *Hylopsar* von Boetticher 1940 for the species more often treated as *L. cupreocauda* and *L. purpuriceps*. We further recommend making *L. corruscus* the sole member of *Notopholia* Roberts 1922, a genus for which *corruscus* is the type. Finally, we recommend subsuming all *Spreo* into *Lamprotornis*.

Acknowledgments

Genetic samples were kindly provided by: Alain Henache (Muséum National Histoire Naturelle, Parc de Clères); Brian Arbogast (Humboldt State University); Charles Foley (Tarangire Elephant Project); Chris Iles (Birdtrek); David Rimlinger (San Diego Zoo); David Willard and John Bates (Field Museum of Natural History); Denis O'Meally (The Australian Museum); Donna Dittmann, Van Remsen, and Frederick Sheldon (Museum of Natural Science, Louisiana State University); Eldredge Bermingham (Smithsonian Tropical Research Institute); Françoise Chapman Baudelot; Jill Marvin (Jardin Zoologique du Quebec); Jon Fjeldsa (Zoological Museum, University of Copenhagen); Jose Veiga (Museo Nacional de Ciencias Naturales); Kim Bostwick (Cornell University

Museum of Vertebrates); Lee Howell (Chessington World of Adventures); Lee Schoen (Houston Zoological Garden); Manuel Marin; Michael Braun (National Museum of Natural History); Michael Wells (Busch Gardens); Mike Myers (Audubon Zoo); Muchai Muchane (National Museums of Kenya); Pablo Tubaro (Museo Argentino de Ciencias Naturales Bernardino Rivadavia); Paul Sweet, Angelique Corthals, and Joel Cracraft (American Museum of Natural History); Robert Ricklefs (University of Missouri); Salit Kark (Hebrew University of Jerusalem); Seth Eiseb (National Museums of Namibia); Sharon Birks (Burke Museum, University of Washington); Susan Congdon (Disney's Animal Kingdom, Walt Disney World); and Tim Osborne (Tandela Ridge). For their help in arranging and conducting field collecting expeditions we thank Wilson Nderitu Watetu (Mpala Research Centre); Muchai Muchane and the staff of the Ornithology Department at the National Museums of Kenya; Belinda Low (Lewa Wildlife Conservancy); Samuel Andanje and Richard Bagine (Kenya Wildlife Service); the Laikipia Wildlife Forum; and the Mpala Research Centre. For laboratory assistance we thank Amanda Talaba, Isabella Fiorentino, Laura Stenzler, and especially Brynn McCleery. We thank John Klicka and an anonymous reviewer for helpful comments on the manuscript. This research was funded by grants from the National Science Foundation (DEB-0515981) and the Chapman Fund of the American Museum of Natural History.

Appendix A

Taxa included in this study, tissue types, collecting localities, institutional sources, and GenBank Accession Numbers

Taxon	Museum source ^a and sample no.	Type ^b	Locality ^c	Mitochondrial coding genes				Nuclear intron loci			
				NDII ^d	COI	COII	ATPases	Fib-5	Fib-7	Rho-1	TGFB2-4
<i>Rhabdornis mysticalis</i>	ZMUC-119523	T	Philippines, Cayapa, Baliuag	EF468190	EF486342	EF484316	EF486775	—	EF471842	EF472854	EF484113
<i>Rhabdornis inornatus</i>	FMNH-357586	T	Philippines, Mindanao, Mt. Kitanglad, Baungon	EF468189	EF484215	EF484315	EF486774	EF468321	EF471841	EF472853	EF484112
<i>Aplonis metallica</i>	UWBM-63222	T	Solomon Islands, Choiseul Island, Choiseul Prov.	EF468151	EF484181	EF484282	EF486740	EF468289	EF471808	EF472823	EF484079
<i>Aplonis cantoroides</i>	UWBM-63221	T	Solomon Islands, Choiseul Island, Choiseul Prov.	EF468146	EF484176	EF484277	EF486735	EF468284	EF471803	EF472818	EF484074
<i>Aplonis insularis</i>	AMNH-6574	T	Solomon Islands, Rennell Island, Tahatmatangi	EF468150	EF484180	EF484281	EF486739	EF468288	EF471807	EF472822	EF484078
<i>Aplonis brunneicapillus</i>	UWBM-60250	T	Solomon Islands, Guadalcanal Island, Gold River	EF468145	EF484175	EF484276	EF486734	EF468283	EF471802	EF472817	EF484073
<i>Aplonis grandis</i>	UWBM-67899	T	Solomon Islands, New Georgia, Arara	EF468149	EF484179	EF484280	EF486738	EF468287	EF471806	EF472821	EF484077
<i>Aplonis panayensis</i>	UWBM-64932	T	Captive bird (JBP)	EF468153	EF484183	EF484284	EF486742	EF468291	EF471810	EF472825	EF484081
<i>Aplonis minor</i>	FMNH-357661	T	Philippines, Mindanao, Mt. Kitanglad, Baungon	EF468152	EF484182	EF484283	EF486741	EF468290	EF471809	EF472824	EF484080
<i>Aplonis pelzelni</i>	AM-17550	T	Captive bird (TZP)	EF468154	EF484184	EF484285	EF486743	EF468292	EF471811	EF472826	EF484082
<i>Aplonis tabuensis</i>	UWBM-42839	T	Tonga, Eua, Houma	EF468155	EF484185	EF484286	EF486744	EF468293	EF471812	EF472827	EF484083
<i>Aplonis cinerascens</i>	UWBM-42817	T	Cook Islands, Rarotonga, Avarua	EF468147	EF484177	EF484278	EF486736	EF468285	EF471804	EF472819	EF484075
<i>Mino kreffti</i>	UWBM-76294	T	Solomon Islands, New Georgia, Lambet	EF468161	EF484191	EF484291	EF486750	EF468299	EF471818	EF472832	EF484089
<i>Mino anais</i>	LSUMNS-B20541	T	Captive bird (PAC)	EF468160	EF484190	EF484290	EF486749	EF468298	EF471817	EF472831	EF484088
<i>Basilornis celebensis</i>	CUMV-51469	T	Captive bird (HZG)	EF468156	EF484186	EF486341	EF486745	EF468294	EF471813	EF472828	EF484084
<i>Basilornis miranda</i>	FMNH-357664	T	Philippines, Mindanao, Mt. Kitanglad, Baungon	EF468157	EF484187	EF484287	EF486746	EF468295	EF471814	EF472829	EF484085
<i>Sarcops calvus</i>	FMNH-358605	T	Philippines, Sibuyan, Goangan	EF468163	EF484193	EF484293	EF486752	EF468301	EF471820	EF472834	EF484091
<i>Streptocitta albicollis</i>	CLOFBP-393013	F	Captive bird (SDZ)	EF468162	EF484192	EF484292	EF486751	EF468300	EF471819	EF472833	EF484090
<i>Enodes erythrophris</i>	AMNH-299958	S	Indonesia, Sulawesi	EF468227 ^d							
<i>Enodes erythrophris</i>	AMNH-299946	S	Indonesia, Sulawesi, Rujukan	EF468228							
<i>Scissirostrum dubium</i>	LSUMNS-B20447	T	Captive bird (PAC)	EF468164	EF484194	EF484294	EF486753	EF468302	EF471821	EF472835	EF484092
<i>Saroglossa spilopectera</i>	AMNH-203507	S	Thailand, Um Parig	EF468220							
<i>Saroglossa aurata</i>	FMNH-384699	T	Madagascar, Toliara, Sakaraha	EF468142	EF484172	EF484273	EF486731	EF468280	EF471799	EF472814	EF484070
<i>Ampeliceps coronatus</i>	BG-020148	F	Captive bird (BG)	EF468148	EF484178	EF484279	EF486737	EF468286	EF471805	EF472820	EF484076
<i>Gracula ptilogenys</i>	CUMV-15269	S	Sri Lanka, PundaLaya	EF468237							
<i>Gracula r. religiosa</i>	LSUMNS-B27008	T	Captive bird (HZG)	EF468159	EF484189	EF484289	EF486748	EF468297	EF471816	EF472830	EF484087
<i>Gracula religiosa indica</i>	CLOFBP-AKD06	F	Captive bird (MNHNPC)	EF468158	EF484188	EF484288	EF486747	EF468296	EF471815	EF484086	—
<i>Acridotheres grandis</i>	AMNH-9614	T	Malaysia, Kuala Lumpur	EF468168	EF484198	EF484298	EF486757	EF468305	EF471824	EF472838	EF484095
<i>Acridotheres cristatellus</i>	NMNH-B3778	T	Philippines, Luzon Island, Cagayan Prov.	EF468165	EF484195	EF484295	EF486754	EF468303	EF471822	EF472836	EF484093
<i>Acridotheres javanicus</i>	UWBM-67528	T	Captive bird (JBP)	EF468169	EF484199	EF484299	EF486758	EF468306	EF471825	EF472839	EF484096
<i>Acridotheres fuscus</i>	AMNH-9618	T	Malaysia, Kuala Lumpur	EF468166	EF484196	EF484296	EF486755	EF468304	EF471823	EF472837	EF484094
<i>Acridotheres ginginianus</i>	CLOFBP-AKD04	F	Captive bird (PAC)	EF468167	EF484197	EF484297	EF486756	—	—	—	—
<i>Acridotheres tristis</i>	UWBM-42794	T	Cook Islands, Mangaia, Lake Tiriara	EF468170	EF484200	EF484300	EF486759	EF468307	EF471826	EF472840	EF484097
<i>Leucopsar rothschildi</i>	UWBM-CHC001	T	Captive bird (WPZ)	EF468176	EF484205	EF484305	EF486764	EF468311	EF471831	EF472843	EF484102

<i>Sturnus nigricollis</i>	AMNH-105473	T	Captive bird (WCS)	EF468174 ^d																
<i>Sturnus nigricollis</i>	NMNH-B5709	T	Myanmar, Sagaing Division, Kan Blu, Kyat Thin	EF468173	EF484203	EF484303	EF486762	EF468309	EF471829	EF472842	EF484100									
<i>Sturnus contra</i>	AMNH-409725	S	China, Dalu	EF468175																
<i>Sturnus philippensis</i>	AMNH-790490	S	Phillippines, Mt. Calaviti	EF468180 ^d																
<i>Sturnus philippensis</i>	CLOFBP-BTK4	F	Captive bird (PAC)	EF468179	EF484208	EF484308	EF486767	EF468314	EF471834	EF472846	EF484105									
<i>Sturnus sinensis</i>	CLOFBP-AKD07	F	Captive bird (MNHNPC)	EF468183 ^d																
<i>Sturnus sinensis</i>	CLOFBP-BTK3	F	Captive bird (PAC)	EF468184																
<i>Sturnus malabaricus</i>	NMNH-B5708	T	Myanmar, Sagaing Division, Kan Blu, Kyat Thin	EF468178	EF484207	EF484307	EF486766	EF468313	EF471833	EF472845	EF484104									
<i>Sturnus albofrontatus</i>	AMNH-265248	S	Sri Lanka, Newara Eliya	EF468244																
<i>Sturnus pagodarum</i>	LSUMNS-B37263	T	Captive bird (PAC)	EF468187	EF484213	EF484313	EF486772	EF468319	EF471839	EF472851	EF484110									
<i>Sturnus roseus</i>	UWBM-46226	T	Kazakhstan, Almaty Oblsyy, Alma Ata	EF468181	EF484209	EF484309	EF486768	EF468315	EF471835	EF472847	EF484106									
<i>Sturnus sericeus</i>	CLOFBP-BTK1	F	Captive bird (PAC)	EF468182	EF484210	EF484310	EF486769	EF468316	EF471836	EF472848	EF484107									
<i>Sturnus cineraceus</i>	UWBM-47190	T	Russia, Khabarovskiy Krai, Khurmuli	EF468177	EF484206	EF484306	EF486765	EF468312	EF471832	EF472844	EF484103									
<i>Sturnus vulgaris</i>	CUMV-44167	T	USA, New York, Ithaca	EF468186	EF484212	EF484312	EF486771	EF468318	EF471838	EF472850	EF484109									
<i>Sturnus unicolor</i>	CLOFBP-3251760	B	Spain, Madrid Prov., Collado Villalba	EF468185	EF484211	EF484311	EF486770	EF468317	EF471837	EF472849	EF484108									
<i>Creatophora cinerea</i>	CLOFBP-DRRWS1	B	Kenya, Rift Valley Prov., Mpala Res. Centre	EF468172 ^d																
<i>Creatophora cinerea</i>	UWBM-70373	T	South Africa, Free State, Springfontein	EF468171	EF484201	EF484301	EF486760	EF468308	EF471827	EF472841	EF484098									
<i>Lamprotornis nitens</i>	CLOFBP-4A14817	B	Namibia, Tandala Ridge, Windpoort Farm	EF468122 ^d																
<i>Lamprotornis nitens</i>	UWBM-70405	T	South Africa, KwaZulu/Natal Prov., Ulundi	EF468121	EF484151	EF484252	EF486710	EF468261	EF471780	EF472798	EF484051									
<i>Lamprotornis chalybaeus</i>	CLOFBP-04208	B	Kenya, Eastern Prov., Lewa Wildlife Cons.	EF468112 ^d																
<i>Lamprotornis chalybaeus</i>	CLOFBP-04222	B	Kenya, Rift Valley Prov., Mpala Res. Centre	EF468111 ^d																
<i>Lamprotornis chalybaeus</i>	CLOFBP-09564	B	Kenya, Rift Valley Prov., Mpala Res. Centre	EF468113	EF484143	EF484244	EF486702	EF468254	EF471773	EF472792	EF484043									
<i>Lamprotornis chloropterus</i>	AMNH-764912	S	Uganda, Lendju, Mt. Matagi, Lake Albert	EF468232																
<i>Lamprotornis chalcurus</i>	NMK-4913	S	Uganda, Yalogi Gulu	EF468240 ^d																
<i>Lamprotornis chalcurus</i>	CUMV-30003	S	Nigeria, Northern Region, Kishi	EF468238																
<i>Lamprotornis splendidus</i>	FMNH-385397	T	Uganda, Southern Prov., Ngoto Swamp	EF468128	EF484158	EF484259	EF486717	EF468267	EF471786	EF472803	EF484057									
<i>Lamprotornis ornatus</i>	AMNH-266276	S	Sao Tome and Principe, Principe	EF468229																
<i>Lamprotornis iris</i>	CLOFBP-981868	F	Captive bird (DAK)	EF468119 ^d																
<i>Lamprotornis iris</i>	LSUMNS-B20774	T	Captive bird (PAC)	EF468118	EF484148	EF484249	EF486707	EF468259	EF471778	EF472796	EF484048									
<i>Lamprotornis purpureus</i>	CLOFBP-AKD01	F	Captive bird (MNHNPC)	EF468124	EF484154	EF484255	EF486713	EF468263	EF471782	EF472799	EF484053									
<i>Lamprotornis purpuroptera</i>	CLOFBP-04217	B	Kenya, Rift Valley Prov., Lake Bogoria	EF468125 ^d																
<i>Lamprotornis purpuroptera</i>	ZMUC-122452	T	Uganda, Queen Elizabeth	EF468126	EF484156	EF484257	EF486715	EF468265	EF471784	EF472801	EF484055									
<i>Lamprotornis caudatus</i>	LSUMNS-B19352	T	Captive bird (SAZ)	EF468110	EF484140	EF484241	EF486699	EF468251	EF471770	EF472790	EF484040									
<i>Lamprotornis regius</i>	CLOFBP-AKD02	F	Captive bird (DAK)	EF468127	EF484157	EF484258	EF486716	EF468266	EF471785	EF472802	EF484056									
<i>Lamprotornis mevesii</i>	CUMV-32262	S	Botswana, Bechuanaland, Tuli Block	EF468239 ^d																

(continued on next page)

Appendix A (continued)

Taxon	Museum source ^a and sample no.	Type ^b	Locality ^c	Mitochondrial coding genes				Nuclear intron loci			
				NDII ^d	COI	COII	ATPases	Fib-5	Fib-7	Rho-1	TGFB2-4
<i>Lamprotornis mevesii</i>	CLOFBP-NB1154	T	Namibia, Kunene Region	EF468120	EF484150	EF484251	EF486709	EF468260	EF471779	EF472797	EF484050
<i>Lamprotornis australis</i>	CLOFBP-056487	B	Namibia, Otjiwarongo District, Utsig Farm	EF468108 ^d							
<i>Lamprotornis australis</i>	CLOFBP-5772	T	Namibia, Otjiwarongo District, Utsig Farm	EF468109	EF484139	EF484240	EF486698	EF468250	EF471769	EF472789	EF484039
<i>Lamprotornis acuticaudus</i>	AMNH-347958	S	Zambia, Kasempa	EF468221							
<i>Lamprotornis corruscus</i>	NMK-16136	S	Kenya, Kipende, Wenje, Tana River	EF468231 ^d							
<i>Lamprotornis corruscus</i>	ZMUC-119491	T	Kenya, Malindi, Sokoke Forest	EF468114	EF484144	EF484245	EF486703	EF468255	EF471774	EF472793	EF484044
<i>Lamprotornis superbus</i>	CLOFBP-04209	B	Kenya, Eastern Prov., Lewa Wildlife Cons.	EF468130 ^d							
<i>Lamprotornis superbus</i>	CLOFBP-41313	B	Kenya, Rift Valley Prov., Mpala Res. Centre	EF468129	EF484159	EF484260	EF486718	EF468268	EF471787	EF472804	EF484058
<i>Lamprotornis hildebrandti</i>	CLOFBP-04206	B	Kenya, Eastern Prov., Lewa Wildlife Cons.	EF468117 ^d							
<i>Lamprotornis hildebrandti</i>	CLOFBP-04224	B	Kenya, Rift Valley Prov., Mpala Res. Centre	EF468116	EF484146	EF484247	EF486705	EF468257	EF471776	EF472795	EF484046
<i>Lamprotornis shelleyi</i>	NMK-15450	S	Kenya, E. Tenar, Tsavo	EF468215							
<i>Lamprotornis pulcher</i>	AMNH-822528	S	Mali, Timbuktu	EF468233 ^d							
<i>Lamprotornis pulcher</i>	CLOFBP-HOU1	F	Captive bird (HZG)	EF468123	EF484153	EF484254	EF486712	—	—	—	—
<i>Lamprotornis purpureiceps</i>	CLOFBP-NMK29	S	Uganda, Bwamba Forest, Mongiro	EF468214 ^d							
<i>Lamprotornis purpureiceps</i>	NMK-15364	S	Uganda, Bwamba, Makitengya	EF468225							
<i>Lamprotornis cupreocauda</i>	CUMV-15304	S	Ghana, Gold Coast, Winnebah	EF468230							
<i>Lamprotornis unicolor</i>	NMK-15473	S	Tanzania, Dodoma	EF468241 ^d							
<i>Lamprotornis unicolor</i>	CLOFBP-AKD03	F	Tanzania, Tarengire National Park	EF468131	EF484161	EF484262	EF486720	—	—	—	—
<i>Lamprotornis fischeri</i>	CLOFBP-04216	T	Kenya, Eastern Prov., Shaba	EF468115	EF484145	EF484246	EF486704	EF468256	EF471775	EF472794	EF484045
<i>Cinnyricinclus femoralis</i>	NMK-4889	S	Kenya, Chyulu	EF468217							
<i>Cinnyricinclus leucogaster</i>	UWBM-72577	T	Malawi, Mwanza District Mwanza	EF488683 ^d							
<i>Cinnyricinclus leucogaster</i>	LSUMNS-B22550	T	Captive bird (HZG)	EF488682	EF484136	EF484237	EF486695	EF468247	EF471766	EF472788	EF484036
<i>Spreo bicolor</i>	UWBM-70392	T	South Africa, Free State, Harrismith	EF468143	EF484173	EF484274	EF486732	EF468281	EF471800	EF472815	EF484071
<i>Spreo albicapillus</i>	CLOFBP-06001	T	Kenya, Eastern Prov., Kalacha	EF468141	EF484171	EF484272	EF486730	EF468279	EF471798	EF472813	EF484069
<i>Onychognathus morio</i>	CLOFBP-C7837	B	Kenya, Rift Valley Prov., Mpala Res. Centre	EF468133 ^d							
<i>Onychognathus morio</i>	UWBM-71314	T	South Africa, KwaZulu/Natal Prov., Melmoth	EF468132	EF484162	EF484263	EF486721	EF468270	EF471789	EF472805	EF484060
<i>Onychognathus tenuirostris</i>	FMNH-356559	T	Uganda, Western Prov., Rwenzori Mts.	EF468135	EF484165	EF484266	EF486724	EF468273	EF471792	EF472807	EF484063
<i>Onychognathus fulgidus</i>	AMNH-827360	S	Uganda, Bwamba Forest, Ntotoro	EF468224 ^d							

<i>Onychognathus fulgidus</i>	CUMV-33750	S	Uganda, Kibale Forest, Fort Portal	EF468223															
<i>Onychognathus walleri</i>	FMNH-439575	T	Malawi, Wilindi Forest, Chitipa	EF468137	EF484167	EF484268	EF486726	EF468275	EF471794	EF472809	EF484065								
<i>Onychognathus blythii</i>	AMNH-669364	S	Somalia, Sheilem	EF468245 ^d															
<i>Onychognathus blythii</i>	AMNH-669371	S	Somalia, Golis, Gidial Valley	EF468243															
<i>Onychognathus frater</i>	AMNH-669276	S	Socotra, Hornbill	EF468246 ^d															
<i>Onychognathus frater</i>	AMNH-669278	S	Socotra, Celilo Pass	EF468235															
<i>Onychognathus tristamii</i>	CLOFBP-ISR2	B	Israel, Southern District, Masada	EF468136	EF484166	EF484267	EF486725	EF468274	EF471793	EF472808	EF484064								
<i>Onychognathus nabouroup</i>	CUMV-32851	S	Namibia, Erongo, Homeb	EF468222															
<i>Onychognathus salvadorii</i>	CLOFBP-04211	T	Kenya, Eastern Prov., Shaba	EF468134	EF484164	EF484265	EF486723	EF468272	EF471791	EF472806	EF484062								
<i>Onychognathus albirostris</i>	CUMV-15258	S	Ethiopia, Lenafe	EF468236															
<i>Poeoptera stuhlmanni</i>	NMK-18046	S	Kenya, Nandi Forest	EF468218															
<i>Poeoptera kenricki</i>	ZMUC-123520	T	Tanzania, Udzungwa Forest, Iringa	EF468138	EF484168	EF484269	EF486727	EF468276	EF471795	EF472810	EF484066								
<i>Poeoptera lugubris</i>	AMNH-10691	T	Central African Rep., Sanga-Mbare, Bayanga	EF468139	EF484169	EF484270	EF486728	EF468277	EF471796	EF472811	EF484067								
<i>Pholia sharpii</i>	FMNH-356553	T	Uganda, Western Prov., Rwenzori Mts.	EF468140	EF484170	EF484271	EF486729	EF468278	EF471797	EF472812	EF484068								
<i>Grafisia torquata</i>	AMNH-162929	S	Cameroon, Pawa	EF468234 ^d															
<i>Grafisia torquata</i>	AMNH-162928	S	Cameroon, Pawa	EF468226															
<i>Speculipastor bicolor</i>	NMK-16455	S	Kenya, Kekeongole	EF468219 ^d															
<i>Speculipastor bicolor</i>	CLOFBP-06004	T	Kenya, Eastern Prov., Kalacha	EF468144	EF484174	EF484275	EF486733	EF468282	EF471801	EF472816	EF484072								
<i>Neocichla gutturalis</i>	NMK-15433	S	Tanzania, Itigi	EF468242 ^d															
<i>Neocichla gutturalis</i>	NMK-15432	S	Tanzania, Itigi	EF468216															
<i>Buphagus erythrorhynchus</i>	NMK-15490	S	Kenya, Eastern Prov., Archer's Post	EF468213															
<i>Buphagus africanus</i>	CLOFBP-DRR4	B	Kenya, Rift Valley Prov., Mpala Res. Centre	EF468188	EF484214	EF484314	EF486773	EF468320	EF471840	EF472852	EF484111								
<i>Dumatella carolinensis</i>	STRI-BHDCA4	T	Bahamas, Grand Bahama Island	EF468192	EF484216	EF484317	EF486776	EF468323	EF471844	EF472856	EF484115								
<i>Melanoptila glabrirostris</i>	LSUMNS-B0081	T	Mexico, Quintana Roo, Isla Cozumel	EF468197	EF484221	EF484322	EF486781	EF468328	EF471849	EF472861	EF484120								
<i>Mimus polyglottos</i>	LSUMNS-B21369	T	USA, California, San Bernardino	EF468202	EF484226	EF484327	EF486786	EF468333	EF471854	EF472866	EF484125								
<i>Mimus gilvus</i>	STRI-CCMG11	T	Trinidad, Chacachacare Island	EF468196	EF484220	EF484321	EF486780	EF468327	EF471848	EF472860	EF484119								
<i>Mimus gundlachi</i>	STRI-JAMGU1	T	Jamaica, Portland Ridge	EF468198	EF484222	EF484323	EF486782	EF468329	EF471850	EF472862	EF484121								
<i>Mimus thenca</i>	MNH-2676	T	Chile	EF468204	EF484228	EF484329	EF486788	EF468335	EF471856	EF472868	EF484127								
<i>Mimus longicaudatus</i>	LSUMNS-B5229	T	Peru, Lambayeque	EF468200	EF484224	EF484325	EF486784	EF468331	EF471852	EF472864	EF484123								
<i>Mimus saturninus</i>	CUMV-50582	T	Argentina, Buenos Aires Prov.	EF468203	EF484227	EF484328	EF486787	EF468334	EF471855	EF472867	EF484126								
<i>Mimus patagonicus</i>	CUMV-50577	T	Argentina, Jujuy Prov.	EF468201	EF484225	EF484326	EF486785	EF468332	EF471853	EF472865	EF484124								
<i>Mimus triurus</i>	CUMV-MACH14	T	Argentina, Buenos Aires Prov.	EF468205	EF484229	EF484330	EF486789	EF468336	EF471857	EF472869	EF484128								
<i>Nesomimus parvulus</i>			Arbogast et al. (2006)	AY311587															
<i>Nesomimus trifasciatus</i>			Arbogast et al. (2006)	AY311551															
<i>Nesomimus macdonaldi</i>			Arbogast et al. (2006)	AY311566															
<i>Nesomimus melanotis</i>			Arbogast et al. (2006)	AY311577															
<i>Oreoscoptes montanus</i>	LSUMNS-B19513	T	USA, California, Barstow	EF468206	EF484230	EF484331	EF486790	EF468337	EF471858	EF472870	EF484129								
<i>Mimodes graysoni</i>			Barber et al. (2004)	AY758199															

(continued on next page)

Appendix A (continued)

Taxon	Museum source ^a and sample no.	Type ^b	Locality ^c	Mitochondrial coding genes				Nuclear intron loci			
				NDII ^d	COI	COII	ATPases	Fib-5	Fib-7	Rho-1	TGFB2-4
<i>Toxostoma rufum</i>	LSUMNS-B0490	T	USA, Louisiana, Cameron Parish	EF468209	EF484233	EF484334	EF486793	EF468340	EF471861	EF472873	EF484132
<i>Toxostoma cinereum</i>	LSUMNS-B6745	T	Mexico, Baja California	EF468208	EF484232	EF484333	EF486792	EF468339	EF471860	EF472872	EF484131
<i>Toxostoma curvirostre</i>			Barber et al. (2004)	AY758201							
<i>Ramphocinclus brachyurus</i>	STRI-SLRBR2	T	St. Lucia	EF468207	EF484231	EF484332	EF486791	EF468338	EF471859	EF472871	EF484130
<i>Melanotis caerulescens</i>	LSUMNS-B0022	T	Mexico, Puebla	EF468193	EF484217	EF484318	EF486777	EF468324	EF471845	EF472857	EF484116
<i>Melanotis hypoleucus</i>	CUMV-44026	T	Mexico, Chiapas	EF468199	EF484223	EF484324	EF486783	EF468330	EF471851	EF472863	EF484122
<i>Allenia fusca</i>	STRI-DOMFU3	T	Dominica, Springfield	EF468195	EF484219	EF484320	EF486779	EF468326	EF471847	EF472859	EF484118
<i>Margarops fuscatus</i>	STRI-BUMFT1	T	Antigua and Barbuda, Barbuda	EF468194	EF484218	EF484319	EF486778	EF468325	EF471846	EF472858	EF484117
<i>Cinclocerthia ruficauda</i>	STRI-GUCRU1	T	Guadaloupe	EF468191							
<i>Bombycilla cedrorum</i>	CUMV-50897	T	USA, New York, Tompkins County	EF468210	EF484234	EF484335	EF486794	EF468341	EF471862	EF472874	EF484133
<i>Catharus guttatus</i>	CUMV-50482	T	USA, New York, Nassau County	EF468211	EF484235	EF484336	EF486795	EF468342	EF471863	EF472875	EF484134
<i>Myadestes townsendi</i>	LSUMNS-B20975	T	USA, California, San Bernardino County	EF468212	EF484236	EF484337	EF486796	EF468343	EF471864	EF472876	EF484135

^a Institutional sources of samples: AM: Australian Museum, Sydney, Australia; AMNH: American Museum of Natural History, New York, NY, USA; CUMV: Cornell University Museum of Vertebrates, Ithaca, NY, USA; FMNH: Field Museum of Natural History, Chicago, IL, USA; LSUMNS: Louisiana State University Museum of Natural Science, Baton Rouge, LA, USA; NMK: National Museums of Kenya, Nairobi, Kenya; NMN: National Museum of Namibia, Windhoek, Namibia; NMNH: National Museum of Natural History, Washington, DC, USA; UWBM: University of Washington Burke Museum, Seattle, WA, USA; ZMUC: Zoological Museum University of Copenhagen, Copenhagen, Denmark.

^b Sample types: B = blood; F = feather from live bird; T = frozen or buffer-preserved tissue; S = toe-pad shaving from museum skin.

^c Avicultural collections abbreviated: BG: Bush Gardens, Tampa, FL, USA; DAK: Disney's Animal Kingdom, Lake Buena Vista, FL, USA; HZG: Houston Zoological Garden, Houston, TX, USA; JBP: Jurong Bird Park, Singapore; TZP: Taronga Zoological Park, Sydney, Australia; MNHNPC: Muséum National Histoire Naturelle, Parc de Clères, Clères, France; PAC: Private avicultural collection; SAZ: San Antonio Zoo, San Antonio, TX, USA; SDZ: San Diego Zoo, San Diego, CA, USA; WCS: Wildlife Conservation Society, New York, NY, USA; WPZ: Woodland Park Zoo, Seattle, WA, USA.

^d Replicated conspecific specimens for which NDII samples were included in preliminary analyses, but which were not included in the phylogenetic trees reported here.

References

- Alström, P., Ericson, P.G.P., Olsson, U., Sundberg, P., 2006. Phylogeny and classification of the avian superfamily Sylvioidea. *Mol. Phylogenet. Evol.* 38, 381–397.
- Amadon, D., 1943. The genera of starlings and their relationships. *Am. Mus. Novitates* 1247, 1–16.
- Amadon, D., 1956. Remarks on the starlings, family Sturnidae. *Am. Mus. Novitates* 1803, 1–41.
- Amadon, D., 1962. Family Sturnidae. In: Mayr, E., Greenway, J.C., Jr. (Eds.), *Check-list of Birds of the World*, vol. XV. Museum of Comparative Zoology, Cambridge, MA, pp. 75–121.
- American Ornithologists' Union, 1998. *Checklist of North American Birds*, seventh ed. American Ornithologists' Union, Washington, DC.
- Arbogast, B.S., Drovetski, S.V., Curry, R.L., Boag, P.T., Seutin, G., Grant, P.R., Grant, B.R., Anderson, D.J., 2006. The origin and diversification of Galapagos mockingbirds. *Evolution* 60, 370–382.
- Banks, R.C., Cicero, C., Dunn, J.L., Kratter, A.W., Rasmussen, P.C., Remsen Jr., J.V., Rising, J.D., Stotz, D.F., 2002. Forty-third supplement to the American Ornithologists' Union Checklist of North American Birds. *Auk* 119, 897–906.
- Barber, B.R., Martinez-Gomez, J.E., Peterson, A.T., 2004. Systematic position of the Socorro mockingbird *Mimus graysoni*. *J. Avian Biol.* 35, 195–198.
- Barker, F.K., 2004. Monophyly and relationships of wrens (Aves: Troglodytidae): a congruence analysis of heterogeneous mitochondrial and nuclear DNA sequence data. *Mol. Phylogenet. Evol.* 32, 486–504.
- Barker, F.K., Cibois, A., Schikler, P., Cracraft, J., 2004. Phylogeny and diversification of the largest avian radiation. *Proc. Natl. Acad. Sci. USA* 101, 11040–11045.
- Beecher, W.J., 1953. A phylogeny of the oscines. *Auk* 70, 270–333.
- Bergsten, J., 2005. A review of long-branch attraction. *Cladistics* 21, 163–193.
- Brewer, D., 2001. *Wrens, Dippers, and Thrashers*. Christopher Helm, London.
- Christians, J.K., Evanson, M., Aiken, J.J., 2001. Seasonal decline in clutch size in European starlings: a novel randomization test to distinguish between the timing and quality hypotheses. *J. Anim. Ecol.* 70, 1080–1087.
- Cibois, A., Cracraft, J., 2004. Assessing the passerine “Tapestry”: phylogenetic relationships of the Muscicapoidea inferred from nuclear DNA sequences. *Mol. Phylogenet. Evol.* 32, 264–273.
- Clegg, S.M., Degnan, S.M., Kikkawa, J., Moritz, C., Estoup, A., Owens, I.P.F., 2002. Genetic consequences of sequential founder events by an island-colonizing bird. *Proc. Natl. Acad. Sci. USA* 99, 8127–8132.
- Cody, M.L., 2005. Family Mimidae (mockingbirds and thrashers). In: del Hoyo, J., Elliott, A., Christie, D.A. (Eds.), *Handbook of the Birds of the World, Cuckoo-shrikes to Thrushes*, vol. 10. Lynx Edicions, Barcelona, pp. 448–495.
- Cordero, P.J., Vinuela, J., Aparicio, M., Veiga, J.P., 2001. Seasonal variation in sex ratio and sexual egg dimorphism favouring daughters in first clutches of the spotless starling. *J. Evol. Biol.* 14, 829–834.
- Craig, A.J.F.K., 1997. A phylogeny for the African starlings (Sturnidae). *Ostrich* 68, 114–116.
- Craig, A.J.F.K., 1998. The timing of moult, morphology, and an assessment of the races of the Redwinged Starling. *Bonn. Zool. Beitr.* 39, 347–360.
- Craig, A.J.F.K., Hartley, A.H., 1985. The arrangement and structure of feather melanin granules as a taxonomic character in African starlings (Sturnidae). *Auk* 102, 629–632.
- Craig, A.J.F.K., Hulley, P.E., 1992. Biogeography and sympatry of Redwinged and Pale-winged Starlings in southern Africa. *J. Afr. Zool.* 106, 313–326.
- de la Cruz-Cardiel, P.J., Deceuninck, B., Peros, S.J., Elenma-Rossello, J.A., 1997. Allozyme polymorphism and interspecific relationships in the Common starling (*Sturnus vulgaris*) and Spotless starling (*S. unicolor*) (Aves: Sturnidae). *J. Zool. Syst. Evol. Res.* 35, 75–79.
- Curry, R.L., Anderson, D.J., 1987. Interisland variation in blood drinking by Galápagos Mockingbirds. *Auk* 104, 517–521.
- Davis, J., Miller, A.H., 1960. Family Mimidae. In: Mayr, E., Greenway, J.C., Jr. (Eds.), *Check-list of Birds of the World*, vol. IX. Museum of Comparative Zoology, Cambridge, MA, pp. 440–458.
- Derrickson, K.C., 1988. Variation in repertoire presentation in Northern Mockingbirds. *Condor* 90, 592–606.
- Dickinson, E.C., 2003. *The Howard and Moore Complete Checklist of the Birds of the World*, third ed. Christopher Helm, London.
- Duffy, D.L., Ball, G.F., 2002. Song predicts immunocompetence in male European starlings (*Sturnus vulgaris*). *Proc. R. Soc. Lond. B* 269, 847–852.
- Eberhard, J.P., Bermingham, E., 2004. Phylogeny and biogeography of the *Amazona ochrocephala* (Aves: Psittacidae) complex. *Auk* 121, 318–332.
- Ericson, P.G.P., Johansson, U.S., 2003. Phylogeny of Passerida (Aves: Passeriformes) based on nuclear and mitochondrial sequence data. *Mol. Phylogenet. Evol.* 29, 126–138.
- Feare, C., 1984. *The Starling*. Oxford University Press, Oxford.
- Feare, C., Craig, A., 1999. *Starlings and Mynas*. Christopher Helm, London.
- Felsenstein, J., 1978. Cases in which parsimony or compatibility methods will be positively misleading. *Syst. Zool.* 27, 401–410.
- Forbes, H.O., 1898. On an apparently new, and supposed to be now extinct, species of bird from the Mascarene Islands, provisionally referred to the genus *Necropsar*. *Bull. Liverpool Mus.* 1, 29–70.
- Fry, C.H., Keith, S., Urban, E.K. (Eds.), 2000. *The Birds of Africa*, vol. VI. Academic Press, London.
- Gentner, T.Q., Margoliash, D., 2003. Neuronal populations and single cells representing learned auditory objects. *Nature* 424, 669–674.
- Hall, B.P., Moreau, R.E., 1970. *An Atlas of Speciation in African Passerine Birds*. Trustees of the British Museum, London.
- Huelsenbeck, J.P., Ronquist, F., 2001. MRBAYES: Bayesian inference of phylogeny. *Bioinformatics* 17, 754–755.
- Hunt, J.S., Bermingham, E., Ricklefs, R.E., 2001. Molecular systematics and biogeography of Antillean thrashers, tremblers, and mockingbirds (Aves: Mimidae). *Auk* 118, 35–55.
- Kessing, B., Croom, H., Martin, A., McIntosh, C., McMillan, W.O., Palumbi, S.P., 1989. *The Simple Fool's Guide to PCR*, version 1.0. Special Publication of the Department of Zoology, University of Hawaii, Honolulu.
- Klicka, J., Voelker, G., Spellman, G.M., 2005. A molecular phylogeny of the “true thrushes” (Aves: Turdinae). *Mol. Phylogenet. Evol.* 34, 486–500.
- Komdeur, J., Wiersma, P., Magrath, M., 2002. Paternal care and male mate-attraction effort in the European starling is adjusted to clutch size. *Proc. R. Soc. Lond. B* 269, 1253–1261.
- Kroodsma, D.E., Byers, B.E., 1991. The function(s) of bird song. *Am. Zool.* 31, 318–328.
- Lovette, I.J., 2004. Molecular phylogeny and plumage signal evolution in a trans Andean and circum Amazonian avian species complex. *Mol. Phylogenet. Evol.* 32, 512–523.
- Maddison, W.P., Maddison, D.R., 2005. *MacClade: Analysis of Phylogeny and Character Evolution*, version 4.08. Sinauer Associates, Sunderland, MA.
- Malcarney, H.L., Martinez, D.R.C., Apanius, V., 1994. Sucrose intolerance in birds: simple nonlethal diagnostic methods and consequences for assimilation of complex carbohydrates. *Auk* 111, 170–177.
- Olson, S.L., Fleischer, R.C., Fisher, C.T., Bermingham, E., 2005. Expunging the ‘Mascarene starling’ *Necropsar leguati*: archives, morphology, and molecules topple a myth. *Bull. BOC* 125, 31–42.
- Pinxten, R., De Ridder, E., Balthazart, J., Ens, M., 2002. Context-dependent effects of castration and testosterone treatment on song in male European starlings. *Horm. Behav.* 42, 307–318.
- Polo, V., Veiga, J.P., Cordero, P.J., Viñuela, J., Monaghan, P., 2004. Female starlings adjust primary sex ratio in response to aromatic plants in the nest. *Proc. R. Soc. Lond. B* 271, 1929–1933.

- Primmer, C.R., Borge, T., Lindell, J., Sætre, G.-P., 2002. Single-nucleotide polymorphism characterization in species with limited available sequence information: high nucleotide diversity revealed in the avian genome. *Mol. Ecol.* 11, 603–612.
- Prychitko, T.M., Moore, W.S., 1997. The utility of DNA sequences of an intron from the beta-fibrinogen gene in phylogenetic analysis of woodpeckers (Aves: Picidae). *Mol. Phylogenet. Evol.* 8, 193–204.
- Ricklefs, R.E., Williams, J.B., 1984. Daily energy-expenditure and water-turnover rate of adult European Starlings (*Sturnus vulgaris*) during the nesting cycle. *Auk* 101, 707–716.
- Ronquist, F., Huelsenbeck, J.P., 2003. MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19, 1572–1574.
- Rubenstein, D.R., 2007a. Stress hormones and sociality: integrating social and environmental stressors. *Proc. R. Soc. Lond. B* 274, 967–975.
- Rubenstein, D.R., 2007b, in press. Temporal but not spatial environmental variation drives adaptive offspring sex allocation in a plural cooperative breeder. *Am. Nat.*
- Sgariglia, E.A., Burns, K.J., 2003. Phylogeography of the California Thrasher (*Toxostoma redivivum*) based on nested-clade analysis of mitochondrial-DNA variation. *Auk* 120, 346–361.
- Sheldon, F.H., Gill, F.B., 1996. A reconsideration of songbird phylogeny, with emphasis on the evolution of titmice and their sylvioid relatives. *Syst. Biol.* 4, 473–495.
- Sibley, C.G., Ahlquist, J.E., 1980. The relationships of the “primitive insect eaters” (Aves: Passeriformes) as indicated by DNA × DNA hybridization. In: Nohring, R. (Ed.) *Proceedings of the 17th International Ornithological Conference*, Berlin, pp. 1215–1220.
- Sibley, C.G., Ahlquist, J.E., 1984. The relationships of the starlings (Sturnidae: Sturnini) and the mockingbirds (Sturnidae: Mimini). *Auk* 101, 230–243.
- Sibley, C.G., Ahlquist, J.E., 1990. *Phylogeny and Classification of Birds*. Yale University Press, New Haven.
- Sibley, C.G., Monroe, B.L., 1990. *Distribution and Taxonomy of Birds of the World*. Yale University Press, New Haven.
- Stallcup, W.B., 1961. Relationships of some families of the suborder Passeres (songbirds) as indicated by comparisons of tissue proteins. *J. Grad. Res. Center Southern Methodist Univ.* 29, 43–65.
- Swofford, D.L., 2002. PAUP*. *Phylogenetic Analysis Using Parsimony (*and Other Methods)*, version 4. Sinauer Associates, Sunderland, MA.
- Voelker, G., Spellman, G.M., 2004. Nuclear and mitochondrial DNA evidence of polyphyly in the avian superfamily Muscicapoidae. *Mol. Phylogenet. Evol.* 30, 386–394.
- Willerslev, E., Cooper, A., 2005. Ancient DNA. *Proc. R. Soc. Lond. B* 972, 3–16.
- Wolters, H.E., 1982. *Die Vogelarten der Erde*. Paul Parey, Hamburg, Germany.
- Zink, R.M., Blackwell-Rago, R.C., 2000. Species limits and recent population history in the Curve-billed Thrasher. *Condor* 102, 881–886.
- Zink, R.M., Blackwell, R.C., Rojas-Soto, O., 1997. Species limits in the Le Conte’s Thrasher. *Condor* 99, 132–138.
- Zink, R.M., Dittmann, D.L., Klicka, J., Blackwell-Rago, R.C., 1999. Evolutionary patterns of morphometrics, allozymes, and mitochondrial DNA in thrashers (genus *Toxostoma*). *Auk* 116, 1021–1038.
- Zink, R.M., Kessen, A.E., Line, T.V., Blackwell-Rago, R.C., 2001. Comparative phylogeography of some aridland bird species. *Condor* 103, 1–10.
- Zuccon, D., Cibois, A., Pasquet, E., Ericson, P.G.P., 2006. Nuclear and mitochondrial sequence data reveal the major lineages of starlings, mynas and related taxa. *Mol. Phylogenet. Evol.* 41, 333–344.