

# Empirical likelihood based hypothesis testing

JOHN H.J. EINMAHL<sup>1</sup> and IAN W. McKEAGUE<sup>2</sup>

<sup>1</sup>*Department of Econometrics & OR, Tilburg University, P.O. Box 90153, 5000 LE Tilburg, Netherlands. E-mail: j.h.j.einmahl@uvt.nl*

<sup>2</sup>*Department of Statistics, Florida State University, Tallahassee FL 32306-4330, USA. E-mail: mckeague@stat.fsu.edu*

Omnibus tests for various nonparametric hypotheses are developed using the empirical likelihood method. These include tests for symmetry about zero, changes in distribution, independence and exponentiality. The approach is to localize the empirical likelihood using a suitable ‘time’ variable implicit in the null hypothesis and then form an integral of the log-likelihood ratio statistic. The asymptotic null distributions of these statistics are established. In simulation studies, the proposed statistics are found to have greater power than corresponding Cramér–von Mises type statistics.

*Keywords:* change point; distribution-free; exponentiality; independence; nonparametric likelihood ratio; symmetry; two-sample problem

## 1. Introduction

We develop an approach to omnibus hypothesis testing based on the empirical likelihood method. This method is known to be desirable and natural for deriving nonparametric and semi-parametric confidence regions for mostly finite-dimensional parameters; see Owen (2001) for an excellent account and an extensive bibliography on the topic. Just a few of the papers cited by Owen, however, consider problems of simultaneous inference, and none as far as we know has made a detailed study of omnibus hypothesis testing beyond the case of a simple null hypothesis.

Our approach is based on localizing the empirical likelihood using one or more ‘time’ variables implicit in the given null hypothesis. An omnibus test statistic is then constructed by integrating the log-likelihood ratio over those variables. We consider the proposed procedure to be potentially more efficient than the corresponding, often used, Cramér–von Mises type statistics. Four nonparametric problems will be studied in this way: testing for symmetry about zero; testing for a change in distribution (and the two-sample problem); testing for independence; and testing for exponentiality. These classical problems have been extensively studied in the literature, but use of the empirical likelihood approach in such contexts appears to be new. Actually, in Owen (2001) testing for symmetry and testing for independence are described as ‘Challenges for empirical likelihood’, since the standard method does not work properly here. Our localization approach, however, appears to be a

convenient adaptation, which makes empirical likelihood suitable for dealing with these fundamental statistical problems as well.

We first recall the case of a simple null hypothesis. Given independent and identically distributed (i.i.d.) observations  $X_1, \dots, X_n$  with distribution function  $F$ , consider  $H_0 : F = F_0$ , where  $F_0$  is a completely specified (continuous) distribution function. Define the localized empirical likelihood ratio

$$R(x) = \frac{\sup\{L(\tilde{F}) : \tilde{F}(x) = F_0(x)\}}{\sup\{L(\tilde{F})\}},$$

where  $L(\tilde{F}) = \prod_{i=1}^n (\tilde{F}(X_i) - \tilde{F}(X_i-))$ . The empirical distribution function  $F_n$  attains the supremum in the denominator, and the supremum in the numerator is attained by putting mass  $F_0(x)/(nF_n(x))$  on each observation up to and including  $x$  and mass  $(1 - F_0(x))/(n(1 - F_n(x)))$  on each observation beyond  $x$ . This easily leads to

$$\log R(x) = nF_n(x) \log \frac{F_0(x)}{F_n(x)} + n(1 - F_n(x)) \log \frac{1 - F_0(x)}{1 - F_n(x)}$$

and, provided  $0 < F_0(x) < 1$ ,

$$-2 \log R(x) = \frac{n(F_n(x) - F_0(x))^2}{F_0(x)(1 - F_0(x))} + o_p(1) \xrightarrow{\mathcal{D}} \chi_1^2 \quad (1.1)$$

under  $H_0$ . This is a special case of Owen's nonparametric version of the classical Wilks's theorem.

For an omnibus test (consistent against any departure from  $H_0$ ), however, we need to look at  $-2 \log R(x)$  simultaneously over a range of  $x$ -values. Taking the integral with respect to  $F_0$  leads to the statistic

$$T_n = -2 \int_{-\infty}^{\infty} \log R(x) dF_0(x).$$

If instead of integrating in  $T_n$ , we take the supremum over all  $x$ , we obtain essentially the statistic of Berk and Jones (1979), who showed that their statistic is more efficient in Bahadur's sense than any weighted Kolmogorov–Smirnov statistic. Li (2000) has introduced an extension of Berk and Jones's approach for a composite null hypothesis that  $F$  belongs to a parametric family of distributions. In that case,  $R(x) = R_\theta(x)$  for a parameter  $\theta$ , and Li suggests replacing the unknown  $\theta$  in Berk and Jones's statistic by its maximum likelihood estimator under the null hypothesis.

Clearly  $T_n$  is distribution-free and its small-sample null distribution can be approximated easily by simulation. Moreover, from (1.1) and a careful application of empirical process theory, it can be shown (cf. the proof of Theorem 1) that

$$T_n \xrightarrow{\mathcal{D}} \int_0^1 \frac{B^2(t)}{t(1-t)} dt$$

under  $H_0$ , where  $B$  is a standard Brownian bridge. Under  $H_0$ ,  $T_n$  is asymptotically equivalent to the Anderson–Darling statistic

$$A_n^2 = n \int_{-\infty}^{\infty} \frac{(F_n(x) - F_0(x))^2}{F_0(x)(1 - F_0(x))} dF_0(x)$$

and the limit distribution may be calculated using a series representation of Anderson and Darling (1952).

We investigate statistics of the form  $T_n$  for a variety of nonparametric hypotheses beyond the case of a simple null hypothesis. Testing for symmetry around zero can be handled using  $F(-x) = 1 - F(x-)$  and localizing at  $x > 0$ . To test for exponentiality, we localize using the memorylessness property of the exponential distribution. Our method also applies to the two-sample problem, and, more generally, to the nonparametric change-point problem; in that case, we localize at  $(x, t)$ , where  $t$  is the proportion of observation time before the change point. Testing for independent components in a bivariate distribution function  $F$  can be handled using  $F(x, y) = F(x, \infty)F(\infty, y)$ , with localization at  $(x, y)$ .

The paper is organized as follows. In Sections 2–5 we examine the four nonparametric testing problems mentioned above and derive likelihood ratio test statistics of the form  $T_n$ . Using empirical process techniques, we derive the limiting distribution of  $T_n$  in each case. Section 6 contains simulation results comparing the small-sample performance of each  $T_n$  with a corresponding Cramér–von Mises type statistic, and Section 7 is a concluding discussion. Proofs are postponed to Section 8. Tables of selected critical values for  $T_n$  are given in the Appendix.

## 2. Testing for symmetry

Much has been written on testing the symmetry of a distribution around either a known or unknown point of symmetry, some recent contributions being Diks and Tong (1999), Mizushima and Nagao (1998), Ahmad and Li (1997), Modarres and Gastwirth (1996), Nikitin (1996a), and Dykstra *et al.* (1995). Early papers include Butler (1969), Orlov (1972), Rothman and Woodroffe (1972), Srinivasan and Godio (1974), Hill and Rao (1977) and Lockhart and McLaren (1985).

Many of the papers cited above consider the case of a known point of symmetry and use a Cramér–von Mises type test statistic. We also assume that the point of symmetry is known, so without loss of generality it is assumed to be zero. Let  $X_1, \dots, X_n$  be i.i.d. with continuous distribution function  $F$ . The null hypothesis of symmetry about zero is

$$H_0 : F(-x) = 1 - F(x-), \quad \text{for all } x > 0.$$

The local likelihood ratio statistic is defined by

$$R(x) = \frac{\sup\{L(\tilde{F}) : \tilde{F}(-x) = 1 - \tilde{F}(x-)\}}{\sup\{L(\tilde{F})\}}, \quad x > 0.$$

As in the Introduction, the unrestricted likelihood in the denominator is maximized by setting  $\tilde{F} = F_n$ , the empirical distribution function. The supremum in the numerator can be found by treating  $\tilde{F}$  as a function of  $0 \leq p \leq 1$ , where  $\tilde{F}$  puts mass  $p/2$  on the interval  $(-\infty, -x]$ , mass  $p/2$  on  $[x, \infty)$  and mass  $1 - p$  on  $(-x, x)$ , with those masses divided equally among

the observations in the respective intervals. That is, the masses on the individual observations in the respective intervals are given by

$$\frac{p/2}{n\hat{p}_1}, \frac{p/2}{n\hat{p}_2}, \frac{1-p}{n(1-\hat{p})},$$

where  $\hat{p} = \hat{p}_1 + \hat{p}_2$ ,  $\hat{p}_1 = F_n(-x)$  and  $\hat{p}_2 = 1 - F_n(x-)$ . The numerator of  $R(x)$  is therefore the maximal value of

$$\left(\frac{p/2}{n\hat{p}_1}\right)^{n\hat{p}_1} \left(\frac{p/2}{n\hat{p}_2}\right)^{n\hat{p}_2} \left(\frac{1-p}{n(1-\hat{p})}\right)^{n(1-\hat{p})},$$

which is easily seen to be attained at  $p = \hat{p}$ . We thus obtain

$$\begin{aligned} \log R(x) &= n\hat{p}_1 \log \frac{\hat{p}}{2\hat{p}_1} + n\hat{p}_2 \log \frac{\hat{p}}{2\hat{p}_2} \\ &= nF_n(-x) \log \frac{F_n(-x) + 1 - F_n(x-)}{2F_n(-x)} \\ &\quad + n(1 - F_n(x-)) \log \frac{F_n(-x) + 1 - F_n(x-)}{2(1 - F_n(x-))}, \end{aligned} \quad (2.1)$$

where  $0 \log(a/0) = 0$ . Consider the test statistic

$$\begin{aligned} T_n &= -2 \int_0^\infty \log R(x) d\{F_n(x) - F_n(-x)\} \\ &= -2 \int_0^\infty \log R(x) dG_n(x), \end{aligned}$$

where  $G_n$  is the empirical distribution function of the  $|X_i|$ . Alternatively, we may write

$$T_n = -\frac{2}{n} \sum_{i=1}^n \log R(|X_i|).$$

Clearly,  $T_n$  is distribution-free; selected critical values are provided in Table A.1. The limit distribution of  $T_n$  is given by the following result.

**Theorem 1.** *Let  $F$  be continuous. Then, under  $H_0$ ,*

$$T_n \xrightarrow{\mathcal{D}} \int_0^1 \frac{W^2(t)}{t} dt,$$

where  $W$  is a standard Wiener process.

### 3. Testing for a change point

The nonparametric change-point testing problem has an extensive literature; recent contributions include Gombay and Jin (1999), Aly (1998), Aly and Kochar (1997), Ferger

(1994; 1995; 1996; 1998), McKeague and Sun (1996) and Szyszkowicz (1994). We consider the non-sequential (retrospective) situation with ‘at most one change’; see, for example, Csörgő and Horváth (1987) and Hawkins (1988).

Let  $X_1, \dots, X_n$  be independent, and assume that for some  $\tau \in \{2, \dots, n\}$  and some continuous distribution functions  $F, G$ ,

$$X_1, \dots, X_{\tau-1} \sim F \quad \text{and} \quad X_{\tau}, \dots, X_n \sim G,$$

with  $\tau, F$  and  $G$  unknown. We wish to test the null hypothesis of no change point,  $H_0 : F = G$ . Define the local likelihood ratio test statistic

$$R(t, x) = \frac{\sup\{L(\tilde{F}, \tilde{G}, \tau) : \tilde{F}(x) = \tilde{G}(x), \tau = [nt] + 1\}}{\sup\{L(\tilde{F}, \tilde{G}, \tau) : \tau = [nt] + 1\}}$$

for  $1/n \leq t < 1$  and  $x \in \mathbb{R}$ , with

$$L(\tilde{F}, \tilde{G}, \tau) = \prod_{i=1}^{\tau-1} (\tilde{F}(X_i) - \tilde{F}(X_{i-})) \prod_{i=\tau}^n (\tilde{G}(X_i) - \tilde{G}(X_{i-})).$$

Set  $n_1 = [nt]$ ,  $n_2 = n - [nt]$ , and let  $F_{1n}$  and  $F_{2n}$  be the empirical distribution functions of the first  $n_1$  observations and last  $n_2$  observations, respectively. Let  $F_n$  be the empirical distribution function of the full sample, so  $F_n(x) = (n_1 F_{1n}(x) + n_2 F_{2n}(x))/n$ . Then

$$\begin{aligned} \log R(t, x) &= n_1 F_{1n}(x) \log \frac{F_n(x)}{F_{1n}(x)} + n_1 (1 - F_{1n}(x)) \log \frac{1 - F_n(x)}{1 - F_{1n}(x)} \\ &\quad + n_2 F_{2n}(x) \log \frac{F_n(x)}{F_{2n}(x)} + n_2 (1 - F_{2n}(x)) \log \frac{1 - F_n(x)}{1 - F_{2n}(x)}, \end{aligned} \tag{3.1}$$

where  $0 \log(a/0) = 0$ . Consider the test statistic

$$\begin{aligned} T_n &= -2 \int_{1/n}^1 \int_{-\infty}^{\infty} \log R(t, x) dF_n(x) dt \\ &= -\frac{2}{n} \sum_{i=1}^n \int_{1/n}^1 \log R(t, X_i) dt. \end{aligned}$$

Clearly,  $T_n$  is distribution-free; selected critical values are provided in Table A.2. The limit distribution of  $T_n$  is given by the following result. Let  $W_0$  be a four-sided tied-down Wiener process on  $[0, 1]^2$  defined by  $W_0(t, y) = W(t, y) - tW(1, y) - yW(t, 1) + tyW(1, 1)$ , where  $W$  is a standard bivariate Wiener process.

**Theorem 2.** *Let  $F$  and  $G$  be continuous. Then, under  $H_0$ ,*

$$T_n \xrightarrow{\mathcal{D}} \int_0^1 \int_0^1 \frac{W_0^2(t, y)}{t(1-t)y(1-y)} dy dt.$$

Note that the classical two-sample problem could be handled in a similar way; see Einmahl and Khmaladze (2001) for recent progress on this problem along other lines and

for the references therein. We will briefly describe the two-sample problem here, but we will not provide a proof for this case, since it is similar to but easier than the proof for the change-point problem given in Section 7.

Let  $X_1, \dots, X_n$  be independent, and suppose that  $X_1, \dots, X_{n_1}$  ( $1 \leq n_1 < n$ ) have common continuous distribution function  $F$ , and  $X_{n_1+1}, \dots, X_n$  have common continuous distribution function  $G$ ; here  $F$  and  $G$  are unknown. We wish to test the null hypothesis of equal distributions,  $H_0 : F = G$ . Define the local likelihood ratio test statistic

$$R(x) = \frac{\sup\{L(\tilde{F}, \tilde{G}) : \tilde{F}(x) = \tilde{G}(x)\}}{\sup\{L(\tilde{F}, \tilde{G})\}}, \quad x \in \mathbb{R},$$

with

$$L(\tilde{F}, \tilde{G}) = \prod_{i=1}^{n_1} (\tilde{F}(X_i) - \tilde{F}(X_{i-})) \prod_{i=n_1+1}^n (\tilde{G}(X_i) - \tilde{G}(X_{i-})).$$

Let  $F_{1n}$  and  $F_{2n}$  be the empirical distribution functions of the first  $n_1$  and last  $n_2 := n - n_1$  observations respectively, and let  $F_n$  be the empirical distribution function of the pooled sample  $X_1, \dots, X_n$ . Then  $\log R(x)$  is equal to the right-hand side of (3.1). Consider the test statistic

$$T_n = -2 \int_{-\infty}^{\infty} \log R(x) dF_n(x) = -\frac{2}{n} \sum_{i=1}^n \log R(X_i);$$

again  $T_n$  is distribution-free.

**Theorem 2a.** *Let  $F$  and  $G$  be continuous and assume  $n_1, n_2 \rightarrow \infty$  as  $n \rightarrow \infty$ . Then, under  $H_0$ ,*

$$T_n \xrightarrow{\mathcal{D}} \int_0^1 \frac{B^2(y)}{y(1-y)} dy,$$

with  $B$  a standard Brownian bridge.

## 4. Testing for independence

The wide variety of tests for independence has been surveyed by Martynov (1992, Section 12). Here we consider a test for the independence of two random variables.

Let  $X_1, \dots, X_n$  be i.i.d. bivariate random vectors with distribution function  $F$  and continuous marginal distribution functions  $F_1$  and  $F_2$ . We wish to test the null hypothesis of independence:

$$H_0 : F(x, y) = F_1(x)F_2(y), \quad \text{for all } x, y \in \mathbb{R}.$$

Define the local likelihood ratio test statistic

$$R(x, y) = \frac{\sup\{L(\tilde{F}) : \tilde{F}(x, y) = \tilde{F}_1(x)\tilde{F}_2(y)\}}{\sup\{L(\tilde{F})\}}$$

for  $(x, y) \in \mathbb{R}^2$ , with  $L(\tilde{F}) = \prod_{i=1}^n \tilde{P}(\{X_i\})$ , where  $\tilde{P}$  is the probability measure corresponding to  $\tilde{F}$ . Then

$$\begin{aligned} \log R(x, y) = & nP_n(A_{11})\log \frac{F_{1n}(x)F_{2n}(y)}{P_n(A_{11})} + nP_n(A_{12})\log \frac{F_{1n}(x)(1 - F_{2n}(y))}{P_n(A_{12})} \\ & + nP_n(A_{21})\log \frac{(1 - F_{1n}(x))F_{2n}(y)}{P_n(A_{21})} + nP_n(A_{22})\log \frac{(1 - F_{1n}(x))(1 - F_{2n}(y))}{P_n(A_{22})}, \end{aligned}$$

where  $P_n$  is the empirical measure,  $F_{1n}$  and  $F_{2n}$  are the corresponding marginal distribution functions, and

$$A_{11} = (-\infty, x] \times (-\infty, y],$$

$$A_{12} = (-\infty, x] \times (y, \infty),$$

$$A_{21} = (x, \infty) \times (-\infty, y],$$

$$A_{22} = (x, \infty) \times (y, \infty).$$

Consider the test statistic

$$T_n = -2 \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \log R(x, y) dF_{1n}(x) dF_{2n}(y).$$

Clearly,  $T_n$  is distribution-free; selected critical values are provided in Table A.3. The limit distribution of  $T_n$  is given by the following result.

**Theorem 3.** *Let  $F_1, F_2$  be continuous. Then, under  $H_0$ ,*

$$T_n \xrightarrow{\mathcal{D}} \int_0^1 \int_0^1 \frac{W_0^2(u, v)}{u(1-u)v(1-v)} du dv,$$

where  $W_0$  is a four-sided tied-down Wiener process on  $[0, 1]^2$ .

The limit distribution above agrees with that in the change-point problem.

## 5. Testing for exponentiality

In this section we develop a likelihood ratio based test for exponentiality motivated by the memorylessness property of the exponential distribution. Cramér–von Mises type tests based on this property have been proposed by Angus (1982) and Ahmad and Alwasel (1999); we refer to these papers for references to the earlier literature.

Let  $X_1, \dots, X_n$  be i.i.d. non-negative random variables with distribution function  $F$ ,  $F(0-) = 0$ , and survival function  $S = 1 - F$ . Consider the null hypothesis

$$H_0 : S(x) = \exp(-\lambda x), \quad x \geq 0, \text{ for some } \lambda > 0.$$

The local likelihood ratio statistic based on the memorylessness property of the exponential distribution is

$$R(x, y) = \frac{\sup\{L(\tilde{S}) : \tilde{S}(x+y) = \tilde{S}(x)\tilde{S}(y)\}}{\sup\{L(\tilde{S})\}}$$

for  $x > 0$ ,  $y > 0$ , where

$$L(\tilde{S}) = \prod_{i=1}^n (\tilde{S}(X_{i-}) - \tilde{S}(X_i)).$$

Let  $F_n$  denote the empirical distribution function. It follows by a straightforward calculation that

$$\log R(x, y) = N_1 \log \frac{n(1-a)}{N_1} + N_2 \log \frac{n(a-b)}{N_2} + N_3 \log \frac{nb(1-a)}{N_3} + N_4 \log \frac{nab}{N_4}$$

where  $N_1 = nF_n(x \wedge y)$ ,  $N_2 = n(F_n(x \vee y) - F_n(x \wedge y))$ ,  $N_3 = n(F_n(x+y) - F_n(x \vee y))$ ,  $N_4 = n(1 - F_n(x+y))$ , and

$$a = \frac{N_2 + N_3 + 2N_4}{n + N_3 + N_4}, \quad b = \left( \frac{N_3 + N_4}{n - N_1} \right) a.$$

Consider the test statistic

$$T_n = -2 \int_0^\infty \int_0^\infty \log R(x, y) \hat{\lambda}^2 e^{-\hat{\lambda}(x+y)} dx dy,$$

with  $\hat{\lambda} = n / \sum_{i=1}^n X_i$ . This statistic is distribution-free (under  $H_0$  its distribution does not depend on the parameter  $\lambda$ ). Selected critical values for  $T_n$  obtained by simulation are displayed in Table A.4.

The asymptotic null distribution of  $T_n$  is given in the following result. Based on this result, selected critical values for the large-sample case are presented in the last row of Table A.4. Comparison of Tables A.1–A.4 shows that the convergence of  $T_n$  is much slower here than in the previous sections.

**Theorem 4.** Under  $H_0$ ,

$$T_n \xrightarrow{\mathcal{D}} 2 \int_0^1 \int_t^1 \frac{st}{(1-s)(1+t)} \left\{ \frac{B(st)}{st} - \frac{B(s)}{s} - \frac{B(t)}{t} \right\}^2 ds dt,$$

where  $B$  is a standard Brownian bridge.

## 6. Simulation results

In this section we present simulation results comparing the small-sample performance of the proposed likelihood ratio statistic  $T_n$  with that of a corresponding Cramér–von Mises type



statistic  $C_n$ . In each case the powers are based on 10 000 samples, and exact critical values are used (see the Appendix for the  $T_n$  critical values).

For the symmetry test, we compared  $T_n$  with

$$C_n = n \int_0^\infty \{1 - F_n(x-) - F_n(-x)\}^2 dG_n(x);$$

see Rothman and Woodroffe (1972). The alternatives are  $N(0.3, 1)$  and chi-squared centred about the mean; see Table 1.

For the change-point test, we compared  $T_n$  with

$$C_n = n \int_{1/n}^1 \int_{-\infty}^\infty \{F_{1n}(x) - F_{2n}(x)\}^2 dF_n(x) dt;$$

see Csörgő and Horváth (1988). The results are presented in Table 2.

For the test of independence, we compared  $T_n$  with

$$C_n = n \int_{-\infty}^\infty \int_{-\infty}^\infty \{F_n(x, y) - F_{1n}(x)F_{2n}(y)\}^2 dF_{1n}(x) dF_{2n}(y);$$

see Deheuvels (1981) or Martynov (1992, Section 12). The alternatives are bivariate normal with correlation  $\rho$ , and  $(U, \beta U + V)$ , where  $U, V$  are i.i.d. uniform on  $(0, 1)$ , for various values of  $\rho$  and  $\beta$ ; see Table 3.

**Table 1.** Power comparison of tests for symmetry at level  $\alpha = 0.05$  for  $n = 50$  and  $\alpha = 0.01$  for  $n = 100$

Alternative	$n = 50$		$n = 100$	
	$T_n$	$C_n$	$T_n$	$C_n$
$N(0.3, 1)$	0.539	0.516	0.629	0.600
Centred $\chi_1^2$	0.893	0.732	0.988	0.872
Centred $\chi_2^2$	0.505	0.433	0.647	0.495
Centred $\chi_3^2$	0.322	0.307	0.332	0.297

**Table 2.** Power comparison of tests for a change point,  $n = 50$ ,  $\alpha = 0.05$

$F$	$G$	$\tau = 11$		$\tau = 21$	
		$T_n$	$C_n$	$T_n$	$C_n$
$N(0, 1)$	$N(0, 16)$	0.210	0.129	0.735	0.356
$\text{Unif}(0, 1)$	$\text{Unif}(0.3, 1.3)$	0.512	0.446	0.837	0.661
$\text{Exp}(1)$	$\text{Exp}(2)$	0.236	0.229	0.418	0.333
$\text{Exp}(1)$	$\text{Exp}(3)$	0.506	0.479	0.789	0.683

**Table 3.** Power comparison of tests for independence at level  $\alpha = 0.05$ 

Alternative	$n = 20$		$n = 50$	
	$T_n$	$C_n$	$T_n$	$C_n$
$\rho = 0.4$	0.357	0.341	0.761	0.728
$\rho = 0.5$	0.550	0.520	0.937	0.915
$\beta = 0.5$	0.437	0.389	0.904	0.826
$\beta = 0.6$	0.573	0.523	0.974	0.935

**Table 4.** Power comparison of tests for exponentiality

Alternative	$n = 20$		$n = 30$	
	$T_n$	$C_n$	$T_n$	$C_n$
$\chi_4^2$	0.675	0.624	0.717	0.678
LN(0.8)	0.638	0.560	0.696	0.618
LN(1.0)	0.227	0.181	0.201	0.144
Weibull(1.5)	0.619	0.588	0.666	0.638

For the test of exponentiality, we compared  $T_n$  with

$$C_n = n \int_0^\infty \int_0^\infty \{S_n(x+y) - S_n(x)S_n(y)\}^2 \hat{\lambda}^2 e^{-\hat{\lambda}(x+y)} dx dy;$$

see Angus (1982). We used levels  $\alpha = 0.10$  for  $n = 20$ , and  $\alpha = 0.05$  for  $n = 30$ . The alternatives were chi-squared, lognormal and Weibull; see Table 4. The lognormal distribution with corresponding normal parameters  $\mu = 0$  and  $\sigma$  is denoted LN( $\sigma$ ); the Weibull distribution with scale parameter 1 and shape parameter  $c$  is denoted Weibull( $c$ ).

The proposed statistics show consistent improvement over the corresponding Cramér–von Mises statistics in *all* cases.

## 7. Discussion

We have developed a rather general localized empirical likelihood approach for testing certain composite nonparametric null hypotheses. We use integral type statistics to establish appropriate limit results. These statistics are somewhat related to Anderson–Darling type statistics, but have the advantage that the implicitly present weight function is automatically determined by the empirical likelihood. Clearly our tests are consistent (against all fixed alternatives). The proofs of our main results (see Section 8) require delicate arguments

concerning weighted empirical processes to handle ‘edge’ effects in the localized empirical likelihood.

Our approach is tractable in the four cases we have examined because the null hypothesis is expressed in terms of a relatively simple functional equation involving the distribution function(s). Another example in which our approach appears to be useful is in testing bivariate symmetry. More complex null hypotheses, however, might be difficult to handle via our localized empirical likelihood technique. In that sense the goodness-of-fit tests for *parametric* models in Li’s (2000) extension of Berk and Jones (1979) are complementary to the present paper (but, in contrast to our approach, the limit distribution is intractable). However, in the case of testing for exponentiality our test is simpler and more natural. For that case both Li’s and our approach can be extended to randomly censored data. Li’s approach is not applicable to the other cases we have considered.

An interesting direction for future research would be to investigate the Bahadur efficiency of  $T_n$ . Nikitin (1996a; 1996b) has studied the Bahadur efficiency of various types of supremum-norm statistics in the contexts of testing for symmetry and exponentiality, but it is not clear how to handle statistics of the form  $T_n$ .

## 8. Proofs

We use the following general strategy in each proof. First, we establish the limit distribution for a version of the test statistic in which the range of integration is restricted to a region where the integrand can be approximated uniformly in terms of an empirical process; that region is carefully chosen to avoid a ‘problematic boundary’ where the approximation breaks down. Second, we show that the contribution from the part of the test statistic close to the boundary is asymptotically negligible. The first proof is presented in full detail, but to save space we skip some details in subsequent proofs and concentrate on the key points.

**Proof of Theorem 1.** The problematic boundary is  $\infty$  in this case. For a given  $0 < \varepsilon < 1$ , split the range of integration in the test statistic into the bounded interval  $[0, x_\varepsilon]$  and its complement, where  $F$  has mass  $1 - \varepsilon$  on  $[-x_\varepsilon, x_\varepsilon]$  and mass  $\varepsilon/2$  on each side, by symmetry. Decompose the test statistic as  $T_n = T_{1n} + T_{2n}$ , and note that it suffices to show that, as  $n \rightarrow \infty$ ,

$$T_{1n} = -2 \int_0^{x_\varepsilon} \log R(x) dG_n(x) \xrightarrow{\mathcal{D}} \int_\varepsilon^1 \frac{W^2(t)}{t} dt, \quad (8.1)$$

and

$$T_{2n} = -2 \int_{x_\varepsilon}^\infty \log R(x) dG_n(x) = O_P(\sqrt{\varepsilon}) \quad (8.2)$$

uniformly in  $\varepsilon$ ; see Billingsley (1968, Theorem 4.2).

First consider the leading term  $T_{1n}$ . From (2.1), a Taylor expansion of  $\log(1 + y)$  and the Glivenko–Cantelli theorem it follows that, almost surely,

$$\begin{aligned}
& \sup_{0 < x \leq x_\varepsilon} \left| \log R(x) + \frac{n}{8} \{-F_n(-x) + 1 - F_n(x-)\}^2 \left( \frac{1}{F_n(-x)} + \frac{1}{1 - F_n(x-)} \right) \right| \\
&= \sup_{0 < x \leq x_\varepsilon} \left| nF_n(-x) \log \left( 1 + \frac{1 - F_n(x-) - F_n(-x)}{2F_n(-x)} \right) \right. \\
&\quad \left. + n(1 - F_n(x-)) \log \left( 1 + \frac{F_n(-x) - (1 - F_n(x-))}{2(1 - F_n(x-))} \right) \right. \\
&\quad \left. + \frac{n}{8} \{-F_n(-x) + 1 - F_n(x-)\}^2 \left( \frac{1}{F_n(-x)} + \frac{1}{1 - F_n(x-)} \right) \right| \\
&\leq \sup_{0 < x \leq x_\varepsilon} \left| \frac{n}{24} (1 - F_n(x-) - F_n(-x))^3 \left( \frac{1}{(F_n(-x))^2} - \frac{1}{(1 - F_n(x-))^2} \right) \right| \\
&\leq \frac{1}{24} \sup_{0 < x \leq x_\varepsilon} (\sqrt{n} \{1 - F_n(x-) - (1 - F(x-)) + F(-x) - F_n(-x)\})^2 \\
&\quad \times \sup_{0 < x \leq x_\varepsilon} \left\{ (1 - F_n(x-) - F_n(-x))^2 \frac{F_n(-x) + 1 - F_n(x-)}{(F_n(-x))^2 (1 - F_n(x-))^2} \right\}.
\end{aligned}$$

Now by the weak convergence of the empirical process  $\sqrt{n}(F_n - F)$ , we immediately obtain that this last bound is  $O_P(1) \cdot o_P(1) = o_P(1)$ .

Set  $U_i = F(X_i)$  and let  $\Gamma_n$  be the empirical distribution function of the  $U_i$ . Then by the uniform approximation of  $\log R(x)$  just obtained, we have

$$\begin{aligned}
T_{1n} &= \int_0^{x_\varepsilon} \left( \frac{n}{4} \{-F_n(-x) + 1 - F_n(x-)\}^2 \left( \frac{1}{F_n(-x)} + \frac{1}{1 - F_n(x-)} \right) + o_P(1) \right) dG_n(x) \\
&= \int_0^{x_\varepsilon} \frac{n}{4} \{-\Gamma_n(F(-x)) + 1 - \Gamma_n(F(x-))\}^2 \left( \frac{1}{\Gamma_n(F(-x))} + \frac{1}{1 - \Gamma_n(F(x-))} \right) \\
&\quad d\{\Gamma_n(F(x)) - \Gamma_n(F(-x))\} + o_P(1) \\
&= \int_{\varepsilon/2}^{1/2} \frac{n}{4} \{-\Gamma_n(t) + 1 - \Gamma_n((1-t)-)\}^2 \left( \frac{1}{\Gamma_n(t)} + \frac{1}{1 - \Gamma_n((1-t)-)} \right) \\
&\quad d\{\Gamma_n(t) - \Gamma_n(1-t)\} + o_P(1) \\
&= \frac{1}{4} \int_{\varepsilon/2}^{1/2} \{\sqrt{n}(t - \Gamma_n(t)) + \sqrt{n}((1-t) - \Gamma_n((1-t)-))\}^2 \left( \frac{1}{\Gamma_n(t)} + \frac{1}{1 - \Gamma_n((1-t)-)} \right) \\
&\quad d\{\Gamma_n(t) - \Gamma_n(1-t)\} + o_P(1), \tag{8.3}
\end{aligned}$$

where we have used the change of variable  $t = F(-x)$ . Consider the uniform empirical process

$$\alpha_n(t) = \sqrt{n}(\Gamma_n(t) - t), \quad t \in [0, 1].$$

Since  $\alpha_n$  converges in distribution to a Brownian bridge, the Skorohod construction ensures almost sure convergence in supremum norm of a sequence of uniform empirical processes to a Brownian bridge  $B$ . That is, keeping the same notation for these new uniform empirical processes,

$$\sup_{0 \leq t \leq 1} |\alpha_n(t) - B(t)| \rightarrow 0 \text{ a.s.}$$

The leading term in (8.3) can then be expressed as

$$\frac{1}{2} \int_{\varepsilon/2}^{1/2} \frac{\{-B(t) - B(1-t)\}^2}{t} d\{\Gamma_n(t) - \Gamma_n(1-t)\} + o(1) \text{ a.s.} \tag{8.4}$$

By the Helly–Bray theorem the main expression in (8.4) converges a.s. to

$$\int_{\varepsilon/2}^{1/2} \frac{\{-B(t) - B(1-t)\}^2}{t} dt \stackrel{\mathcal{D}}{=} \int_{\varepsilon/2}^{1/2} \frac{W^2(2t)}{t} dt = \int_{\varepsilon}^1 \frac{W^2(t)}{t} dt.$$

This settles (8.1).

It remains to show that  $T_{2n}$  is asymptotically negligible in the sense of (8.2). Decompose

$$T_{2n} = -2 \int_{x_\varepsilon}^{V_n \vee x_\varepsilon} \log R(x) dG_n(x) - 2 \int_{V_n \vee x_\varepsilon}^{\infty} \log R(x) dG_n(x) = T_{3n} + T_{4n},$$

where  $V_n = \min(-X_{1:n}, X_{n:n})$  and  $X_{i:n}$  denotes the  $i$ th order statistic. Using  $|\log(1+y) - y| \leq 2y^2$  for  $y \geq -\frac{1}{2}$ , we find that

$$|\log R(x)| \leq \frac{n}{2} (-F_n(-x) + 1 - F_n(x-))^2 \left( \frac{1}{F_n(-x)} + \frac{1}{1 - F_n(x-)} \right)$$

for all  $x$ . This leads to (cf. (8.3))

$$\begin{aligned} T_{3n} &\leq \int_{F(-V_n) \wedge \varepsilon/2}^{\varepsilon/2} \{\alpha_n(t) + \alpha_n((1-t)-)\}^2 \left( \frac{1}{\Gamma_n(t)} + \frac{1}{1 - \Gamma_n((1-t)-)} \right) d(\Gamma_n(t) - \Gamma_n(1-t)) \\ &= \int_{F(-V_n) \wedge \varepsilon/2}^{\varepsilon/2} \frac{\{\alpha_n(t) + \alpha_n((1-t)-)\}^2}{t^{1/2}} \frac{1}{t^{1/2}} \left( \frac{t}{\Gamma_n(t)} + \frac{t}{1 - \Gamma_n((1-t)-)} \right) d(\Gamma_n(t) - \Gamma_n(1-t)). \end{aligned} \tag{8.5}$$

The following four sequences are bounded in probability:

$$\begin{aligned} \sup_{0 < t < 1} \frac{|\alpha_n(t)|}{t^{1/4}}, \quad \sup_{0 < t < 1} \frac{|\alpha_n((1-t)-)|}{t^{1/4}}, \\ \sup_{U_{1:n} \leq t \leq 1} \frac{t}{\Gamma_n(t)}, \quad \sup_{1 - U_{n:n} \leq t \leq 1} \frac{t}{1 - \Gamma_n((1-t)-)}, \end{aligned}$$

in the case of the first two by the Chibisov–O’Reilly theorem, and the last two by Shorack

and Wellner (1986, p. 404). Using these bounds inside the integrand of (8.5), and noting that  $F(-V_n) \geq \max(U_{1:n}, 1 - U_{n:n})$ , we obtain

$$T_{3n} = O_P(1) \int_0^{\varepsilon/2} \frac{1}{t^{1/2}} d(\Gamma_n(t) - \Gamma_n(1-t)).$$

It follows from integration by parts that

$$\begin{aligned} \int_0^{\varepsilon/2} \frac{1}{t^{1/2}} d(\Gamma_n(t) - \Gamma_n(1-t)) &= \int_0^{\varepsilon/2} \frac{1}{t^{1/2}} d(\Gamma_n(t) + 1 - \Gamma_n(1-t)) \\ &= \frac{\Gamma_n(\varepsilon/2) + 1 - \Gamma_n(1 - \varepsilon/2)}{(\varepsilon/2)^{1/2}} + \frac{1}{2} \int_0^{\varepsilon/2} \frac{\Gamma_n(t) + 1 - \Gamma_n(1-t)}{t^{3/2}} dt. \end{aligned}$$

Since  $\sup_{0 < t < 1} \Gamma_n(t)/t = O_P(1)$  (see Shorack and Wellner 1986, p. 404), and similarly  $\sup_{0 < t < 1} (1 - \Gamma_n(1-t))/t = O_P(1)$ , we obtain that  $T_{3n} = O_P(\sqrt{\varepsilon})$ .

Finally, consider  $T_{4n}$ . Note that  $R(x)$  is invariant under a sign change of the observations  $X_i$ . Thus it suffices to evaluate  $T_{4n}$  in the case that  $F_n(V_n) = 1$ , which holds either for the original observations or for the sign-changed observations. This gives

$$T_{4n} \leq -2 \int_{V_n}^{\infty} nF_n(-x) \log \frac{1}{2} dG_n(x) = O(n)(1 - G_n(V_n))^2 = O_P(1/n),$$

uniformly in  $\varepsilon$ . The last equality can be seen by noticing that the number of  $|X_i|$  greater than  $V_n$  is bounded above by a geometric random variable with parameter  $\frac{1}{2}$ .  $\square$

**Proof of Theorem 2.** Write  $U_i = F(X_i)$  and let  $\Gamma_{1n}$ ,  $\Gamma_{2n}$  and  $\Gamma_n$  be the corresponding empirical distribution functions. Let  $0 < \varepsilon < \frac{1}{2}$ . It suffices to show that, as  $n \rightarrow \infty$ ,

$$\begin{aligned} T_{1n} &= -2 \int_{\varepsilon}^{1-\varepsilon} \int_{\varepsilon}^{1-\varepsilon} \log R(t, Q(y)) d\Gamma_n(y) dt \\ &\xrightarrow{\mathcal{D}} \int_{\varepsilon}^{1-\varepsilon} \int_{\varepsilon}^{1-\varepsilon} \frac{W_0^2(t, y)}{t(1-t)y(1-y)} dy dt \end{aligned} \quad (8.6)$$

and

$$T_{2n} = T_n - T_{1n} = O_P(\sqrt{\varepsilon}) \quad (8.7)$$

uniformly in  $\varepsilon$ .

First, consider  $T_{1n}$ . By a Taylor expansion it readily follows that uniformly for  $\varepsilon \leq t$ ,  $y \leq 1 - \varepsilon$ ,

$$\begin{aligned} -2 \log R(t, Q(y)) &= nt(1-t)(\Gamma_{1n}(y) - \Gamma_{2n}(y))^2 \left( \frac{1-t}{\Gamma_{1n}(y)(1-\Gamma_{1n}(y))} + \frac{t}{\Gamma_{2n}(y)(1-\Gamma_{2n}(y))} \right) \\ &\quad \times (1 + o(1)) + o_P(1). \end{aligned}$$

So instead of  $T_{1n}$  we consider

$$\int_{\varepsilon}^{1-\varepsilon} \int_{\varepsilon}^{1-\varepsilon} nt(1-t)(\Gamma_{1n}(y) - \Gamma_{2n}(y))^2 \left( \frac{1-t}{\Gamma_{1n}(y)(1-\Gamma_{1n}(y))} + \frac{t}{\Gamma_{2n}(y)(1-\Gamma_{2n}(y))} \right) d\Gamma_n(y) dt.$$

Set  $Y_n(t, y) = \sqrt{n}t(1-t)(\Gamma_{1n}(y) - \Gamma_{2n}(y))$ . From Csörgő and Horváth (1987) (see also McKeague and Sun 1996), it follows that there exists a sequence  $\{W_{0,n}\}$  of four-sided tied-down Wiener processes such that

$$P \left( \sup_{n^{-1/2} < t, y < 1-n^{-1/2}} |Y_n(t, y) - W_{0,n}(t, y)| > A \frac{(\log n)^{3/4}}{n^{1/4}} \right) \leq Bn^{-\delta}$$

for all  $\delta > 0$ , where  $A = A(\delta)$  and  $B$  are constants. Hence it suffices to consider

$$\begin{aligned} \int_{\varepsilon}^{1-\varepsilon} \int_{\varepsilon}^{1-\varepsilon} \frac{W_{0,n}^2(t, y)}{t(1-t)} \left( \frac{1-t}{\Gamma_{1n}(y)(1-\Gamma_{1n}(y))} + \frac{t}{\Gamma_{2n}(y)(1-\Gamma_{2n}(y))} \right) d\Gamma_n(y) dt \\ \stackrel{\mathcal{D}}{=} \int_{\varepsilon}^{1-\varepsilon} \int_{\varepsilon}^{1-\varepsilon} \frac{W_0^2(t, y)}{t(1-t)y(1-y)} d\Gamma_n(y) dt + o_P(1), \end{aligned}$$

which implies (8.6) by the Helly–Bray theorem.

It remains to prove (8.7). We will only consider the relevant region of the unit square where, in addition, both  $y$  and  $t$  are less than or equal to  $\frac{1}{2}$ , that is, we assume  $n^{-1} \leq t \leq \varepsilon$  and  $0 < y \leq \frac{1}{2}$  or,  $n^{-1} \leq t \leq \frac{1}{2}$  and  $0 < y \leq \varepsilon$ . Denote this L-shaped region by  $A_\varepsilon$ . The other regions can be handled in the same way, by symmetry. We prove that

$$\iint_{A_\varepsilon} \log R(t, Q(y)) d\Gamma_n(y) dt = O_P(\sqrt{\varepsilon}). \tag{8.8}$$

We will split the region  $A_\varepsilon$  in turn into several subregions.

First, we consider the case where  $n^{-1} \leq t \leq n^{-3/5}$  and  $n^{-3/8} \leq y \leq \frac{1}{2}$ . Note that in this region

$$\begin{aligned} \left| n_1 \Gamma_{1n}(y) \log \frac{\Gamma_n(y)}{\Gamma_{1n}(y)} \right| &\leq n^{2/5} \log n, \\ \left| n_1 (1 - \Gamma_{1n}(y)) \log \frac{1 - \Gamma_n(y)}{1 - \Gamma_{1n}(y)} \right| &\leq n^{2/5} \log n, \end{aligned}$$

and with arbitrarily high probability, for large  $n$ ,

$$\begin{aligned} \left| n_2 \Gamma_{2n}(y) \log \frac{\Gamma_n(y)}{\Gamma_{2n}(y)} \right| &\leq |2n_2(\Gamma_n(y) - \Gamma_{2n}(y))| \\ &= \left| \frac{2n_2 n_1}{n} (\Gamma_{1n}(y) - \Gamma_{2n}(y)) \right| \leq 2n^{2/5}, \\ \left| n_2 (1 - \Gamma_{2n}(y)) \log \frac{1 - \Gamma_n(y)}{1 - \Gamma_{2n}(y)} \right| &\leq 2n^{2/5}. \end{aligned}$$

Hence with high probability, for large  $n$ ,

$$\int_{n^{-1}}^{n^{-3/5}} \int_{n^{-3/8}}^{1/2} |\log R(t, Q(y))| d\Gamma_n(y) dt \leq \int_{n^{-1}}^{n^{-3/5}} 3n^{2/5} \log n dt \leq \frac{3 \log n}{n^{1/5}} \rightarrow 0.$$

Now consider the region  $n^{-3/8} \leq t \leq \frac{1}{2}$  and  $0 < y \leq n^{-3/5}$ . In this region we have with high probability, for large  $n$ ,

$$\begin{aligned} \left| n_1 \Gamma_{1n}(y) \log \frac{\Gamma_n(y)}{\Gamma_{1n}(y)} \right| &\leq n_1 \Gamma_{1n}(n^{-3/5}) \log n \leq n \Gamma_n(n^{-3/5}) \log n \\ &\leq 2n^{2/5} \log n, \end{aligned}$$

$$\left| n_2 \Gamma_{2n}(y) \log \frac{\Gamma_n(y)}{\Gamma_{2n}(y)} \right| \leq n \Gamma_n(n^{-3/5}) \log n \leq 2n^{2/5} \log n,$$

$$\left| n_1(1 - \Gamma_{1n}(y)) \log \frac{1 - \Gamma_n(y)}{1 - \Gamma_{1n}(y)} \right| \leq |2n_1(\Gamma_{1n}(y) - \Gamma_n(y))| \leq 2n \Gamma_n(n^{-3/5}) \leq 4n^{2/5},$$

$$\left| n_2(1 - \Gamma_{2n}(y)) \log \frac{1 - \Gamma_n(y)}{1 - \Gamma_{2n}(y)} \right| \leq |2n_2(\Gamma_{2n}(y) - \Gamma_n(y))| \leq 2n \Gamma_n(n^{-3/5}) \leq 4n^{2/5}.$$

Hence with high probability, for large  $n$ ,

$$\int_{n^{-3/8}}^{1/2} \int_0^{n^{-3/5}} |\log R(t, Q(y))| d\Gamma_n(y) dt \leq \int_0^{n^{-3/5}} 5n^{2/5} \log n d\Gamma_n(y) \leq \frac{6 \log n}{n^{1/5}} \rightarrow 0.$$

Next consider the region  $n^{-1} \leq t \leq n^{-3/8}$  and  $0 < y \leq n^{-3/8}$ . In this region

$$\left| n_1 \Gamma_{1n}(y) \log \frac{\Gamma_n(y)}{\Gamma_{1n}(y)} \right| \leq n^{5/8} \log(n^{5/8}),$$

$$\left| n_1(1 - \Gamma_{1n}(y)) \log \frac{1 - \Gamma_n(y)}{1 - \Gamma_{1n}(y)} \right| \leq n^{5/8} \log(n^{5/8}),$$

and with high probability, for large  $n$ ,

$$\left| n_2 \Gamma_{2n}(y) \log \frac{\Gamma_n(y)}{\Gamma_{2n}(y)} \right| \leq 2n^{5/8} \log n,$$

$$\left| n_2(1 - \Gamma_{2n}(y)) \log \frac{1 - \Gamma_n(y)}{1 - \Gamma_{2n}(y)} \right| \leq \left| \frac{2n_2 n_1}{n} (\Gamma_{1n}(y) - \Gamma_{2n}(y)) \right| \leq 2n^{5/8}.$$

Hence

$$\int_{n^{-1}}^{n^{-3/8}} \int_0^{n^{-3/8}} |\log R(t, Q(y))| d\Gamma_n(y) dt \leq \frac{4n^{5/8} \log n}{n^{3/4}} \leq \frac{4 \log n}{n^{1/8}} \rightarrow 0.$$

In order to handle the remaining part of  $A_\varepsilon$  we need two lemmas. The first follows quite easily from Inequality 2 of Shorack and Wellner (1986, pp. 415–416).



**Lemma 1.** Let  $0 < a_n, b_n \leq 1/2$  with  $na_nb_n \rightarrow \infty$  as  $n \rightarrow \infty$ . Then, for any  $\delta > 0$ ,

$$P\left(\sup_{a_n \leq t \leq 1} \left\{ \left( \sup_{b_n \leq y \leq 1} \frac{\Gamma_{1n}(y)}{y} \right) \vee \left( \sup_{b_n \leq y \leq 1} \frac{y}{\Gamma_{1n}(y)} \right) \right\} > 1 + \delta \right) \rightarrow 0.$$

The second lemma follows directly from Komlós *et al.* (1975), in a similar but easier way than in Csörgő and Horváth (1987).

**Lemma 2.** Under the same conditions as in Lemma 1, there exists a sequence  $\{W_{0,n}\}$  of four-sided tied-down Wiener processes such that

$$\sup_{a_n \leq t \leq 1-a_n} \sup_{b_n \leq y \leq 1-b_n} \frac{|Y_n(t, y) - W_{0,n}(t, y)|}{(t(1-t)y(1-y))^{1/4}} \xrightarrow{P} 0.$$

We are now in a position to conclude the proof of Theorem 2. Consider the region  $n^{-3/5} \leq t \leq \varepsilon$  and  $n^{-3/8} \leq y \leq \frac{1}{2}$ . We have by a Taylor expansion and Lemma 1 that with high probability, uniformly over this region, for large  $n$ ,

$$\begin{aligned} |\log R(t, Q(y))| &\leq n_1 \frac{(\Gamma_n(y) - \Gamma_{1n}(y))^2}{\Gamma_{1n}(y)(1 - \Gamma_{1n}(y))} + n_2 \frac{(\Gamma_n(y) - \Gamma_{2n}(y))^2}{\Gamma_{2n}(y)(1 - \Gamma_{2n}(y))} \\ &= \frac{n_1 n_2^2 (\Gamma_{1n}(y) - \Gamma_{2n}(y))^2}{n^2 \Gamma_{1n}(y)(1 - \Gamma_{1n}(y))} + \frac{n_2 n_1^2 (\Gamma_{1n}(y) - \Gamma_{2n}(y))^2}{n^2 \Gamma_{2n}(y)(1 - \Gamma_{2n}(y))}. \end{aligned}$$

We only continue with the first term of this sum; the second is somewhat easier to deal with. By Lemma 1, with high probability and uniformly over the region, the first term is bounded above by

$$\frac{2Y_n^2(t, y)}{ty} \frac{y}{\Gamma_{1n}(y)} \leq \frac{3Y_n^2(t, y)}{ty} = 3 \left( \frac{Y_n(t, y)}{(ty)^{1/4}} \right)^2 \frac{1}{(ty)^{1/2}}.$$

But by Lemma 2,

$$\begin{aligned} \int_{n^{-3/5}}^{\varepsilon} \int_{n^{-3/8}}^{1/2} \left( \frac{Y_n(t, y)}{(ty)^{1/4}} \right)^2 \frac{1}{(ty)^{1/2}} d\Gamma_n(y) dt &\stackrel{D}{=} \int_{n^{-3/5}}^{\varepsilon} \int_{n^{-3/8}}^{1/2} \left( \frac{W_0(t, y)}{(ty)^{1/4}} \right)^2 \frac{1}{(ty)^{1/2}} d\Gamma_n(y) dt + o_P(1) \\ &= O_P(1) \int_0^{\varepsilon} \frac{1}{t^{1/2}} dt + o_P(1) = O_P(\sqrt{\varepsilon}). \end{aligned}$$

Finally, it remains to consider the region  $n^{-3/8} \leq t \leq \frac{1}{2}$  and  $n^{-3/5} \leq y \leq \varepsilon$ . This region, however, can be treated in the same way and yields another term of order  $O_P(\sqrt{\varepsilon})$ . Hence (8.7) is proved.  $\square$

**Proof of Theorem 3.** The proof is somewhat similar to the change-point case. Set  $X_i = (X_{i1}, X_{i2})$  and denote the empirical distribution function of the  $(F_1(X_{i1}), F_2(X_{i2}))$  by  $G_n$ , with marginals  $G_{1n}$  and  $G_{2n}$ . Under  $H_0$ , the distribution of  $(F_1(X_{i1}), F_2(X_{i2}))$  is uniform

on the unit square. Write  $Q_1, Q_2$  for the quantile functions corresponding to  $F_1, F_2$ . Let  $0 < \varepsilon < \frac{1}{2}$ . It suffices to show that, as  $n \rightarrow \infty$ ,

$$\begin{aligned} T_{1n} &= -2 \int_{\varepsilon}^{1-\varepsilon} \int_{\varepsilon}^{1-\varepsilon} \log R(Q_1(u), Q_2(v)) dG_{1n}(u) dG_{2n}(v) \\ &\xrightarrow{D} \int_{\varepsilon}^{1-\varepsilon} \int_{\varepsilon}^{1-\varepsilon} \frac{W_0^2(u, v)}{u(1-u)v(1-v)} du dv \end{aligned} \quad (8.9)$$

and

$$T_{2n} = T_n - T_{1n} = O_P(\sqrt{\varepsilon}) \quad (8.10)$$

uniformly in  $\varepsilon$ .

First, consider  $T_{1n}$ . By a Taylor expansion it readily follows that uniformly for  $\varepsilon \leq u, v \leq 1 - \varepsilon$  (replacing  $(x, y)$  by  $(Q_1(u), Q_2(v))$  in the definition of the  $A_{jk}$ ),

$$\begin{aligned} -2 \log R(Q_1(u), Q_2(v)) &= \frac{n(P_n(A_{11})P_n(A_{22}) - P_n(A_{12})P_n(A_{21}))^2}{u(1-u)v(1-v)} + o_P(1) \\ &= \frac{n(P_n(A_{11}) - G_{1n}(u)G_{2n}(v))^2}{u(1-u)v(1-v)} + o_P(1) \\ &= \frac{(\alpha_n(u, v) - v\alpha_{1n}(u) - u\alpha_{2n}(v))^2}{u(1-u)v(1-v)} + o_P(1), \end{aligned} \quad (8.11)$$

with  $\alpha_n(u, v) = \sqrt{n}(G_n(u, v) - uv)$ ,  $\alpha_{1n}(u) = \sqrt{n}(G_{1n}(u) - u)$ ,  $\alpha_{2n}(v) = \sqrt{n}(G_{2n}(v) - v)$ ,  $0 < u, v < 1$ . So instead of  $T_{1n}$  we consider

$$\int_{\varepsilon}^{1-\varepsilon} \int_{\varepsilon}^{1-\varepsilon} \frac{(\alpha_n(u, v) - v\alpha_{1n}(u) - u\alpha_{2n}(v))^2}{u(1-u)v(1-v)} dG_{1n}(u) dG_{2n}(v),$$

which, by standard empirical process theory and a multivariate version of the Helly–Bray theorem, converges in distribution to

$$\int_{\varepsilon}^{1-\varepsilon} \int_{\varepsilon}^{1-\varepsilon} \frac{(B(u, v) - vB(u, 1) - uB(1, v))^2}{u(1-u)v(1-v)} du dv,$$

where  $B$  is a standard bivariate Brownian bridge: a centred Gaussian process with covariance structure  $EB(u, v)B(\tilde{u}, \tilde{v}) = (u \wedge \tilde{u})(v \wedge \tilde{v}) - uv\tilde{u}\tilde{v}$ ,  $0 < u, \tilde{u}, v, \tilde{v} < 1$ . Observing that

$$\{B(u, v) - vB(u, 1) - uB(1, v), (u, v) \in (0, 1)^2\} \stackrel{D}{=} \{W_0(u, v), (u, v) \in (0, 1)^2\},$$

completes the proof of (8.9).

It remains to prove (8.10). We will only consider integration over the region

$$B_{\varepsilon} = \{(u, v) \in (0, 1)^2 : 0 < u \leq \varepsilon, 0 < v \leq \frac{1}{2}, \text{ or } 0 < u \leq \frac{1}{2}, 0 < v \leq \varepsilon\},$$

because of symmetry arguments; compare the way we handled  $A_\varepsilon$  in the change-point case. Because of a further symmetry argument, namely the symmetry in  $u$  and  $v$ , we will further restrict ourselves to the following three regions which clearly cover  $\{(u, v) \in B_\varepsilon : u \leq v\}$ :

$$B_{\varepsilon,1} = \{(u, v) \in (0, 1)^2 : 0 < u \leq n^{-3/5}, n^{-3/8} \leq v \leq \frac{1}{2}\},$$

$$B_{\varepsilon,2} = \{(u, v) \in (0, 1)^2 : 0 < u \leq v \leq n^{-3/8}\},$$

$$B_{\varepsilon,3} = \{(u, v) \in (0, 1)^2 : n^{-3/5} < u \leq \varepsilon, n^{-3/8} \leq v \leq \frac{1}{2}\}.$$

We almost immediately obtain, along the lines of the change-point case,

$$\iint_{B_{\varepsilon,1} \cup B_{\varepsilon,2}} |\log R(Q_1(u), Q_2(v))| dG_{1n}(u) dG_{2n}(v) = o_P(1), \tag{8.12}$$

where we have (again) used the equalities

$$\begin{aligned} |P_n(A_{11}) - G_{1n}(u)G_{2n}(v)| &= |P_n(A_{11}) - G_{1n}(u)(1 - G_{2n}(v))| \\ &= |P_n(A_{21}) - (1 - G_{1n}(u))G_{2n}(v)| \\ &= |P_n(A_{22}) - (1 - G_{1n}(u))(1 - G_{2n}(v))|. \end{aligned}$$

Moreover, here and later in the proof we use the fact that, uniform over certain classes of rectangles (the  $A_{jk}$ ),  $P_n/P$  converges to 1 in probability. This follows from, for example, Chapters 2 and 3 of Einmahl (1987).

For  $(u, v) \in B_{\varepsilon,3}$  it easily follows that with arbitrarily high probability, uniformly over  $B_{\varepsilon,3}$ , for large  $n$ ,

$$\begin{aligned} |\log R(Q_1(u), Q_2(v))| &\leq \frac{(\alpha_n(u, v) - v\alpha_{1n}(u) - u\alpha_{2n}(v))^2}{u(1-u)v(1-v)} \\ &\leq 12 \left\{ \frac{\alpha_n^2(u, v)}{uv} + \frac{\alpha_{1n}^2(u)}{u} + \frac{\alpha_{2n}^2(v)}{v} \right\}; \end{aligned}$$

cf. (8.11). This yields that indeed

$$-2 \iint_{B_{\varepsilon,3}} \log R(Q_1(u), Q_2(v)) dG_{1n}(u) dG_{2n}(v) = O_P(\sqrt{\varepsilon}),$$

and this, in conjunction with (8.12), yields (8.10). □

**Proof of Theorem 4.** The quantile function of  $F$  is  $Q(u) = -\log(1-u)/\lambda$ , so we have

$$T_n = -4 \int_0^1 \int_0^v \log R(Q(u), Q(v)) \left(\frac{\hat{\lambda}}{\lambda}\right)^2 ((1-u)(1-v))^{\hat{\lambda}/\lambda-1} du dv,$$

and it suffices to show that

$$T_{1n} = -4 \int_{\varepsilon}^{1-\varepsilon} \int_{\varepsilon}^v \log R(Q(u), Q(v)) \left(\frac{\hat{\lambda}}{\lambda}\right)^2 ((1-u)(1-v))^{\hat{\lambda}/\lambda-1} du dv$$

$$\xrightarrow{\mathcal{D}} 2 \int_{\varepsilon}^{1-\varepsilon} \int_t^{1-\varepsilon} \frac{st}{(1-s)(1+t)} \left\{ \frac{B(st)}{st} - \frac{B(s)}{s} - \frac{B(t)}{t} \right\}^2 ds dt \quad (8.13)$$

and

$$T_{2n} = T_n - T_{1n} = O_P(\sqrt{\varepsilon}) \quad (8.14)$$

uniformly in  $0 < \varepsilon < \frac{1}{2}$ .

First, consider (8.13). With  $S_n(u) = 1 - F_n(Q(u))$ , by a Taylor expansion

$$-2 \log R(Q(u), Q(v)) = \frac{n(S_n(u)S_n(v) - S_n(u+v-uv))^2}{u(1-u)(1-v)(2-v)} (1 + o_P(1)) \quad (8.15)$$

uniformly for  $\varepsilon \leq u \leq v \leq 1 - \varepsilon$ . Writing

$$S_n(u)S_n(v) - S_n(u+v-uv) = S_n(u)(S_n(v) - (1-v)) + (S_n(u) - (1-u))(1-v)$$

$$+ ((1-u)(1-v) - S_n(1 - (1-u)(1-v)))$$

and using the weak convergence of the uniform empirical process to a standard Brownian bridge  $B$ , we see that the right-hand side of (8.15) converges weakly on  $\varepsilon \leq u \leq v \leq 1 - \varepsilon$  to

$$\frac{((1-u)(-B(v)) - B(u)(1-v) + B(1 - (1-u)(1-v)))^2}{u(1-u)(1-v)(2-v)}$$

$$\stackrel{\mathcal{D}}{=} \frac{(-(1-u)B(1-v) - (1-v)B(1-u) + B((1-u)(1-v)))^2}{u(1-u)(1-v)(2-v)}. \quad (8.16)$$

Thus, using the change of variables  $s = 1 - u$ ,  $t = 1 - v$ , and noting that  $\hat{\lambda} \xrightarrow{P} \lambda$ , we see that (8.13) follows directly from (8.15) and (8.16).

The proof of (8.14) follows along the lines of the previous proofs, in particular the proof of the change-point case. We note here only that results for weighted empirical processes indexed by intervals, especially Theorem 3.3 in Einmahl (1987), are used to complete the proof.  $\square$

## Appendix

Tables A.1–A.4 provide selected critical values for the four proposed test statistics  $T_n$ . The values are based on 100 000 samples in each case.

**Table A.1.** Test for symmetry

$n$	<i>Percentage points</i>			
	90%	95%	97.5%	99%
10	2.620	3.392	4.272	5.393
15	2.477	3.325	4.195	5.317
20	2.428	3.271	4.138	5.306
30	2.360	3.154	3.989	5.160
50	2.295	3.081	3.902	5.027
100	2.254	3.041	3.880	5.005
150	2.231	3.002	3.836	4.967

**Table A.2.** Test for a change point

$n$	<i>Percentage points</i>			
	90%	95%	97.5%	99%
10	1.420	1.667	1.899	2.141
15	1.496	1.756	2.024	2.355
20	1.529	1.804	2.074	2.423
30	1.556	1.832	2.111	2.485

**Table A.3.** Test for independence

$n$	<i>Percentage points</i>			
	90%	95%	97.5%	99%
10	1.535	1.792	2.020	2.283
15	1.572	1.841	2.103	2.442
20	1.575	1.852	2.126	2.485
50	1.581	1.861	2.154	2.553

**Table A.4.** Test for exponentiality

$n$	Percentage points			
	90%	95%	97.5%	99%
10	0.521	0.734	0.969	1.322
15	0.676	0.906	1.148	1.524
20	0.787	1.004	1.242	1.578
30	0.951	1.155	1.370	1.681
60	1.179	1.390	1.611	1.911
120	1.308	1.522	1.747	2.043
300	1.408	1.631	1.855	2.160
$\infty$	1.467	1.679	1.895	2.192

## Acknowledgements

The first named author carried out part of his research at Eindhoven University of Technology. He was supported in part by a Senior Fulbright Scholarship. The second named author's research was supported in part by National Science Foundation grant 9971784.

## References

- Ahmad, I.A. and Alwasel, I.A. (1999) A goodness-of-fit test for exponentiality based on the memoryless property. *J. Roy. Statist. Soc. Ser. B*, **61**, 681–689.
- Ahmad, I.A. and Li, Q. (1997) Testing symmetry of an unknown density function by kernel method. *J. Nonparametr. Statist.*, **7**, 279–293.
- Aly, E.-E.A.A. (1998) Change point tests for randomly censored data. In B. Szyszkowicz (ed.), *Asymptotic Methods in Probability and Statistics*, pp. 503–513. Amsterdam: North-Holland.
- Aly, E.-E.A.A. and Kochar, S.C. (1997) Change point tests based on  $U$ -statistics with applications in reliability. *Metrika*, **45**, 259–269.
- Anderson, T.W. and Darling, D.A. (1952) Asymptotic theory of certain 'goodness-of-fit' criteria based on stochastic processes. *Ann. Math. Statist.*, **23**, 193–212.
- Angus, J.E. (1982) Goodness-of-fit tests for exponentiality based on a loss-of-memory type functional equation. *J. Statist. Plann. Inference*, **6**, 241–251.
- Berk, R.H. and Jones, D.H. (1979) Goodness-of-fit test statistics that dominate the Kolmogorov statistics. *Z. Wahrscheinlichkeitstheorie Verw. Geb.*, **47**, 47–59.
- Billingsley, P. (1968) *Convergence of Probability Measures*. New York: Wiley.
- Butler, C.C. (1969) A test for symmetry using the sample distribution function. *Ann. Math. Statist.*, **40**, 2209–2210.
- Csörgő, M. and Horváth, L. (1987) Nonparametric tests for the changepoint problem. *J. Statist. Plann. Inference*, **17**, 1–9.
- Csörgő, M. and Horváth, L. (1988) Nonparametric methods for changepoint problems. In P.R.

- Krishnaiah and C.R. Rao (eds), *Quality Control and Reliability*, Handbook of Statistics 7, pp. 403–425. Amsterdam: North-Holland.
- Deheuvels, P. (1981) An asymptotic decomposition for multivariate distribution-free tests of independence. *J. Multivariate Anal.*, **11**, 102–113.
- Diks, C. and Tong, H. (1999) A test for symmetries of multivariate probability distributions. *Biometrika*, **86**, 605–614.
- Dykstra, R., Kochar, S. and Robertson, T. (1995) Likelihood ratio tests for symmetry against one-sided alternatives. *Ann. Inst. Statist. Math.*, **47**, 719–730.
- Einmahl, J.H.J. (1987) *Multivariate Empirical Processes*, CWI Tract 32. Amsterdam: Centre for Mathematics and Computer Science.
- Einmahl, J.H.J. and Khmaladze, E.V. (2001) The two-sample problem in  $\mathbb{R}^m$  and measure-valued martingales. In M. de Gunst, C. Klaassen and A. van der Vaart (eds), *State of the Art in Probability and Statistics: Festschrift for Willem R. van Zwet*, IMS Lecture Notes Monogr. Ser. 36, pp. 434–463. Beachwood, OH: Institute of Mathematical Statistics.
- Ferger, D. (1994) Nonparametric change-point tests of the Kolmogorov–Smirnov type. In H.G. Mueller and D. Siegmund (eds), *Change-Point Problems*, IMS Lecture Notes Monogr. Ser. 23, pp. 145–148. Hayward, CA: Institute of Mathematical Statistics.
- Ferger, D. (1995) Nonparametric tests for nonstandard change-point problems. *Ann. Statist.*, **23**, 1848–1861.
- Ferger, D. (1996) On the asymptotic behavior of change-point estimators in case of no change with applications to testing. *Statist. Decisions*, **14**, 137–143.
- Ferger, D. (1998) Testing for the existence of a change-point in a specified time interval. In W. Kahle, E. Von Collani, J. Franz and U. Jensen (eds), *Advances in Stochastic Models for Reliability, Quality and Safety*, pp. 277–289. Boston: Birkhäuser.
- Gombay, E. and Jin, X. (1999) Sign tests for change under alternatives. *J. Nonparametr. Statist.*, **10**, 389–404.
- Hawkins, D.L. (1988) Retrospective and sequential tests for a change in distribution based on Kolmogorov–Smirnov-type statistics. *Sequential Anal.*, **7**, 23–51.
- Hill, D.L. and Rao, P.V. (1977) Tests of symmetry based on Cramér–von Mises statistics. *Biometrika*, **64**, 489–494.
- Komlós, J., Major, P. and Tusnady, G. (1975) An approximation of partial sums of independent RV's and the sample DF. I. *Z. Wahrscheinlichkeitstheorie Verw. Geb.*, **32**, 111–131.
- Li, G. (2000) A nonparametric likelihood ratio goodness-of-fit test for survival data. Preprint.
- Lockhart, R.A. and McLaren, C.G. (1985) Asymptotic points for a test of symmetry about a specified median. *Biometrika*, **72**, 208–210.
- Martynov, G.V. (1992) Statistical tests based on empirical processes, and related problems. *J. Soviet Math.*, **61**, 2195–2271.
- McKeague, I.W. and Sun, Y. (1996) Transformations of Gaussian random fields to Brownian sheet and nonparametric change-point tests. *Statist. Probab. Lett.*, **28**, 311–319.
- Mizushima, T. and Nagao, H. (1998) A test for symmetry based on density estimates. *J. Japan Statist. Soc.*, **28**, 205–225.
- Modarres, R. and Gastwirth, J.L. (1996) A modified runs test for symmetry. *Statist. Probab. Lett.*, **31**, 107–112.
- Nikitin, Y.Y. (1996a) On Baringhaus–Henze test for symmetry: Bahadur efficiency and local optimality for shift alternatives. *Math. Methods Statist.*, **5**, 214–226.
- Nikitin, Y.Y. (1996b) Bahadur efficiency of a test of exponentiality based on a loss-of-memory type functional equation. *J. Nonparametr. Statist.*, **6**, 13–26.

- Orlov, A.I. (1972) Testing the symmetry of a distribution. *Theory Probab. Appl.*, **17**, 372–377.
- Owen, A.B. (2001) *Empirical Likelihood*. Boca Raton, FL: Chapman & Hall/CRC.
- Rothman, E.N.D. and Woodroffe, M.A. (1972) A Cramér–von Mises type statistic for testing symmetry. *Ann. Math. Statist.*, **43**, 2035–2038.
- Shorack, G.R. and Wellner, J.A. (1986) *Empirical Processes with Applications to Statistics*. New York: Wiley.
- Srinivasan, R. and Godio, L.B. (1974) A Cramér–von Mises type statistic for testing symmetry. *Biometrika*, **61**, 196–198.
- Szyszkowicz, B. (1994) Weak convergence of weighted empirical type processes under contiguous and changepoint alternatives. *Stochastic Process. Appl.*, **50**, 281–313.

Received November 2000 and revised March 2002