CRITICAL APPRAISAL SKILLS PROGRAMME making sense of evidence

11 questions to help you make sense of a Case Control Study

General comments

• Three broad issues need to be considered when appraising a case control study.

Are the results of the study valid?

What are the results?

Will the results help locally?

The 11 questions on the following pages are designed to help you think about these issues systematically.

- The first two questions are screening questions and can be answered quickly. If the answer to those two is "yes", it is worth proceeding with the remaining questions.
- There is a fair degree of overlap between several of the questions.
- You are asked to record a "yes", "no" or "can't tell" to most of the questions.
- A number of italicised hints are given after each question. These are designed to remind you why the question is important. There will not be time in the small groups to answer them all in detail!

[©] Critical Appraisal Skills Programme (CASP) 2004. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise without the prior permission of CASP. However, organisations may reproduce or use the publication for non-commercial educational purposes provided the source is acknowledged. Enquiries concerning reproduction or use in other circumstances should be addressed to CASP.

A/ Are the results of the study valid?

Screening Questions

 1 Did the study address a clearly focused issue? HINT: A question can be focused in terms of: the population studied the risk factors studied whether the study tried to detect a beneficial or harmful effect? 	Yes	Can't tell	No D
2 Did the authors use an appropriate method to answer their question?	Yes	Can't tell	No L
HINT: Consider			
 Is a case control study an appropriate way of answering the question under the circumstances? (Is the outcome rare or harmful?) Did it address the study question? 			

Is it worth continuing?

Detailed Questions

3 Were the cases recruited in an acceptable way?	Yes	Can't tell	No
 HINT: We are looking for selection bias which might compromise the validity of the findings: Are the cases defined precisely? Were the cases representative of a defined population (geographically and/or temporally)? Was there an established reliable <u>system</u> for selecting all the cases? Are they incident or prevalent? Is there something special about the cases? Is the time frame of the study relevant to the disease/exposure? Was there a sufficient number of cases selected? Was there a power calculation? 			

Yes	Can't tell	No
Yes	Can't tell	No
List:		
Yes	Can't tell	No L
	Yes List:	Yes Can't tell Output Output

B/ What are the results?

7. What are the results of this study?	
HINT:	
 What are the bottom line results? Is the analysis appropriate to the design? How strong is the association between exposure and outcome (look at the odds ratio)? Are the results adjusted for confounding and might confounding still explain the association? Has adjustment made a big difference to the OR ?? 	
8. How precise are the results?	
How precise is the estimate of risk?	
 Size of the P-value Size of the confidence intervals Have the authors considered all the important variables? How was the effect of subjects refusing to participate evaluated? 	
 9. Do you believe the results? HINT: Big effect is hard to ignore! Can it be due to chance, bias or confounding? Are the design and methods of this study sufficiently flawed to make the results unreliable? Consider Bradford Hills criteria (e.g. time sequence, dose-response gradient, strength, biological 	Yes No
plausibility)	

Is it worth continuing?

C/ Will the results help me locally?

10. Can the results be applied to the local population?	Yes	Can't tell	No
HINT: Consider whether			
 The subjects covered in the study could be sufficiently different from your population to cause concern. Your local setting is likely to differ much from that of the study. Can you estimate the local benefits and harms? 			
11. Do the results of this study fit with other available evidence?	Yes	Can't tell	No
HINT:			
- Consider all the available evidence from RCTs, systematic reviews, cohort studies and case-control studies as well for consistency.			

One observational study rarely provides sufficiently robust evidence to recommend changes to clinical practice or within health policy decision making.

However, for certain questions observational studies provide the only evidence.

Recommendations from observational studies are always stronger when supported by other evidence. CASP Copyright: Critical Appraisal Skills Programme (CASP) 2004. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise without the prior permission of CASP. However, organisations may reproduce or use the publication for non-commercial educational purposes provided the source is acknowledged. Enquiries concerning reproduction or use in other circumstances should be addressed to CASP.

http://www.phru.nhs.uk/casp/cohort%2012%20questions.pdf