

Problem Set #14

*Don't miss the fourth problem on the second page**Problem 47 has been divided into two problems to properly weight its grade. Minor correction to #47.*

46. Consider a quantum mechanical system with a Hamiltonian H at a temperature T described by a density matrix ρ . The eigenstates and eigenvalues of H are $|\phi_n\rangle$ and E_n . (Since n labels the states, there may be many values of n whose values for E_n are the same.)
- Write down an explicit expression for ρ in terms of $|\phi_n\rangle$, E_n .
 - Write down an explicit expression for the von Neumann entropy σ in terms E_n .
 - If the Helmholtz free energy F is defined as $F = E - kT\sigma$ where E is the total energy of the system, write down an explicit expression for F in terms of E_n .
47. A quantum system is made up of the product of two spin- $\frac{1}{2}$ states: $|\pm\rangle_A \otimes |\pm\rangle_B$, where the factor $|\pm\rangle_{A/B}$ describes a state with $s_z = \pm\frac{\hbar}{2}$. Let $\vec{s}^A$ and $\vec{s}^B$ be the corresponding spin operators and assume a Hamiltonian of the form:

$$H = \frac{\alpha kT}{\hbar} S_z + \frac{\beta kT}{2} \left(\frac{\vec{S}^2}{\hbar^2} + \frac{3}{2} \right)$$

where $\vec{S} = \vec{s}^A + \vec{s}^B$ and the factors of $\frac{1}{2}$ and kT and the constant $\frac{3}{2}$ have been introduced to give simple expressions for the resulting Boltzmann factors.

- If this system is in a mixed state at the temperature T write down the 4×4 density matrix $\rho^{A \otimes B}$ describing that system. (This can be expressed as a sum over four projection operators with explicit coefficients.)
 - What is the average result if the total spin $\vec{S}$ is measured many times on such a system?
 - Find the reduced 2×2 density matrix ρ^A that can be used to calculate the results from measuring operators O^A which act on only the states $|\pm\rangle^A$ that appear in the basis states $|\pm\rangle_A \otimes |\pm\rangle_B$.
 - If only the spin $\vec{s}^A$ is measured multiple times, use your result from part (c) to determine the average result? Does $\langle \vec{S}^{A \otimes B} \rangle = 2\langle \vec{s}^A \rangle$?
48. Use the results of problem 47 to answer the following questions.
- If subsystem associated with the states $|\pm\rangle$ is interpreted as a system with the Hamiltonian $H = \frac{\alpha}{\hbar} s_z^A$ what temperature would you assign to the system?
 - What is the von Neumann entropy $\sigma^{A \otimes B}$ of the system described by the density matrix $\rho^{A \otimes B}$ obtained in part (a)?
 - What is the von Neumann entropy σ^A of the system described by the reduced density matrix ρ^A obtained in part (c)? What is the relation between the magnitudes of the von Neumann entropies $\sigma^{A \otimes B}$ and sum of the entropies of the subsystems A and B which because of symmetry equals $2\sigma^A$?

49. (From Gottfried and Yan, Ch 2, problem 4) Consider a system S composed of two subsystems A and B and let $\rho^{A \otimes B}$ be the density matrix for a configuration of the combined system S . Show that for the reduced density matrix ρ^A for subsystem A to describe a pure state then the density matrix $\rho^{A \otimes B}$ must have the form:

$$\rho^{A \otimes B} = |\phi\rangle_A \langle\phi|_A \otimes \rho^B$$

where $|\phi\rangle_A$ is a normalized state in the Hilbert space for the system A and ρ^B is the reduced density matrix for the subsystem B .