

Problem Set #17

58. Show explicitly that $P_d(t) + \int d^3p P_{\vec{p}}(t) = 1$ in the Wigner-Weisskopf description of the decay of a discrete state $|d\rangle$ into a continuum of states $|\phi(\vec{p})\rangle$. The quantity $P_d(t)$ is the probability of finding the system in the state $|d\rangle$ at time t while $P_{\vec{p}}(t)$ is the probability density corresponding to the states $|\phi(\vec{p})\rangle$. Assume that at $t = 0$, $P_d(0) = 1$ and $P_{\vec{p}}(0) = 0$.
59. A negatively charged muon with mass $m_\mu = 106 \text{ MeV}/c^2$ and an electron with mass $m_e = 0.51 \text{ MeV}/c^2$ orbit a helium nucleus. Initially the muon is in the $2p$, $m_l = 0$, state while the electron is in the $1s$ state. Compute the rate for the decay process in which the muon falls to the $1s$ state while the electron is ejected from the atom. Since $m_\mu \gg m_e$, the Helium nucleus and the muon occupy a very small volume and appear to the electron as an accurate approximation to a hydrogen nucleus. Call this the central charge approximation.
- Treat the deviation between the charge distribution of the muon and this central charge approximation as creating a perturbing electrostatic potential $V(\vec{r})$ which acts on the electron. Write down an expression for the decay rate in terms of a matrix element of $V(\vec{r})$ between the initial and final states where the final state is the product of the $1s$ muon and a plane wave for the ejected electron.
 - Determine this matrix element by first evaluating the factor in the matrix element which involves the muon wave function.
 - Second evaluate the portion of matrix element involving the electron's coordinates.
 - Finally evaluate the decay rate by integrating over the direction of the emitted electron.
60. Consider two degenerate states $|d_1\rangle$ and $|d_2\rangle$ which are coupled to a continuum of states $|\phi_{\vec{p}}\rangle$ by a potential V :

$$\langle \phi_{\vec{p}} | V | d_i \rangle = V_{\vec{p},i}$$

where $H = H_0 + V$ with

$$\begin{aligned} H_0 |d_i\rangle &= E_d |d_i\rangle \quad i = 1, 2 \\ H_0 |\phi_{\vec{p}}\rangle &= E_{\vec{p}} |\phi_{\vec{p}}\rangle \end{aligned}$$

and we assume that $\langle d_i | V | d_j \rangle = \langle \phi_{\vec{p}} | V | \phi_{\vec{p}'} \rangle = 0$. Show using the Wigner-Weisskopf method that the state $\beta_1 |d_1\rangle + \beta_2 |d_2\rangle$ decays with the exponent $-\gamma t/2$ provided the quantities β_1 , β_2 and γ satisfy the following two conditions:

- (a) The column vector $\begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix}$ is an eigenvector of the 2×2 matrix

$$\begin{aligned} \Gamma_{i,j} &= \frac{2\pi}{\hbar} \int d^3p \delta(E_{\vec{p}} - E_d) \langle d_i | V | \phi_{\vec{p}} \rangle \langle \phi_{\vec{p}} | V | d_j \rangle \\ &\quad + \frac{2i}{\hbar} \mathcal{P} \left\{ \int d^3p \frac{\langle d_i | V | \phi_{\vec{p}} \rangle \langle \phi_{\vec{p}} | V | d_j \rangle}{E_d - E_{\vec{p}}} \right\} \end{aligned}$$

- (b) The quantity γ is the corresponding complex eigenvalue.