

## Assignment #4

## Reading:

Oct 5 Kleppner and Kolenkow Note 5.2

Oct 7 Kleppner and Kolenkow 6.5

## Problems:

26. Kleppner and Kolenkow 4.15 (Assume that the relative velocity  $u$  between a jumping man and the railway car is the difference between the man's and car's velocities after the jump.)
27. Kleppner and Kolenkow 5.3
28. Kleppner and Kolenkow 5.4
29. Kleppner and Kolenkow 5.6
30. Kleppner and Kolenkow 5.15
31. Kleppner and Kolenkow 5.19 (Note "mechanical energy" means kinetic and potential.)
32. (Kleppner and Kolenkow 1st edition) Find the forces implied by the following potential energies:
  - (a)  $U = Ax^2 + By^2 + Cz^2$
  - (b)  $U = A \ln(x^2 + y^2 + z^2)$  ( $\ln = \log_e$ )
  - (c)  $U = A \cos(\theta)/r^2$  (Here  $r, \theta$  are plane polar coordinates.)
33. Consider the effect of a radio wave on an ionospheric electron discussed in Example 1.11 in Kleppner and Kolenkow. Use CGS units. To avoid unnecessary complications assume that the quantity  $eE_0/m = 1 \text{ cm/sec}^2$  and that the frequency of the radio wave is  $\omega = 3/\text{sec}$ .
  - (a) If the electron is at rest at  $t = 0$  evaluate the explicit solution to determine how far the electron has moved from its position at the time  $t = 2\pi \text{ sec}$ .
  - (b) Using Python integrate Newton's equation using the simple method of updating the position and velocity from their values at the time  $t = n\Delta t$  to the time  $t = (n + 1)\Delta t$  using the velocity and acceleration at the time  $t = n\Delta t$  described in class. If a time step  $\Delta t = 0.1$  is used, how large a numerical error appears in the solution at the time  $t = 2\pi \text{ sec}$ ? (Simply compare the exact and numerical solutions at  $t = 2\pi \text{ sec}$ .) Show a graph of your Python solution for the time interval 0 to 10 sec.

- (c) Show using Python that the “kick” given the electron when the electric field is turned on at  $t=0$  can be reduced if the oscillating field is slowly turned starting at  $t = -8$  sec using:

$$E(t) = \frac{m}{e} \cdot \frac{1 \text{ cm}}{\text{sec}^2} \begin{cases} 0 & t \leq -8 \text{ sec} \\ \frac{10^{0.5t/\text{sec}}}{0.01+10^{0.5t/\text{sec}}} \sin(\omega t) & -8 \text{ sec} \leq t \leq 0 \\ \sin(\omega t) & 0 \leq t \end{cases} .$$

Do this by comparing the electron’s position at  $t = 2\pi$  sec found for the original problem with the electron’s position at  $t = 2\pi$  sec for this second choice of  $E(t)$  assuming that for this 2<sup>nd</sup> case  $x$  and  $dx/dt$  vanish at  $t = -8$  sec. Show a graph of your Python solution for the time interval -8 to 10 sec.

The Python code needed to solve this problem might be developed by modifying the Jupyter notebook [ProjectileMotion](#) discussed in class.