


Answer each of the following **four (4)** questions. Please give a complete description of your method of solution since *partial credit* will be given.

1. A cylinder of length  $l$  and radius  $R$  carries a uniform current density  $j$ .




- (a) Find the magnetic field for both the regions  $r > R$  and  $r < R$ . [10 points]

- (b) A particle with momentum  $\vec{p}$  and charge  $q$  moves parallel to and within the cylinder, displaced a distance  $r$  from the cylinder's axis. Calculate the additional momentum  $\Delta\vec{p}$  given the particle as it passes through the length  $l$  of the cylinder. (You should assume that  $\Delta p/p \ll 1$  so that the particle's spatial deflection by the magnetic forces can be neglected.) [10 points]


- (c) Including the effects of this  $\Delta\vec{p}$  on the subsequent motion of the particle, show that such particles for all values of  $r$ ,  $0 \leq r \leq R$ , will cross the axis of the cylinder at the same distance  $D$  from the cylinder. Find this focal length  $D$ . [5 points]

2. Consider the combination of two resistors,  $R_1$  and  $R_2$  and a capacitor  $C$  driven by an applied voltage  $V(t) = V_0 \cos(\omega t)$  configured as shown at the right.




- (a) Find the complex impedance  $Z$  seen between the points A and C. [9 points]  
 (b) What is the actual current flowing through  $R_1$  in both amplitude and phase? [8 points]  
 (c) What is the average power dissipated in the resistor  $R_2$ ? [8 points]

3. (a) A mono-energetic, horizontal beam of electrons, each with an energy of 1 electron Volt, is incident on a vertical screen with a horizontal slit of width  $D = 10^{-5}$  cm. Make a rough numerical estimate of the vertical width of the pattern of electrons that strike a vertical screen a distance of 1 meter from the screen. Recall that an electron has mass  $m = 0.911 \cdot 10^{-27}$  gram,  $\hbar = 1.05 \cdot 10^{-27}$  erg sec and  $1 \text{ eV} = 1.602 \cdot 10^{-12}$  erg. [9 points]



- (b) Use the uncertainty principle,  $\Delta r \Delta p \geq \hbar/2$  to show that the orbit of an electron bound to a proton must have a minimum size, recognizing that the orbiting electron will have energy equal to  $p^2/2m - e^2/r$ . Estimate the size of the hydrogen atom in terms of the electron's mass ( $m$ ), charge ( $e$ ) and Planck's constant  $\hbar$  [8 points]

- (c) A particle of mass  $m$  and charge  $q$  is constrained to move in a circle of radius  $R$ . The particle is located by giving its position  $x$  measured along the circumference. If a uniform magnetic field  $B$  is imposed perpendicular to the plane of the circle, the energy operator can be written:



$$H_{\text{op}} = \frac{(p_{\text{op}} - eRB/c)^2}{2m}.$$

Find the allowed energies and the wave functions of the corresponding energy eigenstates for this system. [8 points]

4. A particle of mass  $m$  moves within a very deep square well  $V(x)$  given by

$$V(x) = \begin{cases} V_0 & x \leq 0 \text{ or } L \leq x \\ 0 & 0 \leq x \leq L \end{cases} \quad (1)$$

where  $V_0$  is a very large positive constant.

- (a) Write down the energy operator (Hamiltonian) for this system and determine those allowed energies which are much smaller than  $V_0$  and the corresponding eigenstates. [8 points]
- (b) Consider a state which at  $t = 0$  has the wave function

$$\psi(x, 0) = N \sin\left(\frac{2\pi}{L}x\right) \cos\left(\frac{\pi}{L}x\right). \quad (2)$$

If the energy of this state is measured, what results are possible? With what probabilities will those results be found? [6 points]

- (c) Find the normalization constant  $N$  needed so that the state  $\psi(x, 0)$  has unit probability. [4 points]
- (d) Find the state  $\psi(x, t)$  into which the state above evolves at the time  $t$ . [4 points]
- (e) Determine the smallest non-zero value of  $t$  for which  $\psi(x, t)$  is physically equivalent to  $\psi(x, 0)$ . [3 points]