

(*Science's Compass*, 20 Nov., p. 1420) about the use of Cre-loxP mouse technology for biomedical research reveals many analogies with plant biotechnology. Proprietary research tools such as promoters and transformation systems have found popular use among the global plant biotechnology community in the last decade or more, and many research projects that used these tools are now in a position for commercial exploitation. Academic institutions such as Michigan State University are now exploring the options open to us and approaching the patent holders of some of these technologies to determine how we can proceed to commercialization. To our surprise and dismay, the initial response has been very different from the Cre-loxP agreement described by Block and Curran and has resulted in a scenario where transgenic plants developed with obvious commercial value are effectively vetoed by the patent holders of these "upstream" technologies.

This is an undesirable situation for agricultural biotechnology (in particular, of transgenic plants), in that the holders of these proprietary "upstream" technologies have effective veto power over whom universities can and cannot approach with their technologies for commercial development.

By inhibiting fair competition and innovation, the development of this sector may well be stifled by a select number of companies holding key basic research tool patents. So while Block and Curran present a favorable picture for the mouse in the laboratory, the situation for maize in the field looks very different to us at present.

Colm Lawler

Licensing Assistant, Office of Intellectual Property, Michigan State University, East Lansing, Michigan 48824, USA.

Fred Erbsich

Director, Office of Intellectual Property, Michigan State University

Replacing Ancient Forests

Anne Simon Moffat's article "Temperate forests gain Moffat" (*News Focus*, 13 Nov., p. 1253) might more accurately have been titled "Industrial forests gain, ancient forests and biodiversity continue to lose." Conservationists welcome reforestation in North America, but the working forests of industry or the mongrel successional forests of the suburbs and abandoned farms are not everywhere a fair trade for our native old-growth forests. The continued ecological losses that attend the destruction of bottom-land hardwood forests of the Southeast (1), the native oak woodlands of California (2), or the native temperate rain forests of the Pacific Northwest (3) are hardly rectified

by the proliferation of genetically altered loblolly pine, exotic eucalyptus, or plantation Douglas-fir. The silvacultural trends described by Wernick *et al.* (4) are welcome not only if they can provide timber and fiber or sequester carbon but also if they can help stop the bleeding in our final few ancient forests.

David W. Stahle

Tree-Ring Laboratory, University of Arkansas, Fayetteville, AR 72701, USA. E-mail: dstahle@comp.uark.edu


References

1. *An Ecosystem in Crisis: The Forested Wetlands of the Mississippi River* (The Nature Conservancy of Louisiana, Baton Rouge, LA, 1992).
2. B. M. Pavlik, P. C. Muick, S. Johnson, M. Popper, *Oaks of California* (Cachuma, Los Olivos, CA, 1991).
3. K. Ervin, *Fragile Majesty: The Battle for North America's Last Great Forest* (Mountaineers, Seattle, WA, 1989).
4. I. Wernick *et al.*, *J. Indust. Ecol.* 1, 125 (1997).

Hominid Brain Volume

Having calculated the brain volumes of several australopithecine and early Homo fossil hominid brain endocasts (1–3), I read with considerable interest the report by Glen C. Conroy *et al.* (12 June, p. 1730) and the commentary by Dean Falk (Perspectives, *Science's Compass*, 12 June, p. 1714). Reexamination of these older specimens by other scientists is a welcome enterprise and, needless to say, I hope that my early attempts will be validated. However, it is important to note that my earlier volume estimations were, in fact, significantly smaller than those previously published. The Sts 71 specimen, for which I obtained a value of 428 cubic centimeters (cm^3), had been estimated as somewhere between 480 and 520 cm^3 . My estimate of the Taung child was 404 cm^3 (4), a drop from Raymond Dart's earlier value of 525 to 562 cm^3 .

The following facts should be noted by readers. First, Conroy *et al.*'s citation of my 1983 article (5) is rather late. The original volumes were published in 1970 (1), again in 1972 (2), with specific discussion of Sts 71, and again in 1973 (3). Second, as I pointed out in the 1972 article in particular (2), the Sts 71 cranium was distorted in the occipital region, and the volume I determined was based on correcting the original endocast. I also graded my attempts according to methods used and found Sts 71 to have the lowest rating (C2–3). Neither Conroy *et al.* nor Falk mentions the plastic deformation that causes the planum occipitale to be at right angles to the endoclast, where the mastoid process is practically at the same plane as the occipital planum, a condition I have seen only on this cranium. Third, pouring one-half of 370 cm^3 of water into a cast of Sts 71 without correcting for the distortions



Cranium of Stw 505, showing "virtual endocast"

and shrinkage is, mildly put, without scientific rigor. In 1970 (4), I wrote, "The standard deviation and coefficient of variation I calculated for the gracile forms are possibly too low, and can be attributed to the small sample size and bias created by using certain gracile values and dimensions to reconstruct less complete specimens." Fourth, those who have access to the casts of Sts 5 and Sts 71 will find that the facial measurements (undistorted) of the two crania are nearly identical, while Sts 5 has a cranial volume of 480 cm^3 ; I know of no evidence disputing that figure. It seems highly unlikely that its cranial volume will be some 110 cm^3 more than that for Sts 71.

I look forward to the use of better technology to pursue these difficult reconstructions, but hope that the attempts to do so will be truly scientific.

Ralph L. Holloway

Department of Anthropology, Columbia University, New York, NY 10027, USA. E-Mail: lh2@columbia.edu

References

1. R. L. Holloway, *Nature* 227, 199 (1970).
2. —, in *Functional and Evolutionary Biology of Primates*, R. Tuttle, Ed. (Aldine, Chicago, 1972), p. 185.
3. —, *J. Hum. Evol.* 2, 449 (1973).
4. —, *Science* 168, 966 (1970).
5. —, *Can. J. Anthropol.* 3, 215 (1983).

Response

Holloway has made many important contributions to paleoneurology, and we are therefore pleased that his comment finds our work to be of "considerable interest." He correctly reminds readers that he was one of the first to realize that many of the early endocranial estimates were overestimates, a situation he corrected in a series of important studies, many of which he cites in his comment. Because Holloway reserves his more specific comments for Sts 71, a specimen not particularly germane to the main focus of our report,

namely, determination of endocranial capacity in Stw 505, we will leave further discussion of that interesting specimen for another day. We can report, however, that some of our more recent computed tomography studies have reconfirmed one endocranial capacity estimate mentioned by Holloway, that of 480 cm³ for Sts 5 (Mrs. Ples).

We welcome the input of our colleagues into this most interesting debate about early hominid paleobiology. Once paleoanthropology matures as a science to the point where all interested parties have unfettered access to the same original fossil data, we suspect that many of our apparent differences of interpretation will melt away.

Glenn C. Conroy

Department of Anatomy and Neurobiology and of Anthropology, Washington University School of Medicine, St. Louis, MO 63110, USA. E-mail: conroy@thalamus.wustl.edu

Gerhard W. Weber

Horst Seidler

Institute of Human Biology, University of Vienna, A-1090 Vienna, Austria

Phillip V. Tobias

Department of Anatomical Sciences, University of the Witwatersrand Medical School, Johannesburg, South Africa

CORRECTIONS AND CLARIFICATIONS


In the issue of 11 December, in the article "Genome sequence of the nematode *C. elegans*: A platform for investigating biology" by The *C. elegans* Sequencing Consortium (Special Section, "*C. elegans*: Sequence to biology," p. 2012), figure 3 (p. 2016) was printed incorrectly (see <http://www.sciencemag.org/feature/data/5396-2012.pdf> for correct figure). In the same issue, in the article "*Caenorhabditis elegans* is a nematode" by Mark Baxter (Special Section, "*C. elegans*: Sequence to biology," p. 2041), figure 2 (p. 2043) was printed incorrectly (see <http://www.sciencemag.org/feature/data/5396-2041.pdf> for correct figure).

In the issue of 4 December, in the Research Article "X-ray crystal structure of the Fe-only hydrogenase (Cpl) from *Clostridium pasteurianum* to 1.8 angstrom resolution" by J. W. Peters *et al.* (p. 1853), the top of figure 1A (p. 1854) was cut off (the correct figure can be seen at <http://www.sciencemag.org/feature/data/5395-1853.pdf>). In the same issue, in the report "Oxygen isotope exchange between refractory inclusion in Allende and solar nebula gas" by H. Yurimoto *et al.* (p. 1874), figure 1 was not printed in half tones (see correct figure at <http://www.sciencemag.org/feature/data/5395-1874.pdf>). In the same issue, in the report "Single-molecule enzymatic dynamics" by H. P. Lu *et al.* (p. 1877), two lines of text were covered by figure 1A (p. 1878). Those lines should have read, "the gel. With excess amounts of cholesterol (0.2...." And, again, in the same issue, in the report "Coupling of mitosis to the completion of S phase through Cdc34-mediated degradation of Wee1" by W. M. Michael and J. Newport (p. 1886), the top three parts of figure 1A (p. 1887) and all of figure 1C were missing. And parts of figure 3 (A, B, and C) (p. 1888) were missing (the correct figure can be seen at <http://www.sciencemag.org/feature/data/5395-1886.pdf>).

In the Policy Forum "The science and technology-bereft Department of State" by Anne Keatley Solomon (*Science's Compass*, 27 Nov., p. 1649), the last paragraph in the third column of page 1649 (carrying over onto page 1650) should have read, "Finally, basic S&T literacy for all State Department personnel is fundamental. Department leadership must make clear the relevance of this basic knowledge to Foreign Service officers. A clear signal early would be the inclusion in the entrance examination of questions testing a basic understanding of fundamental scientific concepts and the nature of scientific inquiry."


In the Policy Forum "The science and technology-bereft Department of State" by Anne Keatley Solomon (*Science's Compass*, 27 Nov., p. 1649), the last paragraph in the third column of page 1649 (carrying over onto page 1650) should have read, "Finally, basic S&T literacy for all State Department personnel is fundamental. Department leadership must make clear the relevance of this basic knowledge to Foreign Service officers. A clear signal early would be the inclusion in the entrance examination of questions testing a basic understanding of fundamental scientific concepts and the nature of scientific inquiry."

LOOK AT THESE IMAGES.



NOW, LOOK AT THE PRICE: \$1195.*

THE 1-MILLION-PIXEL DIGITAL CAMERA SYSTEM
No matter what your photomicroscopy application, you cannot get better image quality at a lower price than with a Pixera digital camera system. In fact, to match Pixera's million-pixel resolution, you would have to spend at least 5 times as much. And Pixera gives you everything you need including a full suite of application software so you can visualize, create and communicate.



NOW—THE PERFECT TOOL FOR TELEMEDICINE.
You can capture two- or three-dimensional still or moving images directly to your PC, Mac, or laptop and then manipulate, organize, archive, and communicate those images via video conferencing or e-mail. A new PCI card makes the system unbelievably easy to install and to use. For more information, visit our web site at www.pixera.com or call toll-free 1-888-4 PIXERA.

pixera

*US list price

All images were captured at 1280 x 1024 pixels with a Pixera Professional Digital Camera mounted on a microscope with a standard C-mount adapter.
140 Knowles Drive, Los Gatos, CA 95030 www.pixera.com ©1998 Pixera Corporation.

CALL FOR SYMPOSIUM PROPOSALS

AAAS Annual Meeting and Science Innovation Exposition

February 17-22, 2000 • Washington, D.C.

You are invited to propose a symposium for the AAAS annual meeting.

Symposium Proposal forms, meeting theme track, and additional information available at:

www.aaas.org/meetings/scope

AAAS Meetings Office
1200 New York Avenue, NW
Washington, DC 20005
Phone: 202-326-6450
Fax: 202-289-4021
E-mail: confinfo@aaas.org

March 22, 1999

Deadline for AAAS to receive proposals.



AMERICAN ASSOCIATION FOR THE
ADVANCEMENT OF SCIENCE

Circle No. 19 on Readers' Service Card