

Robust utility maximization with intractable claims

Yunhong Li*

Zuo Quan Xu[†]

Xun Yu Zhou[‡]

April 17, 2023

Abstract

We study a continuous-time expected utility maximization problem in which the investor at maturity receives the value of a contingent claim in addition to the investment payoff from the financial market. The investor knows nothing about the claim other than its probability distribution, hence an “intractable claim”. In view of the lack of necessary information about the claim, we consider a robust formulation to maximize her utility in the worst scenario. We apply the quantile formulation to solve the problem, expressing the quantile function of the optimal terminal investment income as the solution of certain variational inequalities of ordinary differential equations. In the case of an exponential utility, the problem reduces to a (non-robust) rank-dependent utility maximization with probability distortion whose solution is available in the literature.

Keywords: Intractable claim; robust model; quantile formulation; calculus of variations; variational inequalities; rank-dependent utility.

Mathematics Subject Classification (2020) 91B28; 91G10; 35Q91

1 Introduction

The expected utility models have taken a central position in modern financial portfolio selection theory. In a typical expected utility model, an investor looks for the best portfolio in a given financial market to maximize her utility, based on a dynamically expanding body of information and subject to various constraints. There are also the so-called partial information models in which not all the information are available to the investor and she needs to continuously observe some related information processes to

*Department of Applied Mathematics, The Hong Kong Polytechnic University, Kowloon, Hong Kong, China. Email: yunhong.li@connect.polyu.hk.

[†]Department of Applied Mathematics, The Hong Kong Polytechnic University, Kowloon, Hong Kong, China. Email: maxu@polyu.edu.hk.

[‡]Department of Industrial Engineering and Operations Research & The Data Science Institute, Columbia University, New York, NY 10027, USA. Email: xz2574@columbia.edu.

gain better understanding of the underlying market over time, using techniques such as filtering theory and adaptive controls.

In practice, however, there are situations where people, facing future contingent claims (incomes or liabilities) that cannot be understood even if the time is very close to maturity, aim to maximize the utility of a total sum including payoffs from the stock investment as well as those “external” claims. For instance, the claim of an insurance contract is not revealed until the related event occurs, the down payment of a mortgage is only known when the purchase has been made, and the value of the employer’s stock option may have little to do with the general stock market. These expected utility models are essentially different from the typical partial information models mentioned above because, while the probability distributions of these claims can be estimated, little or no information on their *correlations* with the financial market are available. Such a claim, about which we know nothing but its own distribution, is termed an “intractable claim” in Hou and Xu [7].

Because intractable claims are not predictable nor hedgeable until they are realized, classical stochastic control including filtering theory is not applicable. Hou and Xu [7] formulate and investigate a continuous-time Markowitz’s mean–variance model with an intractable claim and the objective to minimize the robust variance in the worst scenario. The special structure of the mean–variance setting allows them to use the completion-of-squares and quantile optimization techniques to derive the optimal solution.

In this paper we consider a continuous-time expected utility model (i.e. Merton’s problem) with an intractable claim. The usual maximization of expected utility is not even well defined because one cannot compute the expected utility of the sum of the investment payoff from the market and the intractable claim without knowing the joint distribution of the two. Instead, we introduce a robust model whose objective is to maximize in the worst scenario, namely, to maximize the smallest possible expected utility of the total terminal income over all possible realizations of the intractable claim. While this formulation is inspired by [7], our model loses the mean–variance structure so the approach there largely fails to work here.

At a first glance, the robust model still involves the expected utility of two random payoffs whose correlation is unknown so the aforementioned difficulty seems to be intact. However, we prove that at the *minimum* expected utility the two random incomes can be separated through the sum of the (marginal) quantile functions of the two; hence the robust measure is computable based on the *respective* information about the market and the claim.

The idea of taking the quantile functions, instead of random variables, as the decision variable to study preference functionals goes back to at least Schied [20]. Jin and Zhou [10] employ the quantile method to overcome the difficulty arising from probability distortion in a continuous-time behavioral portfolio selection model featuring cumulative prospect

theory. He and Zhou [6] put forward a unified theory on this approach that can be used to solve both expected and non-expected utility models and coin the term “quantile formulation”. The theory has been further developed since then; see e.g. [30, 26, 5, 27, 7, 25, 28, 29].

In this paper, we apply the quantile formulation to tackle our problem. From the “first-order condition” of the corresponding calculus of variation for quantile, we derive an ordinary integro–differential equation (OIDE) as well as a variational inequality type of ordinary differential equations (ODEs) that are satisfied by the quantile function of the optimal terminal payoff from the stock market. Surprisingly, when the utility is exponential, we show that our robust model can be reduced to a behavioral rank–dependent utility maximization problem whose solution is available ([25]).

The rest of the paper is organized as follows. In Section 2, we introduce the financial market and formulate the robust expected utility maximization (EUM) problem. We turn the problem into its quantile formulation and study its well-posedness in Section 3. Section 4 presents the solution. In Section 5 we discuss the special case of the exponential utility. Finally, Section 6 concludes.

2 Problem Formulation

Throughout this paper, we fix a probability space $(\Omega, \mathbf{F}, \mathbb{P})$ satisfying the usual assumptions, along with a standard m -dimensional Brownian motion

$$W = \{(W_1(t), \dots, W_m(t))^\top, t \geq 0\}$$

representing the randomness of the financial market under consideration (to be described below), where (and hereafter) A^\top denotes the transpose of a matrix or vector A . Unless otherwise stated, a r.v. is the shorthand for an \mathbf{F} -measurable random variable. We also fix an investment horizon $[0, T]$ where $T > 0$. Let $\{\mathcal{F}_t\}_{t \geq 0}$ be the filtration generated by W complemented by all the \mathbb{P} -null sets, and $L_{\mathcal{F}_T}^0$ be the set of \mathcal{F}_T -measurable random variables. We stress that $\mathcal{F}_T \subsetneq \mathbf{F}$; so there is randomness *outside* of the financial market or, equivalently, not every random variable belongs to $L_{\mathcal{F}_T}^0$.

For any r.v. Y , let F_Y denote its probability distribution function, and Q_F its quantile (function) defined by

$$Q_Y(t) := \inf \left\{ z \in \mathbb{R} \mid F_Y(z) > t \right\}, \quad t \in (0, 1),$$

which is the right-continuous inverse function of F_Y . It is easy to check that every quantile is a right-continuous and increasing (RCI) function on $(0, 1)$.¹ On the other hand, every

¹In this paper by “increasing” we mean “non-decreasing”, and by “decreasing” we mean “non-increasing”.

RCI function Q on $(0, 1)$ is the quantile for the r.v. $Q(U)$, where U is any r.v. that is uniformly distributed on $(0, 1)$. Hence, the set of quantiles is the same as that of RCI functions on $(0, 1)$. We may also use the convention that

$$Q_Y(0) := \lim_{t \rightarrow 0+} Q_Y(t) \quad \text{and} \quad Q_Y(1) := \lim_{t \rightarrow 1-} Q_Y(t);$$

so

$$Q_Y(0) = \text{ess inf } Y \quad \text{and} \quad Q_Y(1) = \text{ess sup } Y.$$

We write $X \sim Y$ if the r.v.'s X and Y follow the same distribution.

The derivative of any absolutely continuous function in this paper is chosen as the right-continuous version. Let $C_{\text{RCI}}^{1-}((0, 1))$ be the set of functions $f : (0, 1) \rightarrow \mathbb{R}$ that are absolutely continuous with derivatives f' being RCI. Clearly, any function $f \in C_{\text{RCI}}^{1-}((0, 1))$ is convex; hence the limits $f(0+) = \lim_{t \rightarrow 0+} f(t)$ and $f(1-) = \lim_{t \rightarrow 1-} f(t)$ exist (but may be infinity).

In what follows, “almost everywhere” (a.e.) and “almost surely” (a.s.) may be suppressed for expositional simplicity whenever no confusion might occur. Finally, we use $|Z|$ to denote $\sqrt{Z^\top Z}$ for any (column) vector Z .

Now we are ready to introduce the financial market and our investment problem.

2.1 Market Model

Consider a continuous-time arbitrage-free financial market where $m + 1$ assets are traded continuously on $[0, T]$. One of the assets is a bond, whose price $S_0(\cdot)$ evolves according to an ordinary differential equation (ODE):

$$\begin{cases} dS_0(t) = r(t)S_0(t) dt, & t \in [0, T], \\ S_0(0) = s_0 > 0, \end{cases}$$

where $r(t)$ is the appreciation rate of the bond at time t . The remaining m assets are stocks, and their prices are modeled by a system of stochastic differential equations (SDEs):

$$\begin{cases} dS_i(t) = S_i(t)[\beta_i(t) dt + \sum_{j=1}^m \sigma_{ij}(t) dW_j(t)], & t \in [0, T], \\ S_i(0) = s_i > 0, \end{cases}$$

where $\beta_i(t)$ is the appreciation rate of the stock i and $\sigma_{ij}(t)$ is the volatility coefficient at time t . Denote by $\beta := \{(\beta_1(t), \dots, \beta_m(t))^\top, t \geq 0\}$ the appreciation rate vector process and by $\sigma := \{(\sigma_{ij}(t))_{m \times m}, t \geq 0\}$ the volatility matrix process, along with the excess return rate vector process $B := \{B(t), t \geq 0\}$ where

$$B(t) := \beta(t) - r(t)\mathbf{1}, \quad t \in [0, T],$$

with $\mathbf{1} = (1, 1, \dots, 1)^\top$ denoting the m -dimensional unit vector.

We impose the following standing assumptions on the market parameters in this paper:

- The processes r , β , B , and σ are all uniformly bounded and $\{\mathcal{F}_t\}_{t \geq 0}$ -progressively measurable on $[0, T]$;
- The process B is not identical to zero; and
- The process σ is non-singular and its inverse process σ^{-1} is uniformly bounded on $[0, T]$.

As is well known, the market is complete under the above assumptions. Define the market price of risk process $\theta := \{\theta(t), t \geq 0\}$ where

$$\theta(t) := \sigma(t)^{-1} B(t), \quad t \in [0, T].$$

Clearly, it is a uniformly bounded and not identical to zero process on $[0, T]$.

2.2 Investment Problem

There is a small investor (“she”) in the market whose transactions have no influence on the asset prices in the market. She has an initial endowment $x > 0$ and invests in the financial market over the time period $[0, T]$. Denote by $\pi_i(t)$ the total market value of her wealth invested in stock i at time t , $i = 1, \dots, m$. We allow short selling so that $\pi_i(t)$ can take negative values. We also assume that the trading of shares takes place continuously in a self-financing fashion (i.e., there is no consumption or income) and the market is frictionless (i.e., transactions do not incur any fees or costs). The investor’s portfolio (process) is $\pi = \{(\pi_1(\cdot), \dots, \pi_m(\cdot))^T, t \geq 0\}$, and the corresponding wealth process X^π evolves according to the wealth SDE (see, e.g., Karatazas and Shreve [12]):

$$dX^\pi(t) = [r(t)X^\pi(t) + \pi(t)^T B(t)] dt + \pi(t)^T \sigma(t) dW(t), \quad t \in [0, T]. \quad (2.1)$$

In our model, the investor makes investment decisions based on the information $\{\mathcal{F}_t\}_{t \geq 0}$ from the financial market, which are strictly less than those represented by \mathbf{F} . So unlike many existing studies in portfolio selection, ours is a partial information model. Accordingly, we call π an admissible portfolio if it is $\{\mathcal{F}_t\}_{t \geq 0}$ -progressively measurable satisfying

$$\int_0^T |\pi(t)|^2 dt < \infty, \quad \text{a.s.}$$

and that the corresponding wealth process X^π is nonnegative. For any admissible portfolio π , the SDE (2.1) admits a unique solution X^π which is called an admissible wealth process, and (X^π, π) is called an admissible pair. Note that (2.1) is linear in X^π and π ; so all the admissible pairs form a convex set. Moreover, all the admissible pairs are $\{\mathcal{F}_t\}_{t \geq 0}$ -progressively measurable.

The original, classical expected utility maximization (EUM) problem is formulated as

$$\sup_{\pi} \mathbb{E}[u(X^{\pi}(T))], \quad (2.2)$$

subject to (X^{π}, π) being an admissible pair with $X^{\pi}(0) = x > 0$,

where u is a given utility function.

In our model, however, in addition to the investment payoff from the financial market, the investor will also receive the value of a claim ϑ (for instance, an excise of her company's stock option, a bonus, the down payment of a mortgage or even a lottery prize) at the maturity date T . She knows nothing about this claim – hence an *intractable* claim – except its *own* distribution before T . In particular, she does not know the *joint* distribution between this claim and the market, nor can she estimate this joint distribution purely based on the market data. Moreover, ϑ is \mathbf{F} -measurable, not \mathcal{F}_T -measurable; so it is not a hedgable contingent claim in the financial market. As a result, it is not meaningful to consider the utility of the total terminal wealth

$$\mathbb{E}[u(X^{\pi}(T) + \vartheta)],$$

because to determine this value the investor needs to know the joint distribution of $X^{\pi}(T)$ and ϑ . Inspired by Hou and Xu [7], we consider instead the worst scenario over all the possible realizations of ϑ , leading to the robust preference

$$J_0(X) := \inf_{Y \sim \vartheta} \mathbb{E}[u(X + Y)], \quad (2.3)$$

where X is the payoff from the financial market, and the infimum is taken over all \mathbf{F} -measurable r.v.'s Y that are distributed the same as ϑ . It turns out, as will be shown later, J_0 does not rely on the joint distribution between X and ϑ .

To summarize, we have the following investment model:

$$V_0(x) := \sup_{\pi} \inf_{Y \sim \vartheta} \mathbb{E}[u(X^{\pi}(T) + Y)], \quad (2.4)$$

subject to (X^{π}, π) being an admissible pair with $X^{\pi}(0) = x > 0$.

The problem can be regarded as a game between the investor (who chooses the best portfolio π and its output X^{π} in the financial market) and the nature (who chooses Y , the worst realization of ϑ , part of which may come from outside of the financial market). In contrast to many classical game models, we can not switch the order of sup and inf in the problem (2.4). It is not a classical control problem either, as we will show in the subsequent analysis that the objective functional is not the standard expectation in stochastic control theory (see, e.g., Yong and Zhou [32]). On the other hand, the classical EUM problem (2.2) can be regarded as the special case of (2.4) where $\vartheta \equiv 0$, in which case the optimal value is denoted by $V_{\text{EU}}(x)$.

To ensure the problem (2.4) to be well-defined and to avoid undue technicalities, we make the following assumption throughout this paper:

Assumption 2.1. *The claim ϑ is almost surely lower bounded, that is,*

$$\text{ess inf } \vartheta > -\infty.$$

Moreover, the utility function u is continuous on $[\min\{0, \text{ess inf } \vartheta\}, \infty)$ and continuously differentiable on $(\min\{0, \text{ess inf } \vartheta\}, \infty)$ with u' being strictly decreasing to 0 at infinity. Furthermore,

$$\mathbb{E}[u(\vartheta)] < \infty.$$

Remark 2.1. *The lower boundedness assumption on ϑ is very mild and reasonable, satisfied in most practical cases. The last assumption $\mathbb{E}[u(\vartheta)] < \infty$ is fulfilled when ϑ has a finite expectation, thanks to the concavity of u .*

Under Assumption 2.1, the utility function u is strictly increasing and strictly concave on $[\min\{0, \text{ess inf } \vartheta\}, \infty)$, under which we will show that the problem (2.4) is well-posed, namely, it has a finite optimal value (see Theorem 3.3 below).

3 Quantile Formulation

In this section we turn the problem (2.4) into a static optimization problem and present its quantile formulation.

3.1 A static optimization problem

Employing the well-established martingale method, we can divide solving the problem (2.4) into two steps. In the first step, we find the optimal solution X^* to the following static optimization problem

$$\begin{aligned} & \sup_X J_0(X), \\ & \text{subject to } X \in \mathcal{A}_x, \end{aligned} \tag{3.1}$$

where \mathcal{A}_x denotes the set of all possible terminal wealth at time T with an initial endowment x . The second step is to find an admissible portfolio π^* that replicates X^* , which will be the optimal portfolio to the original dynamic problem (2.4). As the market is complete by assumption, the second step is standard and can be solved by, say, the backward stochastic differential equation (BSDE) theory. Hence, it suffices to study the problem (3.1), which we will now focus on.

To solve (3.1), we need to first characterize the set \mathcal{A}_x in a more tractable form. The following result is well known (see, e.g., Karatazas and Shreve [12]).

Lemma 3.1. *The set \mathcal{A}_x can be expressed as*

$$\mathcal{A}_x = \left\{ X \in L_{\mathcal{F}_T}^0 : \mathbb{E}[\rho X] \leq x, X \geq 0 \right\}, \quad (3.2)$$

where ρ is called the pricing kernel or the stochastic discount factor, given by

$$\rho := \exp \left(- \int_0^T \left(r(s) + \frac{1}{2} |\theta(s)|^2 \right) ds + \int_0^T \theta(s)^\top dW(s) \right).$$

The constraint $\mathbb{E}[\rho X] \leq x$ in (3.2) is called the budget constraint.

It follows from our standing assumptions that θ is not identical to zero; hence ρ is a genuine random variable (i.e. it is not deterministic). We assume

Assumption 3.1. *The pricing kernel ρ has a finite expectation $\mathbb{E}[\rho] < \infty$.*

This again is a very mild assumption; in particular it is true when θ is a deterministic process.

Let \mathcal{U}_ρ be the set of r.v.'s that are comonotonic with ρ and uniformly distributed on $(0, 1)$. This set is non-empty and $\rho = F_\rho(U)$ for any $U \in \mathcal{U}_\rho$ (see, e.g. Xu [26]). In particular, \mathcal{U}_ρ is a singleton if and only if F_ρ is a continuous function, in which case $\mathcal{U}_\rho = \{F_\rho(\rho)\}$. This happens when, say, both r and θ are deterministic processes.

By virtue of Lemma 3.1, the problem (3.1) is equivalent to

$$\begin{aligned} & \sup_{X \in L_{\mathcal{F}_T}^0} J_0(X), \\ & \text{subject to} \quad \mathbb{E}[\rho X] \leq x, X \geq 0. \end{aligned} \quad (3.3)$$

We employ the quantile formulation method to tackle the problem (3.3), whose key idea is to change the decision variable from X to its quantile. Denote by Q_X the quantile function of a given r.v. X . The following lemma is critical in our quantile formulation.

Lemma 3.2. *We have*

$$J_0(X) = \int_0^1 u(Q_X(t) + Q_\vartheta(t)) dt. \quad (3.4)$$

As a consequence, the functional J_0 is law-invariant and strictly increasing in the sense that $J_0(X) = J_0(Y)$ whenever $X \sim Y$ and one of $J_0(X)$ and $J_0(Y)$ is finite, and $J_0(X_1) > J_0(X_2)$ whenever $X_1 \geq X_2$, $\mathbb{P}(X_1 > X_2) > 0$ and $J_0(X_2) < \infty$. In particular,

$$J_0(X + \varepsilon) > J_0(X)$$

for any constant $\varepsilon > 0$ if $J_0(X) < \infty$.

Proof. Recall that two r.v.'s X and Y are called comonotonic if

$$(X(\omega) - X(\omega'))(Y(\omega) - Y(\omega')) \geq 0 \quad d\mathbb{P} \times d\mathbb{P} \text{ almost surely,}$$

and called anti-comonotonic if X and $-Y$ are comonotonic (see, e.g. Schmeidler [21] and Yaari [31]). It is well known that comonotonic r.v.'s have the minimum convex order (see, e.g., Müller [15], Dhaene et al. [3, 4]);² so the infimum in (2.3) is obtained when X and Y^* are comonotonic with $Y^* \sim \vartheta$, in which case

$$Q_{X+Y^*} = Q_X + Q_{Y^*} \quad \text{a.e..}$$

Consequently,

$$J_0(X) = \int_0^1 u(Q_{X+Y^*}(t)) dt = \int_0^1 u(Q_X(t) + Q_{Y^*}(t)) dt = \int_0^1 u(Q_X(t) + Q_\vartheta(t)) dt.$$

So (3.4) holds.

Next, if $X_1 \geq X_2$ and $\mathbb{P}(X_1 > X_2) > 0$, then $Q_{X_1} \geq Q_{X_2}$, and $Q_{X_1}(t) > Q_{X_2}(t)$ for some $t \in (0, 1)$. By the right-continuity of quantiles, we have $Q_{X_1} > Q_{X_2}$ on some right neighborhood of t . Because u is strictly increasing and $J_0(X_2) < \infty$, we have

$$J_0(X_1) = \int_0^1 u(Q_{X_1}(t) + Q_\vartheta(t)) dt > \int_0^1 u(Q_{X_2}(t) + Q_\vartheta(t)) dt = J_0(X_2),$$

proving the strictly monotonicity. \square

Because J_0 is law-invariant, henceforth we will define, with a slight abuse of notation:

$$J_0(Q_X) := J_0(X) \equiv \int_0^1 u(Q_X(t) + Q_\vartheta(t)) dt.$$

By the expression (3.4), we see that the original problem (2.4) is not a standard stochastic control problem; hence classical stochastic control theory (e.g. Yong and Zhou [32]) is not applicable directly. To overcome this difficulty, we employ a refinement of the quantile formulation approach proposed in [28, 29] to tackle (2.4).

Let \mathcal{Q} denote the set of all quantiles generated by nonnegative r.v.'s, that is,

$$\mathcal{Q} := \{Q : (0, 1) \rightarrow \mathbb{R} \mid Q(\cdot) \text{ is the quantile for some nonnegative r.v. } X\}.$$

It is easy to see

$$\mathcal{Q} = \{Q : (0, 1) \rightarrow [0, \infty) \mid Q(\cdot) \text{ is RCI}\}.$$

Evidently, \mathcal{Q} is a convex set.

3.2 Well-posedness

Before tackling our problem (2.4), it is natural to ask whether the problem is well-posed; that is, whether its optimal value is finite. An infinite optimal value signals a

²For a pair of r.v.'s X and Y , we say that X is smaller than Y in the sense of convex order, if $\mathbb{E}[g(X)] \leq \mathbb{E}[g(Y)]$ for all convex functions g such that both expectations exist.

trivial optimization problem in which trade-offs between competing sub-objectives are not properly modeled. The following result answers the question of well-posedness completely.

Recall that V_{EU} and V_0 are the value functions of the problems (2.2) and (2.4), respectively.

Theorem 3.3. *The following statements are equivalent.*

1. $V_{\text{EU}}(x) < \infty$ for some $x > 0$.
2. $V_{\text{EU}}(x) < \infty$ for all $x > 0$.
3. $V_0(x) < \infty$ for all $x > 0$.
4. $V_0(x) < \infty$ for some $x > 0$.
5. There exists a constant $\lambda > 0$ such that

$$\mathbb{E}\left[u\left((u')^{-1}(\lambda\rho)\right)\right] < \infty, \quad \mathbb{E}\left[\rho(u')^{-1}(\lambda\rho)\right] < \infty.$$

Proof. The proof consists of the following steps: $1 \implies 2 \implies 3 \implies 4 \implies 1$, $1 \implies 5$ and $5 \implies 1$.

$1 \implies 2$: The proof is essentially from [9], which however is reproduced here for reader's convenience. Suppose that $V_{\text{EU}}(x) < \infty$ for some $x > 0$. Fix an arbitrary $y > 0$. Then $Y \in \mathcal{A}_y$ if and only if $X = \frac{x}{y}Y \in \mathcal{A}_x$ by (3.2). If $0 < y \leq x$, then, because u is increasing,

$$V_{\text{EU}}(y) = \sup_{Y \in \mathcal{A}_y} \mathbb{E}[u(Y)] = \sup_{X \in \mathcal{A}_x} \mathbb{E}\left[u\left(\frac{y}{x}X\right)\right] \leq \sup_{X \in \mathcal{A}_x} \mathbb{E}[u(X)] = V_{\text{EU}}(x) < \infty.$$

If $y > x$, then, because u is concave, we have for $k > 1$, $a > 0$:

$$\frac{u(ka) - u(a)}{ka - a} \leq \frac{u(a) - u(0)}{a - 0},$$

i.e.,

$$u(ka) \leq ku(a) - (k-1)u(0).$$

It hence follows

$$V_{\text{EU}}(y) = \sup_{X \in \mathcal{A}_x} \mathbb{E}\left[u\left(\frac{y}{x}X\right)\right] \leq \sup_{X \in \mathcal{A}_x} \mathbb{E}\left[\frac{y}{x}u(X) - \frac{y-x}{x}u(0)\right] = \frac{y}{x}V_{\text{EU}}(x) - \frac{y-x}{x}u(0) < \infty.$$

$2 \implies 3$: Because u is concave, we have for $a, b > 0$:

$$\frac{u(a+b) - u(a)}{b} \leq \frac{u(b) - u(0)}{b},$$

i.e.,

$$u(a+b) \leq u(a) + u(b) - u(0). \tag{3.5}$$

This together with the monotonicity of u yields

$$\begin{aligned}
J_0(X) &= \inf_{Y \sim \vartheta} \mathbb{E}[u(X + Y)] \\
&\leq \inf_{Y \sim \vartheta} \mathbb{E}[u(X + \max\{Y, 0\})] \\
&\leq \inf_{Y \sim \vartheta} \left(\mathbb{E}[u(X)] + \mathbb{E}[u(\max\{Y, 0\}) - u(0)] \right) \\
&= \mathbb{E}[u(X)] + \mathbb{E}[u(\max\{\vartheta, 0\}) - u(0)].
\end{aligned} \tag{3.6}$$

Maximizing both sides over $X \in \mathcal{A}_x$ yields

$$V_0(x) \leq V_{\text{EU}}(x) + \mathbb{E}[u(\max\{\vartheta, 0\}) - u(0)].$$

Thanks to Assumption 2.1, we have

$$\mathbb{E}[u(\max\{\vartheta, 0\}) - u(0)] = \mathbb{E}[u(\vartheta) - u(\min\{\vartheta, 0\})] \leq \mathbb{E}[u(\vartheta)] - u(\min\{\text{ess inf } \vartheta, 0\}) < \infty.$$

The implication thus follows.

$3 \implies 4$: This is trivial.

$4 \implies 1$: Suppose $Q_\vartheta(0) = \text{ess inf } \vartheta < 0$. By the concavity of u , for $x > y \geq 1$, we have

$$\frac{u(x) - u(y)}{x - y} \leq u'(1);$$

so

$$u(y) \geq u(x) + u'(1)(y - x).$$

This together with the monotonicity of u yields

$$\begin{aligned}
J_0(X) &= \inf_{Y \sim \vartheta} \mathbb{E}[u(X + Y)] \\
&\geq \inf_{Y \sim \vartheta} \mathbb{E}[u(X + \text{ess inf } Y)] = \mathbb{E}[u(X + Q_\vartheta(0))] \\
&\geq \mathbb{E}[u(X + Q_\vartheta(0)) \mathbf{1}_{X > 1 + |Q_\vartheta(0)|}] + \mathbb{E}[u(Q_\vartheta(0)) \mathbf{1}_{0 \leq X \leq 1 + |Q_\vartheta(0)|}] \\
&\geq \mathbb{E}[(u(X) + u'(1)Q_\vartheta(0)) \mathbf{1}_{X > 1 + |Q_\vartheta(0)|}] - |u(Q_\vartheta(0))| \\
&\geq \mathbb{E}[u(X) \mathbf{1}_{X > 1 + |Q_\vartheta(0)|}] - u'(1)|Q_\vartheta(0)| - |u(Q_\vartheta(0))| \\
&= \mathbb{E}[u(X)] - \mathbb{E}[u(X) \mathbf{1}_{0 \leq X \leq 1 + |Q_\vartheta(0)|}] - u'(1)|Q_\vartheta(0)| - |u(Q_\vartheta(0))| \\
&\geq \mathbb{E}[u(X)] - |u(1 + |Q_\vartheta(0)|)| - u'(1)|Q_\vartheta(0)| - |u(Q_\vartheta(0))|.
\end{aligned} \tag{3.7}$$

The last inequality also holds when $Q_\vartheta(0) = \text{ess inf } \vartheta \geq 0$, because in this case

$$J_0(X) = \inf_{Y \sim \vartheta} \mathbb{E}[u(X + Y)] \geq \mathbb{E}[u(X + Q_\vartheta(0))] \geq \mathbb{E}[u(X)].$$

Maximizing both sides of (3.7) over $X \in \mathcal{A}_x$ yields

$$V_0(x) \geq V_{\text{EU}}(x) - |u(1 + |Q_\vartheta(0)|)| - u'(1)|Q_\vartheta(0)| - |u(Q_\vartheta(0))|,$$

proving the desired implication.

1 \implies 5 : If $\mathbb{E}[\rho(u')^{-1}(\lambda\rho)] = \infty$ for all $\lambda > 0$, then by Jin, Xu and Zhou [9, Theorem 3.1], $V_{\text{EU}}(x) = \infty$ for all $x > 0$, contradicting Statement 1. So there exists some $\lambda > 0$ such that

$$x_\lambda := \mathbb{E}[\rho(u')^{-1}(\lambda\rho)] < \infty.$$

Then by the definition of V_{EU} , we have

$$\mathbb{E}[u((u')^{-1}(\lambda\rho))] \leq V_{\text{EU}}(x_\lambda).$$

But the right hand side (RHS) is finite by Statement 2, which has been proved to be a consequence of Statement 1. This proves the desired implication.

5 \implies 1 : Let $X_0 := (u')^{-1}(\lambda\rho)$ and $x_0 := \mathbb{E}[\rho X_0]$, then $\mathbb{E}[u(X_0)] < \infty$ and $x_0 < \infty$ by assumption. It thus follows that

$$\begin{aligned} V_{\text{EU}}(x_0) &= \sup_{X \in \mathcal{A}_{x_0}} \mathbb{E}[u(X)] \leq \sup_{X \in \mathcal{A}_{x_0}} \mathbb{E}[u(X) - \lambda\rho X] + \lambda x_0 \\ &\leq \mathbb{E}[u(X_0) - \lambda\rho X_0] + \lambda x_0 = \mathbb{E}[u(X_0)] < \infty, \end{aligned}$$

where the first equality is due to that $\mathbb{E}[\rho X] \leq x_0$ for any $X \in \mathcal{A}_{x_0}$, and the second equality is due to ω -wise maximization. The proof is complete. \square

From now on we assume that the value functions of the problems (2.2) and (2.4) are both finite, which is technically equivalent to that Statement 5 in Theorem 3.3 holds.

Lemma 3.4. *The value function V_0 to the problem (2.4) is concave and strictly increasing on $(0, \infty)$.*

Proof. It follows immediately from the concavity of u that J_0 is concave. The concavity of V_0 thus comes from the convexity of \mathcal{A}_x and concavity of J_0 .

Next, we show

$$\lim_{x \rightarrow \infty} V_0(x) = u(\infty).$$

In fact, $X = \frac{x}{\mathbb{E}[\rho]}$ is a feasible solution to the problem (3.3) with the initial endowment $x > 0$; so

$$\liminf_{x \rightarrow \infty} V_0(x) \geq \lim_{x \rightarrow \infty} J_0\left(\frac{x}{\mathbb{E}[\rho]}\right) = \lim_{x \rightarrow \infty} \int_0^1 u\left(\frac{x}{\mathbb{E}[\rho]} + Q_\vartheta(t)\right) dt \geq u(\infty)$$

by Fatou's lemma. For any feasible solution X to the problem (2.4), since u is increasing,

$$J_0(X) = \int_0^1 u(Q_X(t) + Q_\vartheta(t)) dt \leq u(\infty),$$

which shows that $V_0(x) \leq u(\infty)$. So we conclude $\lim_{x \rightarrow \infty} V_0(x) = u(\infty)$.

We now show that V_0 is strictly increasing. First, it follows from Lemma 3.2 that V_0 is increasing. Suppose V_0 was not strictly increasing, then it would be a constant

on some interval. Because V_0 is concave and finite at every point, V_0 must reach its (finite) maximum on that interval. Because V_0 is increasing and $\lim_{x \rightarrow \infty} V_0(x) = u(\infty)$, we conclude $V_0(x) = u(\infty)$ for some $x > 0$. This is impossible if $u(\infty) = \infty$ as $V_0(x) < \infty$. Hence $V_0(x) = u(\infty) < \infty$. By the monotone convergence theorem, there exists $t_0 \in (0, 1)$ such that

$$\int_{t_0}^1 Q_\rho(1-s) ds > \frac{1}{2} \int_0^1 Q_\rho(1-s) ds = \frac{1}{2} \mathbb{E}[\rho].$$

Let X be any feasible solution to the problem (3.3). Then it follows from the Hardy–Littlewood inequality and the non-negativity and monotonicity of quantiles that

$$\begin{aligned} x \geq \mathbb{E}[\rho X] &\geq \int_0^1 Q_X(s) Q_\rho(1-s) ds \geq \int_{t_0}^1 Q_X(s) Q_\rho(1-s) ds \\ &\geq Q_X(t_0) \int_{t_0}^1 Q_\rho(1-s) ds \geq \frac{1}{2} Q_X(t_0) \mathbb{E}[\rho], \end{aligned}$$

which yields

$$J_0(X) = \int_0^1 u(Q_X(s) + Q_\vartheta(s)) ds \leq \int_0^{t_0} u\left(\frac{2x}{\mathbb{E}[\rho]} + Q_\vartheta(s)\right) ds + \int_{t_0}^1 u(\infty) ds.$$

Maximizing both sides over $X \in \mathcal{A}_x$ leads to

$$V_0(x) \leq \int_0^{t_0} u\left(\frac{2x}{\mathbb{E}[\rho]} + Q_\vartheta(s)\right) ds + \int_{t_0}^1 u(\infty) ds < \int_0^{t_0} u(\infty) ds + \int_{t_0}^1 u(\infty) ds = u(\infty),$$

where the last inequality is due to the strictly monotonicity of u and the fact that $u(\infty) < \infty$. This contradicts that $V_0(x) = u(\infty)$. The proof is thus complete. \square

Thanks to the strictly monotonicity in Lemma 3.2, we can now apply [26, Theorem 9] to obtain the following result.

Proposition 3.5. *A random variable $X^* \in L_{\mathcal{F}_T}^0$ is an optimal solution to the problem (3.1) if and only if it can be expressed as*

$$X^* = \overline{Q}(1 - U)$$

where $U \in \mathcal{U}_\rho$ and \overline{Q} is an optimal solution to the following quantile optimization problem

$$\begin{aligned} &\sup_{Q \in \mathcal{Q}} J_0(Q) \\ &\text{subject to } \int_0^1 Q(t) Q_\rho(1-t) dt = x. \end{aligned} \tag{3.8}$$

Moreover, the optimal value of the problem (3.8) is equal to $V_0(x)$, the optimal value of the problem (2.4).

By virtue of this result, solving our original stochastic control problem (2.4) reduces to solving the quantile optimization problem (3.8).

4 Solution

We now solve (3.8), which is a constrained concave optimization (or calculus of variation) problem. The first step is to remove the constraint using the Lagrange method.

Lemma 4.1. *A quantile \overline{Q} is an optimal solution to (3.8) if and only if there exists a constant $\lambda > 0$ such that \overline{Q} satisfies*

$$\int_0^1 \overline{Q}(t) Q_\rho(1-t) dt = x, \quad (4.1)$$

and solves the following optimization problem

$$V_\lambda = \sup_{Q \in \mathcal{Q}} J_\lambda(Q), \quad (4.2)$$

where

$$J_\lambda(Q) := \int_0^1 \left[u(Q(t) + Q_\vartheta(t)) - \lambda Q(t) Q_\rho(1-t) \right] dt.$$

Moreover,

$$V_0(x) = V_\lambda + \lambda x.$$

Proof. \implies : Because V_0 is concave, for each given $x > 0$, we have

$$V_0(y) - \lambda y \leq V_0(x) - \lambda x, \quad y > 0, \quad (4.3)$$

where

$$\lambda = \liminf_{\Delta \rightarrow 0+} \frac{V_0(x + \Delta) - V_0(x)}{\Delta} \geq 0,$$

due to the monotonicity of V_0 . By the concavity and finiteness of V_0 ,

$$\lambda = \liminf_{\Delta \rightarrow 0+} \frac{V_0(x + \Delta) - V_0(x)}{\Delta} \leq \frac{V_0(x) - V_0(x/2)}{x - x/2} < \infty.$$

If $\lambda = 0$, then (4.3) would indicate that x is a global maximizer of V_0 , contradicting the strictly monotonicity of V_0 from Lemma 3.4. So we conclude $0 < \lambda < \infty$.

Suppose \overline{Q} is an optimal solution to (3.8). Then it clearly satisfies (4.1). Also,

$$V_0(x) = J_0(\overline{Q}) = J_\lambda(\overline{Q}) + \lambda \int_0^1 \overline{Q}(t) Q_\rho(1-t) dt = J_\lambda(\overline{Q}) + \lambda x. \quad (4.4)$$

For any $Q \in \mathcal{Q}$, let

$$Q_n(t) = \min\{Q(t), n\},$$

and

$$y_n = \int_0^1 Q_n(t) Q_\rho(1-t) dt \leq \int_0^1 n Q_\rho(1-t) dt = n \mathbb{E}[\rho] < \infty.$$

Then, by definition,

$$\begin{aligned} J_\lambda(Q_n) &= J_0(Q_n) - \lambda \int_0^1 Q_n(t) Q_\rho(1-t) dt = J_0(Q_n) - \lambda y_n \\ &\leq V_0(y_n) - \lambda y_n \leq V_0(x) - \lambda x = J_\lambda(\bar{Q}) \end{aligned}$$

in view of (4.3) and (4.4). It hence follows

$$J_\lambda(\bar{Q}) \geq J_\lambda(Q_n) \geq \int_0^1 u(Q_n(t) + Q_\vartheta(t)) dt - \lambda \int_0^1 Q(t) Q_\rho(1-t) dt.$$

Because

$$u(Q_n(t) + Q_\vartheta(t)) \geq u(Q_\vartheta(t)) \geq u(Q_\vartheta(0)),$$

Fatou's lemma yields

$$\begin{aligned} J_\lambda(\bar{Q}) &\geq \liminf_{n \rightarrow \infty} \int_0^1 u(Q_n(t) + Q_\vartheta(t)) dt - \lambda \int_0^1 Q(t) Q_\rho(1-t) dt \\ &\geq \int_0^1 u(Q(t) + Q_\vartheta(t)) dt - \lambda \int_0^1 Q(t) Q_\rho(1-t) dt = J_\lambda(Q). \end{aligned}$$

This indicates that \bar{Q} is an optimal solution to (4.2).

\Leftarrow : Suppose there exists $\lambda > 0$ such that \bar{Q} satisfies (4.1) and is an optimal solution to (4.2). Then, for any feasible solution $Q \in \mathcal{Q}$ to (3.8),

$$\begin{aligned} J_0(Q) &= J_\lambda(Q) + \lambda \int_0^1 Q(t) Q_\rho(1-t) dt = J_\lambda(Q) + \lambda x \\ &\leq J_\lambda(\bar{Q}) + \lambda x = J_\lambda(\bar{Q}) + \lambda \int_0^1 \bar{Q}(t) Q_\rho(1-t) dt = J_0(\bar{Q}). \end{aligned}$$

Because \bar{Q} is a feasible solution to (3.8), the above inequality shows that it is optimal to (3.8). \square

Our problem now reduces to first solving the unconstrained optimization (4.2), and then finding $\lambda > 0$ such that the resulting optimal solution satisfies (4.1). The quantile optimization (4.2) is a concave optimization problem, which can be tackled by calculus of variations following Xu [28, 29].

4.1 Well-posedness of (4.2)

Before solving (4.2), we need to first address its well-posedness.

Lemma 4.2. *The problem (4.2) is well-posed if and only if*

$$\mathbb{E}[u((u')^{-1}(\lambda\rho)) - \lambda\rho(u')^{-1}(\lambda\rho)] < \infty. \quad (4.5)$$

Proof. By the estimates (3.6) and (3.7), we have

$$-C_1 \leq J_0(X) - \mathbb{E}[u(X)] \leq C_2, \quad \forall X,$$

where

$$\begin{aligned} C_1 &:= |u(1 + |Q_\vartheta(0)|)| + u'(1)|Q_\vartheta(0)| + |u(Q_\vartheta(0))|, \\ C_2 &:= \mathbb{E}[u(\max\{\vartheta, 0\})] - u(0). \end{aligned}$$

Hence,

$$-C_1 \leq J_\lambda(Q) - \int_0^1 [u(Q(t)) - \lambda Q(t)Q_\rho(1-t)] dt \leq C_2, \quad \forall Q \in \mathcal{Q}.$$

Maximizing both sides over $Q \in \mathcal{Q}$ yields

$$-C_1 \leq V_\lambda - \sup_{Q \in \mathcal{Q}} \int_0^1 [u(Q(t)) - \lambda Q(t)Q_\rho(1-t)] dt \leq C_2. \quad (4.6)$$

Let

$$Q_0(t) := \lim_{s \rightarrow t+} (u')^{-1}(\lambda Q_\rho(1-s)).$$

Then $Q_0 \in \mathcal{Q}$. Applying the first-order condition to the maximum of the function $u(x) - \lambda x Q_\rho(1-t)$ in x , we have

$$u(Q(t)) - \lambda Q(t)Q_\rho(1-t) \leq u((u')^{-1}(\lambda Q_\rho(1-t))) - \lambda (u')^{-1}(\lambda Q_\rho(1-t))Q_\rho(1-t).$$

The RHS is, for a.e. $t \in (0, 1)$, equal to

$$u(Q_0(t)) - \lambda Q_0(t)Q_\rho(1-t).$$

Hence, we conclude that

$$\sup_{Q \in \mathcal{Q}} \int_0^1 [u(Q(t)) - \lambda Q(t)Q_\rho(1-t)] dt = \int_0^1 [u(Q_0(t)) - \lambda Q_0(t)Q_\rho(1-t)] dt.$$

For $U \in \mathcal{U}_\rho$, we have $Q_\rho(U) = \rho$ and

$$Q_0(1-U) = (u')^{-1}(\lambda Q_\rho(U)) = (u')^{-1}(\lambda \rho), \quad \text{a.s.},$$

so

$$\sup_{Q \in \mathcal{Q}} \int_0^1 [u(Q(t)) - \lambda Q(t)Q_\rho(1-t)] dt = \mathbb{E}[u((u')^{-1}(\lambda \rho)) - \lambda \rho (u')^{-1}(\lambda \rho)]. \quad (4.7)$$

This together with (4.6) implies that the optimal value of the problem (4.2) is finite if and only if the condition (4.5) holds. The proof is complete. \square

Remark 4.1. *Because*

$$\frac{\partial}{\partial \lambda} \mathbb{E}[u((u')^{-1}(\lambda \rho)) - \lambda \rho (u')^{-1}(\lambda \rho)] = -\mathbb{E}[\rho (u')^{-1}(\lambda \rho)] < 0,$$

the condition (4.5) holds if and only if either $\lambda > \underline{\lambda}_1$ or $\lambda \geq \underline{\lambda}_1$, where

$$\underline{\lambda}_1 = \inf \left\{ \lambda > 0 : \mathbb{E} \left[u \left((u')^{-1}(\lambda \rho) \right) - \lambda \rho (u')^{-1}(\lambda \rho) \right] < \infty \right\}.$$

4.2 Characterization of solution

We now investigate the problem (4.2). In view of Lemma 4.2, we only need to study the case under the well-posedness condition (4.5), which is henceforth assumed.

Because u is strictly concave, (4.2) admits at most one optimal solution. The following result characterizes the unique optimal solution, if it exists.

Lemma 4.3 (Optimality condition). *A quintile $\overline{Q} \in \mathcal{Q}$ is the unique optimal solution to (4.2) if and only if it satisfies*

$$\int_0^1 \left[u'(\overline{Q}(t) + Q_\vartheta(t)) - \lambda Q_\rho(1-t) \right] (Q(t) - \overline{Q}(t)) dt \leq 0, \quad \forall Q \in \mathcal{Q}. \quad (4.8)$$

Proof. Suppose \overline{Q} is the unique optimal solution to (4.2). For any $Q \in \mathcal{Q}$, $\varepsilon \in (0, 1)$, define

$$Q_\varepsilon(t) = \overline{Q}(t) + \varepsilon(Q(t) - \overline{Q}(t)), \quad t \in (0, 1).$$

Then $Q_\varepsilon \in \mathcal{Q}$. Recalling that \overline{Q} is optimal and applying Fatou's lemma, we get

$$\begin{aligned} 0 &\geq \liminf_{\varepsilon \rightarrow 0^+} \frac{1}{\varepsilon} \left[\int_0^1 u(Q_\varepsilon(t) + Q_\vartheta(t)) - \lambda Q_\varepsilon(t) Q_\rho(1-t) dt \right. \\ &\quad \left. - \int_0^1 u(\overline{Q}(t) + Q_\vartheta(t)) - \lambda \overline{Q}(t) Q_\rho(1-t) dt \right] \\ &\geq \int_0^1 \left[u'(\overline{Q}(t) + Q_\vartheta(t)) - \lambda Q_\rho(1-t) \right] (Q(t) - \overline{Q}(t)) dt, \end{aligned}$$

leading to (4.8).

Conversely, suppose $\overline{Q} \in \mathcal{Q}$ satisfies (4.8). Because u is concave, we have the elementary inequality $u(y) - u(x) \leq u'(x)(y - x)$ for any $x, y \in \mathbb{R}$. It follows that

$$\begin{aligned} &u(Q(t) + Q_\vartheta(t)) - \lambda Q(t) Q_\rho(1-t) - \left[u(\overline{Q}(t) + Q_\vartheta(t)) - \lambda \overline{Q}(t) Q_\rho(1-t) \right] \\ &\leq \left[u'(\overline{Q}(t) + Q_\vartheta(t)) - \lambda Q_\rho(1-t) \right] (Q(t) - \overline{Q}(t)), \quad \forall Q \in \mathcal{Q}. \end{aligned}$$

Integrating both sides in above and using (4.8), we conclude \overline{Q} is optimal to (4.2). \square

Before proceeding further, we impose the following technical assumption in the rest of this paper.

Assumption 4.1. *The quantiles of ϑ and ρ satisfy*

$$\lim_{t \rightarrow 0^+} \frac{u'(Q_\vartheta(t))}{Q_\rho(1-t)} = 0.$$

This assumption together with $\mathbb{E}[\rho] < \infty$ implies

$$\mathbb{E}[u'(\vartheta)] = \int_0^1 u'(Q_\vartheta(t)) dt < \infty. \quad (4.9)$$

Corollary 4.4. *The optimal solution \overline{Q} to the problem (4.2) must satisfy $\overline{Q}(0) = 0$.*

Proof. Because $\overline{Q} \geq 0$ and u is concave, Assumption 4.1 implies

$$u'(\overline{Q}(t) + Q_\vartheta(t)) - \lambda Q_\rho(1-t) \leq u'(Q_\vartheta(t)) - \lambda Q_\rho(1-t) < 0, \quad \text{for } t \in (0, \varepsilon),$$

with $\varepsilon > 0$ being sufficiently small. Suppose $\overline{Q}(0) > 0$, then we may assume $\overline{Q}(t) > 0$ for $t \in (0, \varepsilon)$ by the right-continuity of \overline{Q} . Now let $Q(t) = \overline{Q}(t) \mathbb{1}_{t \geq \varepsilon}$. Then $Q \in \mathcal{Q}$ and

$$\int_0^1 [u'(\overline{Q}(t) + Q_\vartheta(t)) - \lambda Q_\rho(1-t)] (Q(t) - \overline{Q}(t)) dt > 0,$$

which contradicts (4.8). The proof is complete. \square

It is hard to use the condition (4.8) to find the optimal solution to the problem (4.2) because one would have to compare \overline{Q} with all the other quantiles in \mathcal{Q} , a task as difficult as solving (4.2). Our next step is to find an equivalent condition to (4.8) that can be easily verified and utilized. To this end, let

$$H(t) := \int_t^1 [\lambda Q_\rho(1-s) - u'(\overline{Q}(s) + Q_\vartheta(s))] ds.$$

Using $\mathbb{E}[\rho] < \infty$, (4.9) and the monotonicity of u' , one can easily show that H is a continuous function on $[0, 1]$. In terms of H , the inequality in (4.8) now reads

$$\int_0^1 H'(t) (Q(t) - \overline{Q}(t)) dt \leq 0.$$

By taking $Q = 2\overline{Q}$ and $Q = \frac{1}{2}\overline{Q}$ in above, we obtain the following two conditions

$$\int_0^1 H'(t) \overline{Q}(t) dt = 0, \quad (4.10)$$

and

$$\int_0^1 H'(t) Q(t) dt \leq 0. \quad (4.11)$$

Suppose $H(a) < 0$ for some $a \in [0, 1)$. Let $Q(t) = \mathbb{1}_{t \in [a, 1]}$, then $Q \in \mathcal{Q}$ and

$$\int_0^1 H'(t) Q(t) dt = -H(a) > 0,$$

contradicting (4.11). Hence, $H \geq 0$ on $[0, 1]$ by continuity.

Suppose $H > 0$ on an interval $(a, b] \subset (0, 1)$ and $\overline{Q}(b) > \overline{Q}(a)$. Then since $H \geq 0$,

$H(1) = 0$ and by integration by parts, we have

$$\int_0^1 H'(t) \overline{Q}(t) dt = H(0) \overline{Q}(0) + \int_{(0,1)} H(t) d\overline{Q}(t) \geq \int_{(a,b]} H(t) d\overline{Q}(t) > 0,$$

contradicting (4.10). This implies that \overline{Q} is constant on every subinterval of $\{t \in (0, 1) \mid H(t) > 0\}$, and consequently $\overline{Q}' = 0$ on $\{t \in (0, 1) \mid H(t) > 0\}$.

Together with Corollary 4.4, we conclude that if \overline{Q} is the optimal solution to the problem (4.2), then

$$\begin{cases} \min \left\{ \overline{Q}'(t), \int_t^1 [\lambda Q_\rho(1-s) - u'(\overline{Q}(s) + Q_\vartheta(s))] ds \right\} = 0, & \text{for } t \in (0, 1); \\ \overline{Q}(0) = 0. \end{cases} \quad (4.12)$$

This is a so-called ordinary integro-differential equation (OIDE) with boundary condition for \overline{Q} . It is easy to check that (4.12) implies (4.10) and (4.11), which is equivalent to (4.8). Therefore, it follows from Lemma 4.3 that the solution to (4.12) is indeed the optimal solution to (4.2). Clearly, in comparison to (4.8), the condition (4.12) is easier to verify and use because it only depends on \overline{Q} itself.

Summarizing the preceding analysis we present the following main result.

Theorem 4.5 (Characterization of optimal solution I). *A function \overline{Q} is the unique optimal solution to the problem (4.2) if and only if it is an RCI function on $(0, 1)$ that satisfies the OIDE (4.12).*

Corollary 4.6. *If Q_ρ is continuous, then so is \overline{Q} , the optimal solution to the problem (4.2). Moreover, we have either $\overline{Q}'(t) = 0$ or*

$$\overline{Q}(t) = (u')^{-1}(\lambda Q_\rho(1-t)) - Q_\vartheta(t) \quad (4.13)$$

for every $t \in (0, 1)$. In particular, the latter case can only happen when Q_ϑ is continuous at t .

Proof. We have shown earlier that on each subinterval of $\{t \in (0, 1) \mid H(t) > 0\}$, \overline{Q} is a constant, and hence continuous there. Because quantiles are right-continuous, \overline{Q} is continuous at 0 as well.

Now suppose $H(t) = 0$ at some $t \in (0, 1)$. Because quantiles are RCI, u' and Q_ρ are continuous, we have

$$\begin{aligned} \lim_{\Delta \rightarrow 0+} \frac{H(t+\Delta) - H(t)}{\Delta} &= u'(\overline{Q}(t) + Q_\vartheta(t)) - \lambda Q_\rho(1-t), \\ \lim_{\Delta \rightarrow 0+} \frac{H(t) - H(t-\Delta)}{\Delta} &= u'(\overline{Q}(t-) + Q_\vartheta(t-)) - \lambda Q_\rho(1-t). \end{aligned}$$

Notice that t is a global minimizer of H , so the first limit is nonnegative and the second one is nonpositive, giving

$$u'(\overline{Q}(t) + Q_{\vartheta}(t)) - \lambda Q_{\rho}(1 - t) \geq u'(\overline{Q}(t-) + Q_{\vartheta}(t-)) - \lambda Q_{\rho}(1 - t).$$

Because u' is strictly decreasing, it follows that

$$\overline{Q}(t) + Q_{\vartheta}(t) \leq \overline{Q}(t-) + Q_{\vartheta}(t-).$$

But quantiles are increasing, the above should be an equation, which let us conclude \overline{Q} and Q_{ϑ} are both continuous at t . This further shows that H is differentiable at t , and

$$0 = H'(t) = u'(\overline{Q}(t) + Q_{\vartheta}(t)) - \lambda Q_{\rho}(1 - t).$$

Consequently,

$$\overline{Q}(t) = (u')^{-1}(\lambda Q_{\rho}(1 - t)) - Q_{\vartheta}(t),$$

completing the proof. \square

As much as OIDEs such as (4.12) are more accessible than the general condition (4.8), they are still hard to analyze in general. Next, we further simplify the condition by turning (4.12) into an ODE.

Denote

$$\Phi(t) = \int_t^1 u'(\overline{Q}(s) + Q_{\vartheta}(s)) \, ds. \quad (4.14)$$

Then $\Phi \in C_{\text{RCI}}^{1-}((0, 1))$ and

$$\overline{Q}(t) = (u')^{-1}(-\Phi'(t)) - Q_{\vartheta}(t), \quad H(t) = -\Phi(t) + \lambda \int_t^1 Q_{\rho}(1 - s) \, ds.$$

Noting that Φ'' exists almost everywhere because $\Phi \in C_{\text{RCI}}^{1-}((0, 1))$, we can rewrite the OIDE (4.12) as

$$\min \left\{ \frac{-\Phi''(t)}{u''((u')^{-1}(-\Phi'(t)))} - Q'_{\vartheta}(t), -\Phi(t) + \lambda \int_t^1 Q_{\rho}(1 - s) \, ds \right\} = 0, \quad \text{a.e. } t \in (0, 1). \quad (4.15)$$

Using the following simple fact that $\min\{a, b\} = 0$ if and only if $\min\{ak, b\ell\} = 0$ for any $k, \ell > 0$, together with the boundary conditions, we end up with the following equation

for Φ :

$$\begin{cases} \min \left\{ \Phi''(t) + Q'_\vartheta(t) u''((u')^{-1}(-\Phi'(t))), -\Phi(t) + \lambda \int_t^1 Q_\rho(1-s) ds \right\} = 0, & \text{a.e. } t \in (0, 1), \\ \Phi(1-) = 0, \quad \Phi'(0+) = -u'(Q_\vartheta(0)). \end{cases} \quad (4.16)$$

In (4.16), if Φ' or Q_ϑ is not differentiable at some point $t \in (0, 1)$, we convent

$$\Phi''(t) + Q'_\vartheta(t) u''((u')^{-1}(-\Phi'(t))) = +\infty,$$

in which case the equation reduces to

$$\Phi(t) = \lambda \int_t^1 Q_\rho(1-s) ds,$$

or $\Phi'(t) = -\lambda Q_\rho(1-t)$ whenever Q_ρ is continuous at t . Also note that it can happen that $u'(Q_\vartheta(0)) = +\infty$.

Equation (4.16) is a system of variational inequalities of ODEs, or a single-obstacle second-order ODE with mixed boundary conditions. The connection between (4.16) and the optimal solution to the problem (4.2) is given as follows.

Theorem 4.7 (Characterization of optimal solution II). *We have the following assertions.*

(1). *If \overline{Q} is the unique optimal solution to the problem (4.2), then*

$$\Phi(s) = \int_s^1 u'(\overline{Q}(t) + Q_\vartheta(t)) dt \quad (4.17)$$

is a solution to (4.16) in $C_{\text{RCI}}^{1-}((0, 1))$.

(2). *If Φ is a solution to (4.16) in $C_{\text{RCI}}^{1-}((0, 1))$, then*

$$\overline{Q}(t) = (u')^{-1}(-\Phi'(t)) - Q_\vartheta(t)$$

is the unique optimal solution to the problem (4.2).

As a consequence, (4.16) admits at most one solution in $C_{\text{RCI}}^{1-}((0, 1))$.

Proof. (1). Because \overline{Q} solves (4.2), we have (4.15). Consequently, we get the first equation in (4.16). The first boundary condition $\Phi(1-) = 0$ is evident. It follows from Corollary 4.4 that $\overline{Q}(0) = 0$, which implies the second boundary condition $\Phi'(0+) = -u'(Q_\vartheta(0))$. Moreover, as quantiles are RCI and u' is continuously decreasing, it is easy to verify that $\Phi \in C_{\text{RCI}}^{1-}((0, 1))$.

(2). Because Φ solves (4.16) in $C_{\text{RCI}}^{1-}((0, 1))$, it satisfies (4.15). Hence

$$\overline{Q}(t) = (u')^{-1}(-\Phi'(t)) - Q_\vartheta(t)$$

satisfies (4.12), which in turn is equivalent to (4.8). As $\Phi \in C_{\text{RCI}}^{1-}((0, 1))$ and Q_ϑ is RCI, \overline{Q} is right-continuous. By (4.12), \overline{Q} is also increasing; so \overline{Q} is a quantile. It follows then from Lemma 4.3 that \overline{Q} is the unique optimal solution to (4.2). \square

5 A Special Case of Exponential Utility

We have reduced the original problem to solving the equation (4.16). Such an equation can be in general solved numerically, but more concrete solutions are possible for special cases. In this section we study such a case where the utility function $u(x) = -e^{-\alpha x}$ for some $\alpha > 0$ and Q_ϑ is continuous. Then (4.16) becomes

$$\begin{cases} \min \left\{ \Phi''(t) + \alpha Q'_\vartheta(t) \Phi'(t), -\Phi(t) + \lambda \int_t^1 Q_\rho(1-s) ds \right\} = 0, & \text{a.e. } t \in (0, 1), \\ \Phi(1-) = 0, \quad \Phi'(0+) = -\alpha e^{-\alpha Q_\vartheta(0)}. \end{cases} \quad (5.1)$$

Note the second-order differential operator is now linear, which enables us to give a more explicit solution.

For any constant $c \geq 0$, consider the following integral equation for χ :

$$\chi(t) = c \int_t^1 e^{\alpha Q_\vartheta(\chi(s))} ds, \quad t \in [0, 1].$$

Clearly, any solution $\chi(t)$ is continuous in t . Let $\overline{\chi} = \chi/c$. Then

$$\overline{\chi}(t) = \int_t^1 e^{\alpha Q_\vartheta(c\overline{\chi}(s))} ds, \quad t \in [0, 1].$$

It follows from the monotone convergence theorem that $\overline{\chi}(t)$ is increasing in c ; hence $\chi(t)$ is strictly increasing in c for every $t \in [0, 1]$. Since Q_ϑ is continuous, $\overline{\chi}(t)$ and thus $\chi(t)$ are continuous in c for any fixed $t \in [0, 1]$. If $c = 0$, then $\chi(0) = 0$. If $c > e^{-\alpha Q_\vartheta(0)}$, then, since $\chi \geq 0$ and Q_ϑ is increasing,

$$\chi(0) = c \int_0^1 e^{\alpha Q_\vartheta(\chi(s))} ds \geq c \int_0^1 e^{\alpha Q_\vartheta(0)} ds > 1.$$

So there exists a constant $c > 0$ such that $\chi(0) = 1$, leading to the following equations

$$\chi'(t) = -ce^{\alpha Q_\vartheta(\chi(t))}, \quad \chi(0) = 1, \quad \chi(1) = 0.$$

Then

$$\chi''(t) = -ce^{\alpha Q_\vartheta(\chi(t))} \alpha Q'_\vartheta(\chi(t)) \chi'(t) = \alpha Q'_\vartheta(\chi(t)) (\chi'(t))^2, \quad \text{a.e. } t \in (0, 1).$$

Let $\Psi(t) = \Phi(\chi(t))$, then $\Psi'(t) = \Phi'(\chi(t)) \chi'(t)$ and

$$\Psi''(t) = \Phi''(\chi(t)) (\chi'(t))^2 + \Phi'(\chi(t)) \chi''(t) = \left(\Phi''(\chi(t)) + \alpha Q'_\vartheta(\chi(t)) \Phi'(\chi(t)) \right) (\chi'(t))^2.$$

Replacing t by $\chi(t)$ in (5.1) and using the above facts, we get

$$\begin{cases} \min \left\{ \Psi''(t), -\Psi(t) + \lambda \int_{\chi(t)}^1 Q_\rho(1-s) ds \right\} = 0, & \text{a.e. } t \in (0, 1), \\ \Psi(0+) = 0, \quad \Psi'(1-) = \Phi'(\chi(1-))\chi'(1-) = c\alpha. \end{cases} \quad (5.2)$$

This indicates that Ψ is the largest convex function dominated by the map

$$t \mapsto \lambda \int_{\chi(t)}^1 Q_\rho(1-s) ds$$

on $(0, 1)$ and satisfies the boundary conditions $\Psi(0+) = 0$ and $\Psi'(1-) = c\alpha$. This leads to an analytical expression of the solution to the problem (4.2): $\Phi(t) = \Psi(\chi^{-1}(t))$.

Because of the special form of the exponential utility function u , we can solve (4.2) in an alternative way. Indeed, rewrite (4.2) as

$$V_\lambda = \sup_{Q \in \mathcal{Q}} \int_0^1 \left[u(Q(t)) |u(Q_\vartheta(t))| - \lambda Q(t) Q_\rho(1-t) \right] dt.$$

Thanks to the triangle inequality and Assumption 2.1, we have

$$\begin{aligned} \mathbb{E}[|u(\vartheta)|] &\leq \mathbb{E}[|u(\vartheta) - u(\text{ess inf } \vartheta)|] + |u(\text{ess inf } \vartheta)| \\ &= \mathbb{E}[u(\vartheta) - u(\text{ess inf } \vartheta)] + |u(\text{ess inf } \vartheta)| < \infty. \end{aligned}$$

Also we may assume $\mathbb{E}[|u(\vartheta)|] > 0$; for otherwise ϑ would be a constant and (2.4) would reduce to the classical EUM problem (2.2) that has been solved in the literature. The above allows us to define

$$w(t) := \frac{1}{\mathbb{E}[|u(\vartheta)|]} \int_0^t |u(Q_\vartheta(1-s))| ds.$$

It is easy to check $w(0) = 0$, $w(1) = 1$ and w is differentiable and increasing; so w is a probability weighting (or distortion) function broadly involved in behavioral finance and insurance among others; see, e.g., Kahneman and Tversky's prospect theory [11], Quiggin's rank-dependent expected utility theory [18] and Wang's distortion insurance premium principle [22, 23, 24].

We now rewrite the problem (4.2) as

$$V_\lambda = \mathbb{E}[|u(\vartheta)|] \sup_{Q \in \mathcal{Q}} \int_0^1 \left[u(Q(t)) w'(1-t) - \lambda Q(t) \frac{Q_\rho(1-t)}{\mathbb{E}[|u(\vartheta)|]} \right] dt. \quad (5.3)$$

Intriguingly, the above is actually a quantile optimization problem arising from a behavioral finance model under the rank-dependent utility theory that has been studied and solved by Xia and Zhou [25] and Xu [27].

Theorem 5.1. *Let $u(x) = -e^{-\alpha x}$, $\alpha > 0$, and let $\delta(\cdot)$ be the concave envelope of the*

following function

$$t \mapsto - \int_0^{w^{-1}(1-t)} \frac{Q_\rho(s)}{\mathbb{E}[|u(\vartheta)|]} ds$$

on $[0, 1]$. Then

$$\overline{Q}(t) := (u')^{-1}(\lambda \delta'(t)), \quad t \in (0, 1),$$

is an optimal solution to (5.3) as well as to (4.2); and

$$X^* := (u')^{-1}(\lambda \delta'(1 - U))$$

is an optimal solution to the problem (3.1), where $U \in \mathcal{U}_\rho$ is arbitrary and $\lambda > 0$ satisfies

$$\int_0^1 (u')^{-1}(\lambda \delta'(t)) Q_\rho(1 - t) dt = x.$$

Indeed, we have the following more explicit expression for X^* without involving the Lagrange multiplier λ .

Corollary 5.2. *Let $U \in \mathcal{U}_\rho$ and*

$$X^* := \frac{1}{\mathbb{E}[\rho]} \left(x + \alpha^{-1} \int_0^1 \ln(\delta'(s)) Q_\rho(1 - s) ds \right) - \alpha^{-1} \ln(\delta'(1 - U)).$$

Then X^ is an optimal solution to (3.1) with the initial endowment $x > 0$.*

Proof. By Proposition 3.5, it is sufficient to prove that the optimal solution \overline{Q} to (5.3) given in Theorem 5.1 satisfies

$$\overline{Q}(t) = \frac{1}{\mathbb{E}[\rho]} \left(x + \alpha^{-1} \int_0^1 \ln(\delta'(s)) Q_\rho(1 - s) ds \right) - \alpha^{-1} \ln(\delta'(t)).$$

Recalling that $u(x) = -e^{-\alpha x}$, $u'(x) = \alpha e^{-\alpha x}$, we get

$$(u')^{-1}(x) = \alpha^{-1} \ln(\alpha) - \alpha^{-1} \ln(x), \quad x > 0;$$

so

$$\begin{aligned} x &= \int_0^1 (u')^{-1}(\lambda \delta'(s)) Q_\rho(1 - s) ds \\ &= \int_0^1 (\alpha^{-1} \ln(\alpha) - \alpha^{-1} \ln(\lambda) - \alpha^{-1} \ln(\delta'(s))) Q_\rho(1 - s) ds \\ &= (\alpha^{-1} \ln(\alpha) - \alpha^{-1} \ln(\lambda)) \mathbb{E}[\rho] - \alpha^{-1} \int_0^1 \ln(\delta'(s)) Q_\rho(1 - s) ds. \end{aligned}$$

Hence

$$\alpha^{-1} \ln(\alpha) - \alpha^{-1} \ln(\lambda) = \frac{1}{\mathbb{E}[\rho]} \left(x + \alpha^{-1} \int_0^1 \ln(\delta'(s)) Q_\rho(1 - s) ds \right).$$

Consequently,

$$\begin{aligned}\overline{Q}(t) &= (u')^{-1}(\lambda\delta'(t)) = \alpha^{-1}\ln(\alpha) - \alpha^{-1}\ln(\lambda) - \alpha^{-1}\ln(\delta'(t)) \\ &= \frac{1}{\mathbb{E}[\rho]} \left(x + \alpha^{-1} \int_0^1 \ln(\delta'(s)) Q_\rho(1-s) ds \right) - \alpha^{-1}\ln(\delta'(t)),\end{aligned}$$

as desired. \square

6 Concluding Remarks

This paper formulates and solves a robust EUM problem with an intractable claim by employing quantile formulation and calculus of variations, with the solution derived in terms of certain variational inequalities of ODEs. The result can be also used to *price* the intractable claim ϑ . Because ϑ is unpredictable and unhedgeable, we are unable to determine its (either upper or lower) hedging price the usual way. However, the results of this paper allow us to determine its price according to the famous utility indifference pricing principle; see e.g. [1, 2, 13, 14, 16, 17, 19].

Specifically, suppose the investor, with an initial endowment x , does not possess the intractable claim initially but is considering buying it. Then her EUM problem before purchasing ϑ is given by (2.2), with the optimal value $V_{\text{EU}}(x)$. If she pays p dollars to buy the claim, then her EUM problem after the purchase becomes (2.4) with x replaced by $x - p$, leading to the optimal value $V_0(x - p)$. The utility indifference pricing theory stipulates that p should be such that the investor is “indifferent” between buying and not buying ϑ , namely, p satisfies

$$V_{\text{EU}}(x) = V_0(x - p). \quad (6.1)$$

Notice

$$\lim_{p \rightarrow -\infty} V_0(x - p) = u(\infty) > V_{\text{EU}}(x),$$

and

$$V_0(0) = \mathbb{E}[u(\vartheta)].$$

It follows from Lemma 3.4 that the function V_0 is continuous and strictly increasing; so (6.1) admit a unique solution $p \leq x$ if and only if $V_{\text{EU}}(x) \geq \mathbb{E}[u(\vartheta)]$. Note the price is negative when the claim is negative.

Beyond the utility indifference pricing, our result can be possibly applied to price the intractable claim under other pricing principle; for instance, the marginal utility-based pricing in [8]. We leave the details to interested readers.

References

- [1] J. CVITANIĆ, W. SCHACHERMAYER, H. WANG: Utility maximisation in incomplete markets with random endowment. *Finance and Stochastics*, 5(2), 259–272 (2001)
- [2] M.H.A. DAVIS: Option pricing in incomplete markets. *Mathematics of Derivative Securities*, 216–227. Cambridge University Press, Cambridge (1997)
- [3] J. DHAENE, M. DENUIT, M. J. GOOVAERTS, R. KAAS, D. VYNCKE: The Concept of Comonotonicity in Actuarial Science and Finance: Theory. *Insurance: Mathematics and Economics*, 31, 3–33 (2002A)
- [4] J. DHAENE, M. DENUIT, M. J. GOOVAERTS, R. KAAS, D. VYNCKE: The Concept of Comonotonicity in Actuarial Science and Finance: Applications. *Insurance: Mathematics and Economics*, 31, 133–161 (2002B)
- [5] X.D. HE, H.Q. JIN, X.Y. ZHOU: Dynamic portfolio choice when risk is measured by Weighted VaR. *Mathematics of Operations Research*, 40, 773–796 (2015)
- [6] X.D. HE, X.Y. ZHOU: Portfolio Choice via Quantiles. *Mathematical Finance*, 21, 203–231 (2011)
- [7] D. HOU, Z.Q. XU: A robust Markowitz mean–variance portfolio selection model with an intractable claim. *SIAM Journal on Financial Mathematics*, 7, 124–151 (2016)
- [8] J. HUGONNIER, D. KRAMKOV, W. SCHACHERMAYER: On utility based pricing of contingent claims in incomplete markets. *Mathematical Finance*, 15(2), 203–212 (2005)
- [9] H.Q. JIN, Z.Q. XU, X.Y. ZHOU: A convex stochastic optimization problem arising from portfolio selection. *Mathematical Finance*, 18, 171–184 (2008)
- [10] H.Q. JIN, X.Y. ZHOU: Behavioral portfolio selection in continuous time. *Mathematical Finance*, 18, 385–426 (2008)
- [11] D. KAHNEMAN, A. TVERSKY: Prospect theory: An analysis of decision under risk. *Econometrica*, 46, 171–185 (1979)
- [12] I. KARATZAS, S. E. SHREVE: *Methods of Mathematical Finance*, Springer-Verlag, New York (1998)
- [13] M. MUSIELA T. ZARIPHOPULOU: An example of indifference prices under exponential preferences. *Finance and Stochastics*, 8(2), 229–239 (2004)
- [14] M. MUSIELA T. ZARIPHOPULOU: A valuation algorithm for indifference prices in incomplete markets. *Finance and Stochastics*, 8(3), 399–414 (2004)

- [15] A. MÜLLER: Stop-loss order for portfolios of dependent risks. *Insurance: Mathematics and Economics*, 21, 219-223 (1997)
- [16] A. OBERMAN, T. ZARIPHOULOU: Pricing early exercise contracts in incomplete markets. *CMS*, 1(1), 75–107 (2003)
- [17] A. PORCHET , N. TOUZI, X .WARIN: Valuation of power plants by utility indifference and numerical computation. *Math Methods Oper Res*, 70(1), 47–75 (2009)
- [18] QUIGGIN: A theory of anticipated utility. *Journal of Economic and Behavioral Organization*, 3(4), 323-343 (1982)
- [19] R. ROUGE, N. EL KAROUÏ : Pricing via utility maximization and entropy. *Math Finance*, 10(2), 259–276 (2000)
- [20] A. SCHIED: On the Neyman-Pearson problem for law-invariant risk measures and robust utility functionals. *Ann Appl Probab*, 14, 1398–1423 (2004)
- [21] D.SCHMEIDLER: Integral representation without additivity. *Proceedings of the American Mathematical Society*, 97, 255-261 (1986)
- [22] S. WANG: Insurance pricing and increased limits rate making by proportional hazards transforms, *Insurance: Mathematics and Economics*, 17, 43-54 (1995)
- [23] S. WANG: Premium calculation by transforming the layer premium density, *ASTIN Bulletin*, 26, 71-92 (1996)
- [24] S. WANG, V. R. YOUNG, H. H. PANJER: Axiomatic Characterization of insurance prices, *Insurance: Mathematics and Economics*, 21, 173-183 (1997)
- [25] J.M. XIA, X.Y. ZHOU: Arrow-Debreu equilibria for rank-dependent utilities. *Mathematical Finance*, 26, 558-588 (2016)
- [26] Z.Q. XU: A Characterization of Comonotonicity and its Application in Quantile Formulation. *J. Math. Anal. Appl.*, 418, 612-625 (2014)
- [27] Z.Q. XU: A note on the quantile formulation. *Mathematical Finance*, 26(3), 589-601 (2016)
- [28] Z.Q. XU: Moral-hazard-free insurance contract design under rank-dependent utility theory. working paper, <https://arxiv.org/abs/1803.02546> (2021)
- [29] Z.Q. XU: Moral-hazard-free insurance: mean-variance premium principle and rank-dependent utility. *Scandinavian Actuarial Journal*, 2023(3), 269-289 (2023)

- [30] Z.Q. XU, X.Y.ZHOU: Optimal stopping under probability distortion. *Annals of Applied Probability*, 23, 251-282 (2013)
- [31] M. E. YAARI: The dual theory of choice under risk. *Econometrica*, 55, 95-115 (1987)
- [32] J.M. YONG, X.Y.ZHOU: *Stochastic Controls: Hamiltonian Systems and HJB Equations*, 43. Springer, New York (1999)