IF: **IN RESIDUAL PHASE OF SCHIZOPHRENIA,** CHECK HERE _____ AND GO TO ***PANIC SYNDROME,*** F. 26

***GENERALIZED ANXIETY DISORDER* (CURRENT ONLY)**

GENERALIZED ANXIETY DISORDER CRITERIA

In the last six months, have you been really nervous?

Do you worry a lot that bad things might happen?

IF YES: What kinds of things do you worry about? (How much do you worry about [EVENTS OR ACTIVITIES])?

During the last six months, would you say that you have worried (more days than not?)

A. Excessive anxiety and worry (apprehensive expectation), occurring more days than not **for at least six months**, about a number of events or activities (such as school performance)

	?	1	2	3
CHI				
PAR				
OTH				
BEST				

When you're worrying this way, do you find that you can't stop yourself?

B. The person finds it difficult to control the worry.

	?	1	2	3
CHI				
PAR				
OTH				
BEST				

Now I'm going to ask you some questions about other feelings in your body (symptoms) that often go along with being nervous.

Thinking about those times in the past six months when you're feeling nervous or worried...

C. The anxiety and worry are associated with at least **one** of the following **six symptoms** (with at least some symptoms present for more days than not **for the past six months)** (three symptoms for adults)**:**

...do you often feel restless, so you can't sit still?

...do you often feel hyped up or edgy?

(1) restlessness or feeling keyed up or on edge

	?	1	2	3
CHI				
PAR				
OTH				
BEST				

...do you often get tired easily?

(2) being easily fatigued

	?	1	2	3
CHI				
PAR				
OTH				
BEST				

In the past six months when you've felt nervous or worried...

...do you have trouble paying attention?

...does you mind go blank?

(3) difficulty concentrating or mind going blank

	?	1	2	3
CHI				
PAR				
OTH				
BEST				

...are you often grouchy or irritable?

(4) irritability

	?	1	2	3
CHI				
PAR				
OTH				
BEST				

...are your muscles often tense?

(5) muscle tension

	?	1	2	3
CHI				
PAR				
OTH				
BEST				

...do you often have trouble falling to sleep or staying asleep?

(6) sleep disturbance (difficulty falling or staying asleep, or restless unsatisfying sleep)

	?	1	2	3
CHI				
PAR				
OTH				
BEST				

?	1	3

↑

AT LEAST ONE "C" SX IS CODED "3" (3 SXS FOR ADULTS)

GO TO ***Panic Syndrome*** F. 26

_____ _____

When was the first time that you started having (SXS OF GAD)?

Age at onset of Generalized Anxiety Syndrome (criteria A, B, C) (CODE 99 IF UNKNOWN)

?=inadequate information 1=absent or false 2=subthreshold 3=threshold or true

CODE BASED ON PREVIOUS INFORMATION.

IF NOT ALREADY KNOWN (Also ask parents): What effect has your/ your child's nervousness or worry or (PHYSICAL SYMPTOMS) had on your/ his or her life?

Just before you began having this anxiety, were you taking any drugs, stimulants or medicines?

(How much coffee, tea, or cola do you drink a day?)

Just before the fears began, were you sick (physically ill)? (What did the doctor say?)

IF A GENERAL MEDICAL CONDITION OR SUBSTANCE MAY BE ETIOLOGICALLY ASSOCIATED WITH THE ANXIETY, GO TO ***SUBSTANCE/ GMC,*** F.35. RETURN HERE TO MAKE RATING OF "1" OR "3."

D. The focus of the anxiety and worry is not confined to the features of another Axis I Disorder, e.g., being embarrassed in public (as in Social Phobia), being contaminated (as in Obsessive Compulsive Disorder), being away from home or close relatives (as in Separation Anxiety Disorder), gaining weight (as in Anorexia Nervosa), or having a serious illness (as in Hypochondriasis), and is not part of Posttraumatic Stress Disorder

? 1 3

↑

GO TO ***Panic Syndrome*** F. 26

E. The anxiety, worry, or physical symptoms cause clinically significant distress or impairment in school, social, occupational, or other important areas of functioning

? 1 3

↑

GO TO ***Panic Syndrome*** F. 26

F. Not due to the direct effects of a substance (e.g., drugs of abuse, medication) or to a general medical condition. DO NOT CODE 3 IF SYMPTOMS OCCUR EXCLUSIVELY DURING A MOOD OR PSYCHOTIC DISORDER, OR PDD

? 1 3

↑ ↑

↑ Due to Mood, Psychotic, or PDD or

Primary Anxiety Disorder

Due to substance use or a GMC

Etiological substances include intoxication with central nervous stimulants (e.g., cocaine, amphetamines, caffeine) or cannabis, hallucinogens, PCP, or alcohol, or withdrawal from central nervous system depressants (e.g., alcohol, sedatives, hypnotics) or cocaine.
Etiological general medical conditions include: hyper- and hypo-thyroidism, hypoglycemia, hyperparathyroidism, pheochromocytoma, congestive heart failure, arrhythmias, pulmonary embolism, chronic obstructive pulmonary disease, pneumonia, hyperventilation, B-12 deficiency, porphyria, CNS neoplasms, vestibular dysfunction, encephalitis.

CONTINUE

GENERALIZED ANXIETY CRITERIA A, B, C, D, E, AND F ARE CODED "3."

1 3

↑

Gen. Anxiety Disorder

CONTINUE